

# In vitro studies on antioxidant, hypolipidemic and cytotoxic potential of *Parmelia perlata*

Hafizur Rahman<sup>1</sup>, Vijaya B<sup>1</sup>, Soumya Ghosh<sup>2</sup>, Geetika Pant<sup>1</sup>, Sibi G<sup>1</sup>

<sup>1</sup>Department of Biotechnology, Indian Academy Degree College, Centre for Research and Post Graduate Studies, Bangalore, INDIA

<sup>2</sup>Department of Microbiology, Indian Academy Degree College, Centre for Research and Post Graduate Studies, Bangalore, INDIA

## Email address:

gsibii@gmail.com (Sibi G.)

## To cite this article:

Hafizur Rahman, Vijaya B, Soumya Ghosh, Geetika Pant, Sibi G. In Vitro Studies on Antioxidant, Hypolipidemic and Cytotoxic Potential of *Parmelia Perlata*. *American Journal of Life Sciences*. Special Issue: Recent Developments in Health Care through Plants and Microbes. Vol. 2, No. 6-1, 2014, pp. 7-10. doi: 10.11648/j.ajls.s.2014020601.12

**Abstract:** New drug preparations of natural origin are in need due to the numerous side effects and resistance development through the continuous and uncontrolled use of synthetic drugs. Lichens are reported to have manifold biological activities and in this study, antioxidant, hypolipidemic and cytotoxic potential of *Parmelia perlata* were assessed. Methanolic extract of *P. perlata* was prepared and used in DPPH assay, total phenolic content and total antioxidant potential assays. In vitro anti-cholesterol and cytotoxic assay using HCT 116 cell lines were performed. The results revealed that *P. perlata* contain high phenolic contents and antioxidant potential. Simvastatin was used as standard drug and the extracts demonstrated 48% anti-cholesterol activity. MTT assay using colon cancer cell lines (HCT 116) produced dose dependent cytotoxic effect with an IC<sub>50</sub> value of 202.1  $\mu\text{g ml}^{-1}$ .

**Keywords:** *Parmelia Perlata*, Antioxidant, Anticancer, Anti-Cholesterol, Colon Cancer

## 1. Introduction

Oxidative stress due to elevated levels of free radicals causes cellular and tissue damage that elicit direct or indirect damage to the body. Cancer is one of the main causes of death all over the world. The world health organization (WHO) estimates that 84 million people would die of cancer between 2005 and 2015 [1]. Hyperlipidemia is caused by a diet high in fat, especially saturated fat and cholesterol. Blood lipids are related to the development of coronary heart diseases. Synthetic antioxidants like butylated hydroxyanisole, butylated hydroxytoluene, tertbutylhydroquinone and propyl gallate have been reported to exert toxic effects [2]. New drug preparations of natural origin are in need due to the numerous side effects and resistance development through the continuous and uncontrolled use of synthetic drugs. Bioactive compounds from natural products have beneficial effects without causing undesired side effects. Among the natural origin, lichens are used in many research. Lichens are symbiotic associations between two entirely different types of microorganisms and produce secondary metabolites that have manifold biological activities [3,4].

Studies on the biological properties of lichens reveals the potential of lichens having antioxidant [5-8], antimicrobial [9-11] and cytotoxic [12, 13] activities.

*Parmelia perlata* (Huds.) Ach. (Parmeliaceae), commonly known as stone flower in India is thallus, foliaceous, membranous leaf like horizontally spreading lobes. The thallus is dirty white or grayish brown nearby 5-10 cm long. It has bitter or saline taste [14]. *Parmelia* is useful in treating a number of ailments [15, 16] and has also been reported to possess antimicrobial [17, 18], antiviral [19], antioxidant [20], antipyretic [21], anticancer [22] and antispasmodic [23] activities. In this study, *Parmelia perlata* was examined for its antioxidant, anti-cholesterol and anticancer activities to derive bioactive compounds from natural origin as great promise for biochemical applications and better human health.

## 2. Materials and Methods

### 2.1. Plant Material Collection and Extraction

Dried samples of *Parmelia perlata* were collected and identified by a Botanist, Bangalore University. The samples were ground into powder and sieved (60 mesh) and about

100 g of was extracted in methanol (1:10 w/v) at 25°C for 24 hours. The extract was centrifuged at 10,000 rpm for 10 minutes and the supernatant was filtered using Whatman No.1 filter paper and concentrated to dryness under reduced pressure in rotary vacuum evaporator. The final extract was stored in air tight containers at 4°C until used.

## 2.2. DPPH Assay

The capacity of the extract to scavenge the stable 2,2'-diphenyl-2-picrylhydrazyl (DPPH) free radical was determined according to the method reported by Gyamfi *et al.*, [24]. In brief, 50 µl of the methanolic plant extract (100 µg ml<sup>-1</sup>) was mixed with 1.8 ml of 0.5 mM DPPH in methanol solution. Methanol (50 µl) only was used as the experimental control. After 30 min of incubation at room temperature, the reduction in the number of DPPH free radicals was measured at 517 nm. The percent inhibition was calculated from the following equation:

$$\text{Inhibition (\%)} = \frac{(\text{Control}_{A517} - \text{Sample}_{A517})}{\text{Control}_{A517}} \times 100$$

## 2.3. Determination of Total Phenolic Content

The amount of total phenolics in the plant extract was determined with Folin-Ciocalteu (FC) reagent [25]. To 200 µl of sample (3 replicates), 1 ml of 1:2 dilution of FC reagent and 800 µl of sodium carbonate (7.5% w/v) were added and the resulting mixture was incubated at room temperature for 30 minutes. The absorbance of the sample was measured at 765 nm using a spectrophotometer and the results were expressed as milligram of gallic acid equivalent per gram of dry weight.

## 2.4. Antioxidant Potential Assay

The Antioxidant potential assay was carried out by phosphomolybdenum reduction assay [26]. To 200µl of plant extract, 1ml of the reagent containing 4 mM ammonium molybdate, 28 mM sodium phosphate and 0.6 M sulphuric acid were added and the resulting mixture was incubated at 37°C for 60 minutes. The absorbance of sample was measured at 695 nm using a spectrophotometer against blank using methanol. The antioxidant potential activity is expressed as mg of ascorbic acid equivalent per gram of dry weight.

## 2.5. In Vitro Anti-Cholesterol Assay

The anti-cholesterol assay was carried out as described by Iswantini *et al.*, [27] and Cholesterol Enzymatic Endpoint Method [28]. Cholesterol was dissolved in chloroform until achieving 25 mg/10 ml. 10 µl of the plant extract (two concentrations) were pipetted into micro titre plate followed by the addition of 2000 µl of Randox reagent and 10 µl of cholesterol as sample. 20 µl of distilled water and 2000 µl of Randox reagent were used as blank. Negative control comprised of 20 µL cholesterol and 2000 µl Randox reagent; standard comprised of 20 µl

simvastatin and 2000 µl Randox reagent. The contents were mixed and incubated for 10 minutes at room temperature. The absorbance was read at 500 nm in a microplate reader against reagent blank.

## 2.6. MTT Assay

The lichen extracts were tested for in vitro cytotoxicity using HCT-116 cells by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay [29]. HCT 116 (70-80%) confluent cell lines were trypsinized followed by viability checking and centrifugation. In a 96 well plate, 5 x 10<sup>4</sup> cells/ well were seeded and incubated for 24 hrs at 37°C in a humidified 5% CO<sub>2</sub> incubator. Plant extracts of varying concentrations ranging from 0-320 µg/ ml (two fold variations) in RPMI media without fetal bovine serum (FBS) and antibiotics were incubated for 24 hrs. After incubation with plant extracts, the media was removed from the wells and added 100µl/well (50 µg/well) of the MTT (5 mg/10 ml of MTT in 1x PBS) working solution was added and incubated for 3 to 4 hours. After incubation with MTT reagent, the media was removed from the wells and added 100 µl of DMSO to rapidly solubilize the formazan. The absorbance for each well was measured at 590 nm in a microtitre plate reader and the percentage inhibition was calculated using the formula.

$$\text{Inhibition (\%)} = \frac{(\text{Control}_{A590} - \text{Sample}_{A590})}{\text{Control}_{A590}} \times 100$$

## 3. Results




Fig 1. Standard curve for antioxidant potential using ascorbic acid




Fig 2. Standard curve for total polyphenols using gallic acid




Fig 3. MTT cytotoxic assay using HCT 116 cell lines

Methanolic extracts of *Parmelia perlata* was prepared by dissolving the concentrated extracts in 98% methanol and used for biological assays. DPPH assay of methanolic extract of *P. perlata* revealed that 32% inhibition thereby confirming the presence of antioxidants. Total antioxidant potential was carried out using ascorbic acid (Fig-1) and was determined as  $1.3 \text{ mg ml}^{-1}$ . The total phenolic content of *P. perlata* was determined spectrophotometrically by the Folin-Ciocalteu method using gallic acid as internal standard (Fig-2) and it was found as  $78.4 \text{ µg ml}^{-1}$ . In vitro anti-cholesterol activity of *P. perlata* was examined using simvastatin as the standard drug for a total half an hour at 10 minutes intervals. The purple colour developed during the reaction was decreased with increasing time and 48% activity was observed at the end of the reaction whereas 94% activity was found with the standard drug, simvastatin. Cytotoxic assay using colon cancer cell lines (HCT 116) revealed the antiproliferative activity of *P. perlata* against HCT 116 cell lines. Loss of viability of the dying cells as evidenced by the morphological changes was scrutinized by microscopy. Depending on the concentration, the lichen extract exhibited different levels of cytotoxicity like cell shrinkage, aggregation and cell death with an IC<sub>50</sub> value of  $202.1 \text{ µg ml}^{-1}$  (Fig-3).

## 4. Discussion

Antioxidants quenching the free radicals generated for specific metabolic requirement in the body. Oxidative damage of tissues and biomolecules occur when the levels of free radicals species exceed the level of antioxidant mechanism eventually leading to disease conditions [30]. Medicinal properties of plants have been investigated for their potential antioxidant activities to counteract metabolic disorders [31]. Plant extracts and plant-derived antioxidant compounds may potentiate the body's antioxidant defense mechanisms [32]. From results, it may be postulated that *Parmelia* was able to reduce the stable free radical DPPH to the yellow-colored diphenylpicrylhydrazine exhibiting free radical scavenging activity. Phenolic compounds have potential antioxidant capabilities and in this study a significant level of total phenols were determined from *P. perlata*.

Approximately 4.8 percent of men and women will be diagnosed with colon and rectum cancer at some point during their lifetime and colon cancer represents 8.2% of all new cancer cases [33]. Anticancer activities of *Parmelia* species against human melanoma, colon carcinoma and breast cancer cell lines were reported in earlier studies [34, 35]. In this study, HCT 116 cell lines were used to determine the antiproliferative activity of *P. perlata* by MTT assay. The results revealed the cytotoxic activity of the extract in a dose dependent manner. The effect of extract concentration was evidenced by microscopy and IC<sub>50</sub> value was found as  $202.1 \text{ µg ml}^{-1}$ . Plant extracts have exhibited anticancer activity against colon cancer in vitro and in vivo [36] and in this study in vitro cytotoxicity of *P. perlata* was determined.

## 5. Conclusion

This study revealed the presence of higher amounts of phenolic compounds in *P. perlata* and its antioxidant activities. The extract has exhibited significant cytotoxicity against HCT 116 and moderate levels of anti-cholesterol activity. In conclusion, *Parmelia* derived bioactive compounds hold great promise for biopharmaceutical applications as antioxidant and cytotoxic agents of new formulations for the benefit of human life.

## References

- [1] F. Danhier, O. Feron, V. Preat, "To exploit the tumor microenvironment: Passive and active tumor targeting of nanocarriers for anti-cancer drug delivery," *J. Control. Release.*, vol. 148, 35-146, 2010.
- [2] W.M. Zhang, B. Li, L. Han, H.D. Zhang, HD, "Antioxidant activities of extracts from areca (*Areca catechu* L.) flower, husk and seed," *Afr J Biotechnol.*, vol. 8: 3887-3892, 2009.
- [3] K. Molnar, E. Farkas, "Current results on biological activities of lichen secondary metabolites: a review," *Journal of Biosciences.*, vol. 65, 157-173, 2010.
- [4] T. Mitrovic, S. Stamenkovic, V. Cvetkovic, M. Nikolic, S. Tosic, D. Stojicic, "Lichens as source of versatile bioactive compounds," *Biologica Nyssana.*, vol. 2, 1-6, 2011.
- [5] B.C. Behera, N. Verma, A. Sonone, U. Makhija, "Determination of antioxidative potential of lichen *Usnea ghattensis* in vitro," *LWT-Food Sci Technol.*, vol. 39, 80-85, 2006.
- [6] M. Gulluce, A. Aslan, M. Sokmen, F. Sahin, A. Adiquzel, G. Agar, A. Sokmen, "Screening the antioxidant and antimicrobial properties of the lichens *Parmelia saxatilis*, *Platismatia glauca*, *Ramalina pollinaria*, *Ramalina polymorpha* and *Umbilicaria nylanderiana*," *Phytomedicine.*, vol. 13, 515-521, 2006.
- [7] H.D. Bhattarai, B. Paudel, S.G. Hong, H.K. Lee, J.H. Yim, "Thin layer chromatography analysis of antioxidant constituents of lichens from Antarctica," *J Nat Med.*, vol. 62, 481-484, 2008.

- [8] B. Paudel, H.D. Bhattarai, J.S. Lee, S.G. Hong, H.W. Shin, et al. "Antioxidant activity of polar lichens from King George Island (Antarctica)," *Polar Biol.*, vol. 31, 605-608, 2008.
- [9] B.R. Rankovic, M.M. Kosani, T.P. Stanojkovic, "Antioxidant, antimicrobial and anticancer activity of the lichens *Cladonia furcata*, *Lecanora atra* and *Lecanora muralis*," *BMC Complement Altern Med.*, vol. 11, 97, 2011.
- [10] A. Karagoz, N. Dogruoz, Z. Zeybek, A. Aslan, "Antibacterial activity of some lichen extracts," *J Med Plants Res.*, vol. 3, 1034-1039, 2009.
- [11] M. Kosanic, B. Rankovic, "Antibacterial and antifungal activity of different lichens extracts and lichen acid," *Res J Biotechnol.*, vol. 6, 23-26, 2011.
- [12] N.B. Perry, M.H. Benn, N.J. Brennan, E.J. Burgess, G. Ellis, D.J. Galloway, S.D. Lorimer, S. Tangney, "Antimicrobial, antiviral and cytotoxic activity of New Zealand lichens," *Lichenologist.*, vol. 31, 627-636, 1999.
- [13] C. Bezivin, S. Tomasi, F. Lohezic-Le Devehat, J. Boustie, J. "Cytotoxic activity of some lichen extracts on murine and human cancer cell lines," *Phytomedicine.*, vol. 10, 499-503, 2003.
- [14] K.R. Kiritkar, B.D. Basu, Indian Medicinal plants. 2<sup>nd</sup> Ed. M.B. Lalit Mohan Basu, Allahabad, vol. 3, 1956.
- [15] P. Halama, C.V. Haluwin, "Antifungal activity of lichen extracts and lichenic acids," *BioControl*, vol. 49, 95-107, 2004.
- [16] T. Tay, T.A. Ozdemir, M. Yilmaz, H. Turk, M. Kivanc, "Evaluation of the antimicrobial activity of the acetone extract of the Lichen *Ramalina farinacea* and its (+)-usnic acid, norstictic acid, and protocetraric acid constituents," *Journal of Biosciences*, vol. 59, 384-388, 2004.
- [17] B. Thippeswamy, N.R. Sushma, K.J. Naveenkumar, "Antimicrobial property of bioactive factor isolated from *Parmelia perlata*," *International Multidisciplinary Research Journal*, vol. 2, 1-5, 2012.
- [18] G. Sibi, V. Apsara, K. Dhananjaya, K.R. Ravikumar and H. Mallesha, "Phytochemical and antibacterial properties of spices against food borne bacteria with special reference to *Parmelia perlata*," *Global J. Bio-Sci. Biotechnol.*, vol. 2, 145-149, 2013.
- [19] C.O. Esimone, K.C. Ofokansi, M.U. Adikwu, E.C. Ibezim, D.O. Abonyi, G.N. Odaibo, D.O. Olaleye, "In vitro evaluation of the antiviral activity of extracts from the lichen *Parmelia perlata* (L.) Ach. against three RNA viruses," *J Infect Dev Ctries*, vol. 1, 315-320, 2007.
- [20] S.B. Patil, V.A. Ghadyale, S.S. Taklikar, C.R. Kulkarni, A.U. Arvindekar, "Insulin secretagogue, alpha-glucosidase and antioxidant activity of some selected spices in streptozotocin-induced diabetic rats," *Plant Foods Hum Nutr*, vol. 66, 85-90, 2011.
- [21] K. Muller, "Pharmaceutically relevant metabolites from lichens," *Appl. Microbiol Biotechnol.*, vol. 56, 9-16, 2001.
- [22] M.M. Kosanic, B.R. Rankovic, T.P. Stanojkovic, "Antioxidant, antimicrobial and anticancer activities of three *Parmelia* species," *J. Sci. Food Agric.*, vol. 92, 1909-1916, 2012.
- [23] J.A. Correia da Silva, "Action of *Parmelia caperata* extracts on smooth muscle organs," *Arch. Farmacol. Toxicol.*, vol. 2, 143-152, 1976.
- [24] M.A. Gyamfi, M. Yonamine, Y. Aniya, "Free radical scavenging action of medicinal herbs from Ghana *Thonningia sanguinea* on experimentally induced liver injuries," *Gen Pharmacol*, vol. 32, 661-667, 1999.
- [25] E.A. Ainsworth, K.M. Gillespie, Estimation of total phenolic content and other oxidation substrates in plant tissues using Folin-Ciocalteu reagent. *Nature Protocols*, vol. 2, 875- 877, 2007.
- [26] P. Prieto, M. Pineda, M. Aguilar M, "Spectrophotometric quantitation of antioxidant capacity through the formation of a Phosphomolybdenum Complex: Specific application to the determination of vitamin E," *Analytical Biochemistry*, vol. 269, 337-341, 1999.
- [27] D. Iswantini, D. Nurenda dan, P. Sugita, "Fractionation and characterization of active compounds from Bangle (*Zingiber Cassumunar* Roxb.) as an activator of the enzyme cholesterol oxidase," *Symposium Nasional Kimia Bahan Alam XV. Himpunan Kimia Bahan Alam. Bogor.*, 2005
- [28] Randox Laboratories, 2009. Cholesterol (Chol) enzymatic endpoint method manual. Available online.
- [29] S.P.M. Crouch, R. Kozlowski, K.J. Slater, J. Fletcher, "The use of ATP bioluminescence as a measure of cell proliferation and cytotoxicity". *J. Immunol. Methods* vol. 160, 81-88, 1993.
- [30] J.M.C. Gutteridge, "Free radicals in Disease Processes: A Compilation of cause and consequence," *Free Radic. Res. Comm.*, vol. 19, 141, 1995.
- [31] B. Auddy, F. Ferreira, L. Blasina, F. Lafon, F. Arredondo, F. Dajas, P.C. Tripathi, T. Seal, B. Mukherjee, "Screening of antioxidant activity of three Indian medicinal plants, traditionally used for the management of neurodegenerative diseases," *J Ethnopharmacol.*, vol. 84, 131-138, 2003.
- [32] B. Sagnia, D. Fedeli, R. Casetti, C. Montesano, G. Falcioni, V. Colizzi, "Antioxidant and Anti-inflammatory activities of extracts from *Cassia alata*, *Eleusine indica*, *Eremomastax speciosa*, *Carica papaya* and *Polyscias fulva* medicinal plants collected in Cameroon," *PLoS One*, vol. 9, e103999, 2014.
- [33] Cancer Statistics, <http://seer.cancer.gov/statfacts/html/colorect.html> accessed on Aug, 2014
- [34] N. Manojlovic, B. Rankovic, M. Kosanic, P. Vasiljevic, T. Stanojkovic, "Chemical composition of three *Parmelia* lichens and antioxidant, antimicrobial and cytotoxic activities of some their major metabolites," *Phytomedicine*, vol. 19, 1166-1172, 2012.
- [35] F. Ari, E. Ulukaya, S. Oran, S. Celikler, S. Ozturk, M.Z. Ozel, "Promising anticancer activity of a lichen, *Parmelia sulcata* Taylor, against breast cancer cell lines and genotoxic effect on human lymphocyte," *Cytotechnology*, 2014.
- [36] K.V. Balan, C. Demetozos, J. Prince, K. Dimas, M. Cladaras, Z. Han, J. H. Wyche, P. Pantazis, "Induction of apoptosis in human colon cancer HCT116 cells treated with an extract of the plant product, chios mastic gum," *In Vivo.*, vol. 19, 93-102, 2005.