

# Iridium (IV) Oxide ( $\text{IrO}_2$ ) Nanoparticles and Cancers

**Alireza Heidari<sup>1,2,3,4,\*</sup>, Margaret Hotz<sup>1,2,3</sup>, Nancy MacDonald<sup>1,2,3</sup>, Victoria Peterson<sup>1,2,3</sup>,  
Angela Caissutti<sup>1,2,3</sup>, Elizabeth Besana<sup>1,2,3</sup>, Jennifer Esposito<sup>1,2,3</sup>, Katrina Schmitt<sup>1,2,3</sup>,  
Ling-Yu Chan<sup>1,2,3</sup>, Francesca Sherwood<sup>1,2,3</sup>, Maria Henderson<sup>1,2,3</sup>, Jimmy Kimmel<sup>1,2,3</sup>**

<sup>1</sup>Faculty of Chemistry, California South University, Irvine, USA

<sup>2</sup>BioSpectroscopy Core Research Laboratory, California South University, Irvine, USA

<sup>3</sup>Cancer Research Institute (CRI), California South University, Irvine, USA

<sup>4</sup>American International Standards Institute, Irvine, USA

## Email address:

Scholar.Researcher.Scientist@gmail.com (A. Heidari), Alireza.Heidari@calsu.us (A. Heidari), Central@aisi-usa.org (A. Heidari)

\*Corresponding author

## To cite this article:

Alireza Heidari, Margaret Hotz, Nancy MacDonald, Victoria Peterson, Angela Caissutti, Elizabeth Besana, Jennifer Esposito, Katrina Schmitt, Ling-Yu Chan, Francesca Sherwood, Maria Henderson, Jimmy Kimmel. Iridium (IV) Oxide ( $\text{IrO}_2$ ) Nanoparticles and Cancers. *American Journal of Physical Chemistry*. Vol. 10, No. 4, 2021, pp. 54-58. doi: 10.11648/j.ajpc.20211004.11

**Received:** July 5, 2021; **Accepted:** August 9, 2021; **Published:** October 30, 2021

**Abstract:** In the current research, roles and applications of Iridium (IV) Oxide ( $\text{IrO}_2$ ) nanoparticles in cancer nanobiotechnology using synchrotron and synchrocyclotron radiations is investigated. The calculation of thickness and optical constants of Iridium (IV) Oxide ( $\text{IrO}_2$ ) roles and applications of Iridium (IV) Oxide ( $\text{IrO}_2$ ) nanoparticles in cancer nanobiotechnology using synchrotron and synchrocyclotron radiations produced using sol-gel method over glassy medium through a single reflection spectrum is presented. To obtain an appropriate fit for reflection spectrum, the classic Drude-Lorentz model for parametric di-electric function is used. The best fitting parameters are determined to simulate the reflection spectrum using Lovenberg-Marquardt optimization method. The simulated reflectivity from the derived optical constants and thickness are in good agreement with experimental results. The results of optimization algorithm of Lovenberg-Marquardt with physical model of Drude-Lorentz for determining optical constants of Iridium (IV) Oxide ( $\text{IrO}_2$ )-roles and applications of Iridium (IV) Oxide ( $\text{IrO}_2$ ) nanoparticles in cancer nanobiotechnology using synchrotron and synchrocyclotron radiations produced using sol-gel method through a single reflection spectrum show that higher doping leads to lower reflectivity and reflection coefficient and also, leads to increase in thickness of thin layer.

**Keywords:** Iridium (IV) Oxide ( $\text{IrO}_2$ ) Nanoparticles, Cancer Nanobiotechnology, Synchrotron and Synchrocyclotron Radiations

## 1. Introduction

Roles and applications of Iridium (IV) Oxide ( $\text{IrO}_2$ ) nanoparticles in cancer nanobiotechnology using synchrotron and synchrocyclotron radiations is investigated. Iridium (IV) Oxide ( $\text{IrO}_2$ ) is a semi-conductor of type n which its 3d level is filling up [1-6] and it belongs to a group of smart materials that reacts to variations of temperature, electrical or magnetic fields and pressure. This oxide can be used as thin films for a wide range of applications including electrical and or optical-thermal switching tools and energy storing covers [7-14]. Therefore, determining optical constants (refractive coefficient,

n, and extinction coefficient, k) of Iridium (IV) Oxide ( $\text{IrO}_2$ ) thin films is essential for designing optoelectrical and optical tools for producing optical covers and similar tools such as multilayer covers and filters [15-21]. The measured experimental parameters including optical reflectivity are used as a function of wavelength to determine optical parameters of thin layers [22-27]. For determining optical parameters, various physical models such as Kuschi, Frouhi-Blumber and Tawk-Lorentz have been suggested to calculate refractive coefficient, n, and extinction coefficient, k, for any thin layer, an appropriate optical model should be selected and used for estimation of real and imaginary di-electric function according to its physical condition [28-34]. To do this, an initial guess is

needed for parameters of di-electric function and thickness which is defined as a range regarding physical characteristics of thin film and the available results in the literature. Iridium (IV) Oxide (IrO<sub>2</sub>)– roles and applications of Iridium (IV) Oxide (IrO<sub>2</sub>) nanoparticles in cancer nanobiotechnology using synchrotron and synchrocyclotron radiations are produced over glassy medium in sol–gel laboratory, Faculty of Chemistry, BioSpectroscopy Core Research Laboratory and Cancer Research Institute (CRI) at California South University, Irvine, California, USA, under similar conditions. Measurement of thin films are performed on four samples of Iridium (IV) Oxide (IrO<sub>2</sub>) as roles and applications of Iridium (IV) Oxide (IrO<sub>2</sub>) nanoparticles in cancer nanobiotechnology using synchrotron and synchrocyclotron radiations with mole ratio of 0.5, 1 and 1.5% of Iridium (IV) Oxide (IrO<sub>2</sub>) [35–40]. Simulation of experimental spectra are performed using a single reflection spectrum of thin films and through Drude–Lorentz physical model in optimization process of Lovenberg–Marquardt. Optical constants such as reflection coefficient,  $n$ , extinction coefficient,  $k$ , and layer thickness are simultaneously determined at wavelength of 400–1100 (nm).

## 2. Modeling, Simulation and Calculation Method

A usual method for describing optical constants of thin films is utilizing classic dispersion relationships based on di-electric function. One of the oldest and most applicable dispersion relationships is Drude–Lorentz di-electric equation which is based on the interaction between light and material. This relationship is shown in Eq. (1):

$$\varepsilon = \varepsilon_{\infty} + \sum_{j=1}^n \frac{f_j E_{0j}^2}{E_{0j}^2 - E^2 + i\Gamma_j E} + \frac{E_p^2}{E^2 + iE_{\tau} E} \quad (1)$$

where  $\varepsilon_{\infty}$ ,  $f_j$ ,  $E_0$  and  $\Gamma_j$  are di-electric constant at high frequencies, resonance amplitude, power and resonance width–band which are recognized as the reason for damping. Damping is due to absorption process which includes transition between two states. The third term is related to Drude model.  $E_p$  is density of Plasma energy and  $E_r$  is incident energy [4]. The complex di-electric function as  $\varepsilon = \varepsilon_1 + i\varepsilon_2$  which describes the reaction of material with electromagnetic waves as a function of photon energy,  $E$ , or wavelength,  $\lambda$ , has a real part  $\varepsilon_1$  and an imaginary part  $\varepsilon_2$ . Real and imaginary parts of complex reflection coefficient, namely  $n(\lambda)$  and  $k(\lambda)$  are related to di-electric function as Eq. (2) [5]:

$$\begin{aligned} n(\lambda) &= \left( \frac{\varepsilon_1 + (\varepsilon_1^2 + \varepsilon_2^2)^{1/2}}{2} \right)^{1/2} \\ k(\lambda) &= \left( \frac{-\varepsilon_1 + (\varepsilon_1^2 + \varepsilon_2^2)^{1/2}}{2} \right)^{1/2} \end{aligned} \quad (2)$$

Reflection spectrum (R) of samples for normal incident is

a function of film thickness  $d$ , medium reflection coefficient  $S$ , incident light wavelength  $\lambda$ , reflection coefficient  $n(\lambda)$  and extinction coefficient  $k(\lambda)$ .


Simulation of the measured reflection data using optimization of objective function, which is the square of difference between the measured reflection spectrum and the calculated one, is defined as:

$$O = (\varepsilon_{\infty}, f, \Gamma, E_0, E_p, E_{\tau}, d) = \sum (R_{\text{meas}} - R_{\text{calc}})^2 \quad (3)$$

where,  $R_{\text{meas}}$  and  $R_{\text{calc}}$  are the measured and theoretical reflection spectrum, respectively. using the fitting parameters obtained from minimization of objective function, dispersion curves of reflection and extinction coefficients can be estimated.

## 3. Results and Discussion

The measured and simulated reflection spectra with fitting parameters of Iridium (IV) Oxide (IrO<sub>2</sub>)–roles and applications of Iridium (IV) Oxide (IrO<sub>2</sub>) nanoparticles in cancer nanobiotechnology using synchrotron and synchrocyclotron radiations at various concentrations of 0.5, 1 and 1.5%, named as a, b, and c, and roles and applications of Iridium (IV) Oxide (IrO<sub>2</sub>) nanoparticles in cancer nanobiotechnology using synchrotron and synchrocyclotron radiations sample, named as p, are shown in Figure 1 in wavelength range of 400–1100 (nm) (visible regions close to infrared) using Drude–Lorentz model for air, film, medium, air system.



**Figure 1.** Results of simulating the reflection spectrum for Iridium (IV) Oxide (IrO<sub>2</sub>)–roles and applications of Iridium (IV) Oxide (IrO<sub>2</sub>) nanoparticles in cancer nanobiotechnology using synchrotron and synchrocyclotron radiations at concentrations of (a) 0.5%, (b) 1%, (c) 1.5% and (p) non-doped.

Comparison of the results were shown that the sample containing 0.5% of Ir (sample a) has shown more reflectivity than samples containing 1% and 1.5% of Iridium (IV) Oxide (IrO<sub>2</sub>) (samples b and c). As can be seen in Figure 1, the reflection of thin films is decreased by increase in mole


concentration of Ir to Iridium (IV) Oxide ( $\text{IrO}_2$ ). This reduction can be attributed to various reasons such as increasing roughness, increasing thickness and increasing the concentration of contaminant. The results of investigation about surface roughness using AFM method confirms the increasing of roughness by increasing the concentration of Ir. Therefore, dispersion of incident light is increased in thin films. Variation of thickness of thin film by increasing the percentage of Ir is effective in variation of reflectivity of thin

films which is due to sol viscosity. Changing the crystalline structure and chemical composition of thin films induced by penetration of Ir ions into the crystalline lattice of Iridium (IV) Oxide ( $\text{IrO}_2$ ) is another effective factor which leads to changing the reflection spectrum. The results of structural analysis using XRD confirms the tendency to be amorphous by increasing the concentration of contaminant.


The best fitting parameters obtained from optimization process and experimental data fitting are listed in Table 1.

**Table 1.** Fitting parameters of di-electric function of DL model.

Parameter	Pure	% 0.5 Iridium (IV) Oxide ( $\text{IrO}_2$ )	1% Iridium (IV) Oxide ( $\text{IrO}_2$ )	% 1.5 Iridium (IV) Oxide ( $\text{IrO}_2$ )
$\epsilon_\infty$	0.85	0.75	0.65	0.55
$E_p$	9.85	8.75	7.65	6.55
$E_\tau$	5.45	5.4	5.35	5.25
$f$	3.35	3.25	3.15	3.05
$E_0$	5.5	5.4	5.3	5.2
$\Gamma$	4.5	4.4	4.3	4.2
$d(\text{nm})$	225	335	445	655



**Figure 2.** Reflection coefficient of Iridium (IV) Oxide ( $\text{IrO}_2$ ) thin films with Ir concentrations of (a) 0.5%, (b) 1%, (c) 1.5% and (p) pure sample.



**Figure 3.** Extinction coefficient of Iridium (IV) Oxide ( $\text{IrO}_2$ ) thin films with Ir concentrations of (a) 0.5%, (b) 1%, (c) 1.5% and (p) pure sample.

As can be seen in Table 1, more increase in Ir leads to increase in  $\Gamma$ ,  $f$ ,  $E_0$  and  $d$  and decrease in other parameters as crystalline structure and inter-atom distance changes in lattice of Iridium (IV) Oxide ( $\text{IrO}_2$ ) thin film. According to A. Heidari [7],  $E_0$  in the range of 2.9–3.1 (eV) shows optical

transition capacity band to displaced state of conducting band which according to the data of Table 1, it can be concluded that optical transition energy (gaff energy) increases with increase in Ir concentration. The calculation results of optical constants including reflection coefficient and extinction coefficient using the parameters of obtained di-electric function from the optimization process of thin films at various concentrations of Iridium (IV) Oxide ( $\text{IrO}_2$ ) as 0.5% (sample a), 1% (sample b) and 1.5% (sample c) are shown in Figures 2 and 3, respectively.

As can be seen in Figure 3, reflection coefficient of samples at 500–1100 (nm) are the same and are decreased by increasing wavelength. By increasing the concentration of Ir, reflection coefficient is totally reduced which is in good agreement with the results related to variations of reflectivity in Figure 2 in which, increasing roughness leads to increase in dispersion and hence, reducing the amount of reflection spectrum. It can be seen in Figure 3 that  $k(\lambda)$  for two samples of p and a are of increasing rate at wavelength range of 400–500 (nm). Further, all samples are of decreasing rate at the range of 500–800 (nm). Totally,  $k(\lambda)$  is reduced by increase in Ir concentration. In other words, optical absorption is reduced in this range and the emerged peaks at extinction coefficient are in agreement with parameters of Drude–Lorentz obtained from the optimization algorithm.

#### 4. Conclusions, Summary, Recommendations, Perspectives, Useful Suggestions and Future Studies

The results of optimization algorithm of Lovenberg–Marquardt with physical model of Drude–Lorentz for determining optical constants of Iridium (IV) Oxide ( $\text{IrO}_2$ )—roles and applications of Iridium (IV) Oxide ( $\text{IrO}_2$ ) nanoparticles in cancer nanobiotechnology using synchrotron and synchrocyclotron radiations produced using sol–gel method through a single reflection spectrum show that higher

doping leads to lower reflectivity and reflection coefficient and also, leads to increase in thickness of thin layer.

## Acknowledgements

This study was supported by the Cancer Research Institute (CRI) Project of Scientific Instrument and Equipment Development, the National Natural Science Foundation of the United States, the International Joint BioSpectroscopy Core Research Laboratory Program supported by the California South University (CSU), and the Key project supported by the American International Standards Institute (AISI), Irvine, California, USA.

## References

- [1] A. Heidari, C. Brown, "Study of Composition and Morphology of Cadmium Oxide (CdO) Nanoparticles for Eliminating Cancer Cells", *J Nanomed Res.*, Volume 2, Issue 5, 20 Pages, 2015.
- [2] A. Heidari, C. Brown, "Study of Surface Morphological, Phytochemical and Structural Characteristics of Rhodium (III) Oxide (Rh<sub>2</sub>O<sub>3</sub>) Nanoparticles", *International Journal of Pharmacology, Phytochemistry and Ethnomedicine*, Volume 1, Issue 1, Pages 15–19, 2015.
- [3] A. Heidari, "An Experimental Biospectroscopic Study on Seminal Plasma in Determination of Semen Quality for Evaluation of Male Infertility", *Int J Adv Technol* 7: e007, 2016.
- [4] A. Heidari, "Extraction and Preconcentration of N-Tolyl-Sulfonyl-Phosphoramid-Saeure-Dichlorid as an Anti-Cancer Drug from Plants: A Pharmacognosy Study", *J Pharmacogn Nat Prod* 2: e103, 2016.
- [5] A. Heidari, "A Thermodynamic Study on Hydration and Dehydration of DNA and RNA-Amphiphile Complexes", *J Bioeng Biomed Sci* 5: 006, 2016.
- [6] A. Heidari, "Computational Studies on Molecular Structures and Carbonyl and Ketene Groups' Effects of Singlet and Triplet Energies of Azidoketene O=C=CH-NNN and Isocyanatoketene O=C=CH-N=C=O", *J Appl Computat Math* 5: e142, 2016.
- [7] A. Heidari, "Study of Irradiations to Enhance the Induces the Dissociation of Hydrogen Bonds between Peptide Chains and Transition from Helix Structure to Random Coil Structure Using ATR-FTIR, Raman and <sup>1</sup>HNMR Spectroscopies", *J Biomol Res Ther* 5: e146, 2016.
- [8] A. Heidari, "Future Prospects of Point Fluorescence Spectroscopy, Fluorescence Imaging and Fluorescence Endoscopy in Photodynamic Therapy (PDT) for Cancer Cells", *J Bioanal Biomed* 8: e135, 2016.
- [9] A. Heidari, "A Bio-Spectroscopic Study of DNA Density and Color Role as Determining Factor for Absorbed Irradiation in Cancer Cells", *Adv Cancer Prev* 1: e102, 2016.
- [10] A. Heidari, "Manufacturing Process of Solar Cells Using Cadmium Oxide (CdO) and Rhodium (III) Oxide (Rh<sub>2</sub>O<sub>3</sub>) Nanoparticles", *J Biotechnol Biomater* 6: e125, 2016.
- [11] A. Heidari, "A Novel Experimental and Computational Approach to Photobiosimulation of Telomeric DNA/RNA: A Biospectroscopic and Photobiological Study", *J Res Development* 4: 144, 2016.
- [12] A. Heidari, "Biochemical and Pharmacodynamical Study of Microporous Molecularly Imprinted Polymer Selective for Vancomycin, Teicoplanin, Oritavancin, Telavancin and Dalbavancin Binding", *Biochem Physiol* 5: e146, 2016.
- [13] A. Heidari, "Anti-Cancer Effect of UV Irradiation at Presence of Cadmium Oxide (CdO) Nanoparticles on DNA of Cancer Cells: A Photodynamic Therapy Study", *Arch Cancer Res.* 4: 1, 2016.
- [14] A. Heidari, "Biospectroscopic Study on Multi-Component Reactions (MCRs) in Two A-Type and B-Type Conformations of Nucleic Acids to Determine Ligand Binding Modes, Binding Constant and Stability of Nucleic Acids in Cadmium Oxide (CdO) Nanoparticles-Nucleic Acids Complexes as Anti-Cancer Drugs", *Arch Cancer Res.* 4: 2, 2016.
- [15] A. Heidari, "Simulation of Temperature Distribution of DNA/RNA of Human Cancer Cells Using Time-Dependent Bio-Heat Equation and Nd: YAG Lasers", *Arch Cancer Res.* 4: 2, 2016.
- [16] A. Heidari, "Quantitative Structure-Activity Relationship (QSAR) Approximation for Cadmium Oxide (CdO) and Rhodium (III) Oxide (Rh<sub>2</sub>O<sub>3</sub>) Nanoparticles as Anti-Cancer Drugs for the Catalytic Formation of Proviral DNA from Viral RNA Using Multiple Linear and Non-Linear Correlation Approach", *Ann Clin Lab Res.* 4: 1, 2016.
- [17] A. Heidari, "Biomedical Study of Cancer Cells DNA Therapy Using Laser Irradiations at Presence of Intelligent Nanoparticles", *J Biomedical Sci.* 5: 2, 2016.
- [18] A. Heidari, "Measurement the Amount of Vitamin D2 (Ergocalciferol), Vitamin D3 (Cholecalciferol) and Absorbable Calcium (Ca<sup>2+</sup>), Iron (II) (Fe<sup>2+</sup>), Magnesium (Mg<sup>2+</sup>), Phosphate (PO<sub>4</sub><sup>-</sup>) and Zinc (Zn<sup>2+</sup>) in Apricot Using High-Performance Liquid Chromatography (HPLC) and Spectroscopic Techniques", *J Biom Biostat* 7: 292, 2016.
- [19] A. Heidari, "Spectroscopy and Quantum Mechanics of the Helium Dimer (He<sup>2+</sup>), Neon Dimer (Ne<sup>2+</sup>), Argon Dimer (Ar<sup>2+</sup>), Krypton Dimer (Kr<sup>2+</sup>), Xenon Dimer (Xe<sup>2+</sup>), Radon Dimer (Rn<sup>2+</sup>) and Ununoctium Dimer (Uuo<sup>2+</sup>) Molecular Cations", *Chem Sci J* 7: e112, 2016.
- [20] A. Heidari, "Human Toxicity Photodynamic Therapy Studies on DNA/RNA Complexes as a Promising New Sensitizer for the Treatment of Malignant Tumors Using Bio-Spectroscopic Techniques", *J Drug Metab Toxicol* 7: e129, 2016.
- [21] A. Heidari, "Novel and Stable Modifications of Intelligent Cadmium Oxide (CdO) Nanoparticles as Anti-Cancer Drug in Formation of Nucleic Acids Complexes for Human Cancer Cells' Treatment", *Biochem Pharmacol (Los Angel)* 5: 207, 2016.
- [22] A. Heidari, "A Combined Computational and QM/MM Molecular Dynamics Study on Boron Nitride Nanotubes (BNNTs), Amorphous Boron Nitride Nanotubes (a-BNNTs) and Hexagonal Boron Nitride Nanotubes (h-BNNTs) as Hydrogen Storage", *Struct Chem Crystallogr Commun* 2: 1, 2016.
- [23] A. Heidari, "Pharmaceutical and Analytical Chemistry Study of Cadmium Oxide (CdO) Nanoparticles Synthesis Methods and Properties as Anti-Cancer Drug and its Effect on Human Cancer Cells", *Pharm Anal Chem Open Access* 2: 113, 2016.

- [24] A. Heidari, "A Chemotherapeutic and Biospectroscopic Investigation of the Interaction of Double-Standard DNA/RNA-Binding Molecules with Cadmium Oxide (CdO) and Rhodium (III) Oxide (Rh<sub>2</sub>O<sub>3</sub>) Nanoparticles as Anti-Cancer Drugs for Cancer Cells' Treatment", *Chemo Open Access* 5: e129, 2016.
- [25] A. Heidari, "Pharmacokinetics and Experimental Therapeutic Study of DNA and Other Biomolecules Using Lasers: Advantages and Applications", *J Pharmacokinet Exp Ther* 1: e005, 2016.
- [26] A. Heidari, "Determination of Ratio and Stability Constant of DNA/RNA in Human Cancer Cells and Cadmium Oxide (CdO) Nanoparticles Complexes Using Analytical Electrochemical and Spectroscopic Techniques", *Insights Anal Electrochem* 2: 1, 2016.
- [27] A. Heidari, "Discriminate between Antibacterial and Non-Antibacterial Drugs Artificial Neural Networks of a Multilayer Perceptron (MLP) Type Using a Set of Topological Descriptors", *J Heavy Met Toxicity Dis.* 1: 2, 2016.
- [28] A. Heidari, "Combined Theoretical and Computational Study of the Belousov-Zhabotinsky Chaotic Reaction and Curtius Rearrangement for Synthesis of Mechlorethamine, Cisplatin, Streptozotocin, Cyclophosphamide, Melphalan, Busulphan and BCNU as Anti-Cancer Drugs", *Insights Med Phys.* 1: 2, 2016.
- [29] A. Heidari, "A Translational Biomedical Approach to Structural Arrangement of Amino Acids' Complexes: A Combined Theoretical and Computational Study", *Transl Biomed.* 7: 2, 2016.
- [30] A. Heidari, "Ab Initio and Density Functional Theory (DFT) Studies of Dynamic NMR Shielding Tensors and Vibrational Frequencies of DNA/RNA and Cadmium Oxide (CdO) Nanoparticles Complexes in Human Cancer Cells", *J Nanomedicine Biotherapeutic Discov* 6: e144, 2016.
- [31] A. Heidari, "Molecular Dynamics and Monte-Carlo Simulations for Replacement Sugars in Insulin Resistance, Obesity, LDL Cholesterol, Triglycerides, Metabolic Syndrome, Type 2 Diabetes and Cardiovascular Disease: A Glycobiological Study", *J Glycobiol* 5: e111, 2016.
- [32] A. Heidari, "Synthesis and Study of 5-[(Phenylsulfonyl)Amino]-1,3,4-Thiadiazole-2-Sulfonamide as Potential Anti-Pertussis Drug Using Chromatography and Spectroscopy Techniques", *Transl Med (Sunnyvale)* 6: e138, 2016.
- [33] A. Heidari, "Nitrogen, Oxygen, Phosphorus and Sulphur Heterocyclic Anti-Cancer Nano Drugs Separation in the Supercritical Fluid of Ozone (O<sub>3</sub>) Using Soave-Redlich-Kwong (SRK) and Pang-Robinson (PR) Equations", *Electronic J Biol* 12: 4, 2016.
- [34] A. Heidari, "An Analytical and Computational Infrared Spectroscopic Review of Vibrational Modes in Nucleic Acids", *Austin J Anal Pharm Chem.* 3 (1): 1058, 2016.
- [35] A. Heidari, C. Brown, "Phase, Composition and Morphology Study and Analysis of Os-Pd/HfC Nanocomposites", *Nano Res Appl.* 2: 1, 2016.
- [36] A. Heidari, C. Brown, "Vibrational Spectroscopic Study of Intensities and Shifts of Symmetric Vibration Modes of Ozone Diluted by Cumene", *International Journal of Advanced Chemistry*, 4 (1) 5-9, 2016.
- [37] A. Heidari, "Study of the Role of Anti-Cancer Molecules with Different Sizes for Decreasing Corresponding Bulk Tumor Multiple Organs or Tissues", *Arch Can Res.* 4: 2, 2016.
- [38] A. Heidari, "Genomics and Proteomics Studies of Zolpidem, Necopidem, Alpidem, Saripidem, Miroprofen, Zolimidine, Olprinone and Abafungin as Anti-Tumor, Peptide Antibiotics, Antiviral and Central Nervous System (CNS) Drugs", *J Data Mining Genomics & Proteomics* 7: e125, 2016.
- [39] A. Heidari, "Pharmacogenomics and Pharmacoproteomics Studies of Phosphodiesterase-5 (PDE5) Inhibitors and Paclitaxel Albumin-Stabilized Nanoparticles as Sandwiched Anti-Cancer Nano Drugs between Two DNA/RNA Molecules of Human Cancer Cells", *J Pharmacogenomics Pharmacoproteomics* 7: e153, 2016.
- [40] A. Heidari, "Biotranslational Medical and Biospectroscopic Studies of Cadmium Oxide (CdO) Nanoparticles-DNA/RNA Straight and Cycle Chain Complexes as Potent Anti-Viral, Anti-Tumor and Anti-Microbial Drugs: A Clinical Approach", *Transl Biomed.* 7: 2, 2016.