

What Is the Effect of Physical Exercise on Nafld/Nash

Maria Fernanda Lopes Ferreira^{1,2}, Icaro Putinhon Caruso^{1,2,3}, Marcelo Andres Fossey^{1,2}, Walnei Fernandes Barbosa⁴, Fatima Pereira De Souza^{1,2,*}

¹Department of Physics, Paulista State University, Sao Jose do Rio Preto, Brazil

²Multiusers Center for Biomolecular Innovation, Paulista State University, Sao Jose do Rio Preto, Brazil

³National Center for Nuclear Magnetic Resonance of Macromolecules, Institute of Medical Biochemistry and National Center for Structure Biology and Bioimaging, Ilha do Fundao, Rio de Janeiro, Brazil

⁴Department Gastroenterology Paulista State University, Botucatu Medical School, Botucatu, Brazil

Email address:

mariafer.lopes.ferreira@gmail.com (M. F. L. Ferreira), icaro.caruso@unesp.br (I. P. Caruso), marcelo.fossey@unesp.br (M. A. Fossey), walnei.fernandes@gmail.com (W. F. Barbosa), fatima.p.souza@unesp (F. P. de Souza)

*Corresponding author

To cite this article:

Maria Fernanda Lopes Ferreira, Icaro Putinhon Caruso, Marcelo Andres Fossey, Walnei Fernandes Barbosa, Fatima Pereira De Souza. What Is the Effect of Physical Exercise on Nafld/Nash. *American Journal of Sports Science*. Vol. 9, No. 1, 2021, pp. 1-7.

doi: 10.11648/j.ajss.20210901.11

Received: November 17, 2020; **Accepted:** January 14, 2021; **Published:** January 22, 2021

Abstract: Non-alcoholic fat liver disease (NAFLD) is a pathologic entity characterized by an excessive accumulation of hepatic lipid without alcohol consumption. Both endurance and strength exercise have shown to be beneficial in NAFLD. PubMed database has been searched for randomized trials and prospective cohort studies in adults aged ≥ 18 , rats and mice that have investigated the effects of at least 4 weeks of exercise only or combination with diet on NAFLD from 2014 to 2019. The review selected have been those in which exercise was clearly described by type, duration, intensity, and frequency, and that the NAFLD diagnosis, as well as the outcome measures, were confirmed through, at least, one of the following methods: biopsy, anthropometric measures, blood testing, ultrasonography imaging, biochemical analysis and Image Resonance Magnetic (MRI). Lifestyle has shown to greatly influence human behavior and health, however, a major factor impacting studies results is the difficulty in controlling people's compliance with a healthier lifestyle, unlike animals that are forced to exercise over time, frequency and intensity, according to the researchers' wish. Therefore, animal studies have shown better results of the effect of exercise on the disease. However, both human and animal studies have reached a positive change in the following parameters: weight, fat percentage, blood glucose, insulin and ALT and AST levels. Physical exercise improves non-alcoholic fat liver disease (NAFLD), however, an optimum approach is still unclear. A regular and long-term systematic practice of exercise leads to better general health, body weight control, and life quality improvement.

Keywords: Physical Exercise, NAFLD, Metabolic Syndrome, NASH

1. Introduction

Non-alcoholic fatty liver disease (NAFLD) is an increasing cause of chronic liver disease worldwide [40]. NAFLD occurs in patients with no history of alcohol abuse, in which there is an excessive increase in fat in the liver [40].

Although the majority of NAFLD patients have the only steatosis without progression, a considerable fraction develops non-alcoholic hepatitis stasis (NASH), which can lead to fibrosis in 20 to 40% of cases, of which 10% of cases can progress to cirrhosis and 1-5% develop hepatic cell

carcinoma (HCC) [39, 41]. It is estimated that about one-third of early-stage NASH cases will progress to stage 3 or 4 fibrosis (cirrhosis) over 5 to 10 years [42], also increasing the risk of mortality as fibrosis progresses for more severe stages [43].

NAFLD can be caused by interactions of environmental and genetic factors or by metabolic complications such as hyperglycemia, diabetes, dyslipidemia, metabolic syndrome, and hypoadiponectinemia, so that these are associated with the development of non-alcoholic hepatitis stasis (NASH) and liver fibrosis [40, 44]. In NASH, the factors

hyperinsulinemia, adipose inflammation, and hypoadiponectinemia deregulate the turnover of one or more hepatic lipid reservoirs, leading to lipid accumulation in the hepatocytes that leads to inflammation, a process known as lipotoxicity [40, 48].

Regardless of having no FDA (Food and Drug Administration)-approved pharmacologic therapy is currently available, lifestyle changes have been suggested as a nonpharmacologic option to combat NAFLD associated with metabolic conditions. The guidelines for NAFLD treatment recommend the combination of daily exercise and caloric restriction, targeting at 3–10% weight loss [5].

The compilation, such as metabolic disorders, contribute to cardiovascular and cerebrovascular events, the major causes of death in NAFLD. However, these findings justify the importance of targeting the skeletal muscle in NAFLD to induce lifestyle modifications that improve cardiovascular and cerebrovascular morbidity and mortality. Studies on physical exercise in NAFLD shows significant benefits on hepatic steatosis, serum transaminases, lipid profile, and glycemic control that can potentially lower cardiovascular and cerebrovascular events [22].

Al-Jiffri, O. *et al.* [2] postulated that there is no proven treatment for patients with NAFLD are currently available. Weight control and reduction, as well as lifestyle modifications with diet changes and increased physical activity, are generally recommended as the first step in the treatment of patients with this condition. Decreasing and maintaining weight reduction can improve NAFLD, but the results of several reports are inconsistent.

This review is intend to understand and compare the effect of physical exercise on humans and rats with NAFLD.

Studies have shown that both endurance and strength exercise are beneficial in NAFLD. Two interesting questions are how does exercise benefit NAFLD and what is/are the mediators of the muscle-liver axis whereby the muscle regulates hepatic metabolism. However, considering the scarcity of studies that suggest a definite and precise answer to these questions, further research is needed in order to widen knowledge into the mechanisms through which physical exercise has a positive impact on NASH patients' life quality [25, 34, 38].

Therefore, the development of a prevention and treatment strategy includes changes in lifestyle, such as caloric restriction, diets with low fat and low fat and low glycemic index, and increasing consumption of foods rich in beneficial bioactive ingredients, and the practice of regular physical activity. Rosa-Caldewell *et al.* [13], shows that vegetables exhibit a variety effects related to their antioxidant, hypoglycemic, and hypolipidemic properties due to the specific characteristics of vegetables proteins, carbohydrates and lipids, and also from several non-nutritional compounds such as polyphenols, phytic acid or α -galactoside oligosaccharides [13, 34].

According to Cho, J. *et al.* [5] Show that caloric restriction solidified its potential in treating NAFLD, however, combined with physical activity resulted in a greater decrease

in body and abdominal adiposity, decreased blood lipids and decreased intrahepatic fat content, together with decreased hepatic glucose production and improved insulin sensitivity. Noting that physical exercise has a very positive beneficial effect as it favors greater weight loss through negative energy balance. Therefore, if caloric restriction alone is not effective, physical exercise should be included in NAFLD treatment.

Autophagy is a cellular catabolic process that causes the degradation of the liver cell components itself, which can improve NAFLD by reversing oxidative stress and mitochondrial dysfunction that contribute to hepatocyte damage and activation of stellate cells and fibrosis decreases the accumulation of liver lipids and fibrosis in NAFLD [5, 27].

According to Wang, B. *et al.* [28], the exercise benefits the NAFLD has mainly autophagy in NAFLD. Physical exercise is a non-pharmacological intervention capable of improving liver function test in NAFLD, ensuring that physical exercise is beneficial through improved autophagy in skeletal muscle and possibly in the liver. Wang, B *et al.* [28], is currently looking for the mechanisms by which exercise stimulates liver autophagy. However, the beneficial of physical exercise cannot be attributed only to the stimulation of autophagy.

Physical exercise increases the secretion of hormone irisin. The moderating increase by the exercise of circulating irisin is likely strongly related to the improvement of body fat mass, insulin resistance and growth hormone. Although irisin increases with exercise, there is a tendency to decrease with exercise maintenance and the progressive loss of its benefits. One way to maintain your production is stimulation through the variation of exercises that must be accompanied by a good professional in the field. Interventions focused on monitoring the NAFLD are weighted, so physical exercise with caloric restriction provide this overall benefit. Long-term studies are needed to establish how exercise produces metabolic organ homeostasis in patients with NAFLD.

Longitudinal studies are needed to stablish the mechanisms by which physical exercise produce homeostasis of the triad of metabolic organs in patients with NAFLD. These searchers will provide the case for optimizing physical exercise and lifestyle changes, including dietary reeducation to reverse liver and non-liver organ dysfunction, and clinical consequences in NAFLD and metabolic syndrome.

2. Study Identification and Selection

The methodology used in this review was through the selection of bibliographic research through scientific articles in the PubMed database. The search terms used to identify the articles were: ("NAFLD", "non-alcoholic fatty liver disease", "NASH" "non-alcoholic fatty liver", "fat", "fatty liver", "steatosis") and ("exercise", "aerobic training", "resistance training", "resistance training", "strength training", "diet").

Animal and human studies, adults ≥ 18 years old of any sex or ethnic origin, were investigated that investigated the effects of at least 4 weeks of exercise or combination with diet in the NAFLD published in the period 2014 to 2019.

NAFLD diagnoses and the response measures, were confirmed by at least one of the following methods: biopsy, anthropometric measurements, blood count, ultrasound, biochemical analysis and magnetic resonance imaging (MRI). Trials that included dietary interventions along with exercise were accepted if they met all criteria.

3. Evaluation of the Effect of Physical Exercise on NAFLD

Studies investigating the effect of physical exercise whether associated to diet (hypercaloric for rats and mice, and hypocaloric for humans) or not, in NAFLD/NASH subjects with insulin sensibility. The selection of the methodology of studies clearly prescribed its intervention, such as the duration, intensity and frequency of physical exercise.

Studies that did not specific exercise prescriptions, outcome measures demonstrating an exercise effect (*i.e.*, measures of fitness and/or strength), and quantitative measures of intrahepatic fat, were not included. Moreover, studies or study arms for which dietary supplements, herbal preparations, nutraceuticals were the intervention to the study were also included. All studies performed calculations of

changes in relation to the percentage of liver fat, and this reduction was compared between the groups that received exercise and those that received a diet plus physical exercise.

The revised animal studies herein yield beneficial results regarding the effect of exercise as shown table 1. Five studies Singh V, et al. [3], Batatinha HAP, et al. [9], Marcinko K, et al. [14], MacLean C, et al. [18], Shojaei-Moradie F, et al. [23] have had a significant improvement of the disease with exercise, which had in their training method an intensity of 10 to 20 m/min with a frequency of 5 times a week for 60 minutes. Rosa-Caldewell, M. E et al. [23], have applied their volunteer training method with 4 weeks of intervention, getting better levels both of AST and TNF- α IL-6.

The remaining animal studies all have had beneficial results, which also have used their method similar to those of significant results, being different from those of Frantz, E. D. C et al. [28] with a frequency of 4 times a week, and Oh S, et al. [16] with 3 times a week.

Overall, the articles have presented similar methods, being different only as to the duration of the study, which has ranged from a minimum of 4 weeks to a maximum of 16 weeks. These studies have shown similar results: improvement in AST, ALT, insulin levels, glucose, TNF- α and IL-6, weight reduction and percentage fat.

Table 1. Methodology and results of the effect of physical exercise on animals.

Author	Method					Results						Conclusion
	N	Intensity	Session length	Session frequency	Time of intervention	Results that there was improvement						
KAWANISHI, N. et al., [11]	34/4	15 – 20 m/min	60 min	5 x week	16 weeks	% Fat	ALT	TNF- α , IL-6	TG	–		Significantly
AAMANNA, L, et al. [23]	26/2			5 x week	4 weeks	% Fat	ALT	AST	–	–		Moderate
BATATINHA, H. A. P. et al., [9]		60% VO ₂ Max.	60 min	5 x week	12 weeks	% Fat	Weight	Glucose	ALT	TNF- α IL-6		Significantly
CHO, J. et al., [2]		12 m/min	50 min	5 x week	16 weeks	% Fat	Weigh	ALT	TG	TNF- α IL-6		Beneficial
GONÇALVES, I. O. PASSOS, E. DIOGO, C. et al. [26]		15-25 m/min	60 min	5 x week	17 weeks	Mitochondrial Improvement						Beneficial
MIOTTO, P. M. et al. [29]	16/8	15 m/min	45 min	5 x week	4 weeks	% Fat	Weigh	TG				Beneficial
ROSA-CALDWELL, M. et al. [13]	20/2	Voluntary roads				AST	TNF- α IL-6	–	–	–		Significantly
WANG, B. Et al. [8]		10 m/min	60 min	4 x week	24 weeks	ALT	TG	TNF- α IL-6	Stress oxidative			Beneficial
WINN, N. C. et al. [33]		Voluntary roads				Weigh	Glucose					Beneficial
XIAO, J. et al. [1]	40/4	10 m/min	60 min	5x week	16 weeks	Weigh	Insulin	AST	Alt	TG	Stress Oxidative	Beneficial
YOSHIMURA, S. et al. [35]		Voluntary roads				% Fat	ALT					Beneficial
KAPRAVELOU, G. et al, [7]	80/10	65-80% V VO ₂ Max. HIT	60 min	5x week	8 weeks	% Fat	AST	ALT	Glucose	TG		Beneficial
MOON, H. Y. et al. [12]	19/2	10 m/min	50 min	5 x week	4 weeks	%Fat	ALT	TG				Significantly
HACZEYNI, F. et al. [48]		Voluntary roads				Weigh	ALT	Insulin				Beneficial
FRANTZ, E. D. C. et al. [38]	32/4	50-75% VO ₂ Max.	60	4 x week	8 weeks	TNF- α IL-6						Beneficial
LINDEN, M. A. et al. [47]	30/10	15/20m/min 40 m/min	60 min 30 min	5 x week	12 weeks	%Fat	Weigh	Insulin	Glucose	ALT	TNF- α IL-6	Beneficial

Author	Method					Results				Conclusion
	N	Intensity	Session length	Session frequency	Time of intervention	Results that there was improvement				
MARCINKO, K. et al. [14]			sprint							
PASSOS, E. PEREIRA, C. D. et al. [31]		15-22 m/min	60 min	3 x week	6 weeks	%Fat	Insulin	ALT		Beneficial
GONÇALES, I. O. et al. [37]		15-25 m/min	60 min	5 x week	8 weeks	Insulin	TNF-α	IL-6		Beneficial
GONÇALVES, I. O. et al. [10]		15-25 m/min	60 min	5 x week	8 weeks	Weigh	ALT	AST	TG	Significantly
		15-25 m/min	60 min	5 x week	8 weeks	%Fat	Stress	Oxidative		Beneficial

N = number of participants, VO₂ oxygen volume

The human studies have yielded 3 types of results as shown table 2, significant, beneficial and those, which have shown no influence of physical exercise on the disease. Zelber-Sagi, S.; Al-Jiffri, O. et al. [33, 2] have found significant improvements in ALT, AST, insulin, total cholesterol and BMI, having in their method an intensity of 60 to 75% HR max, with frequency of 3 times a week and duration of 40-45 minutes each session, for three months.

Five studies have shown no improvement in the disease due to physical exercise. They have had some differences in their methods compared to those studies, which have demonstrated significant results. Oh, S. et al. [20] have applied the exercise at an intensity of 40% H. R Max,

Takahashi A, eta al. [6] have worked with 50% of the maximum strength in bodybuilding exercises, Zelber-Sagi S et al. [15] have applied 5 sprint shots, 2 times a week, with a small number of participants, and Malin, S. K. et al. [36] have used only 7 days of intervention. These non-significant results may have been due to the method applied and also to difficulties in controlling patients compliance with the training program, however Oh, S.; Croci, I. et al. [20, 6] have obtained better results in their diet groups.

In human studies, in general, the exercise sessions frequency have been 3 times a week, lasting 45 to 60 minutes and with large variations in intervention time, ranging from seven days to 16 weeks.

Table 2. Methodology and results of the effect of physical exercise on humans.

Author	Method					Results								Conclusion
	N	Intensity	Session length in times	Session frequency	Time of intervention	Results that there was improvement								
KULLMAN, E. L. et al. [46]	22	85% HR Max.	60 min	7days	7days	Glucose Insulin								Beneficial
OH, S. et al. [16]	72	40% HR Max,	90 min	3x week	3 moths	Diet had more benefits								No Influence
SHOJAE, F. et al. [22]	29	40-60% HR Max.	60 min	4x week	16 weeks	%Fat	AST	ALT	Weigh	IMC	Circumference	Insulin fasting	Beneficial	
TAKAHASHI, A. et al. [6]	53	3x10 rep	30 min	3x week	12 weeks	Weigh	IMC	ALT	AST	Insulin	Cholesterol	IHCL	Beneficial	
TUTINO, V. et al. [32]	142	60-70% HR Max.	45 min	3x week Body building		IMC	HDL	HOMA-IR					Beneficial	
YOSHIMURA, E. et al [34]	89		60 min	3x week	12 weeks	%Fat	TG	TNF- α IL-6	IMC	Visceral fat index			No Influence	
ZELBER-SAGI, S. et al. [15]	64	60-75%	45 min	3X Week Body building	3 moths	Weigh	IMC	Cholesterol Total					Significantly	
HOUGHTON, D. et al. [21]	24	Hard	45-60% HR Max.	3x week	12 weeks	%Fat	TG	Insulin	HTGC				Beneficial	
ZANG, H-J. et al. [36]	208	45-55% 65-80% HR Max	30 min	5x week	12 mouths	Weigh	TG	Circumference	Risk Factors				Beneficial	
TANIGUCHI, H. et al. [20]	27	60-75%	45 min	3-5x week	5 weeks	%Fat	AST	TG	%Fat visceral				Beneficial	
CROCI, I. et al. [17]	16	50% 1RM	60 min	3x week	6 moths	%Fat	Diet had more benefits						No Influence	
PUGH, C. A. J. et al. [45]	21	45-60% HR Max.	45 min	3-5x week	16 weeks	IMC	Weigh	Glucose	Circumference				Beneficial	
MACLEAN, C. et al. [18]	9	Sprint 5 shots	60 min	2x week	6 weeks	%Fat						No Influence		
AL-JIFFRI, O. et al. [5]	100	65-75% HR Max.	40 min	3x week	3 moths	IMC	ALT	AST	Insulin				Significantly	

Author	Method					Results	Conclusion
	N	Intensity	Session length in times	Session frequency	Time of intervention	Results that there was improvement	
MALIN, S. K. et al. [19]	13	85% HR Max.	60 min	7 Days	7 days	Fetuin-A	No influence

HR = heart rate; Fetuin-A = are blood proteins that are produced in the liver and secreted into the bloodstream

In animal studies, the intensity, and frequency of the applied method have been higher than those in human studies; consequently, the results have also shown more improvement in the disease due to physical exercise. This might have been because of difficulties in changing people's lifestyle. Therefore, considering that lifestyle greatly influences human behavior and health, a major factor impacting studies results is the difficulty in controlling people's compliance to a healthier lifestyle, unlike animals which are forced to exercise over time, frequency and intensity, according to the researchers' wish.

The AIT method that the author used in this study is a viable intervention technique to improve plasma and hepatic biochemical parameters as well as hepatic histological alterations inherent to early stages of NAFLD in obese Zucker rats. The training protocol was especially efficient to improve insulin sensitivity and decrease the hepatic lipid content, as well as ameliorating the oxidative stress conditions in this organ [5, 30].

Different methods and parameters have been used to analyses data in animal and human studies. Human studies have been done through anthropometric and laboratory measurements, whereas rats or their livers have been analyzed by immunohistochemical and blood tests. Animal studies have shown better results of the effect of exercise on the disease, while less significant result have been observed in humans. However, both human and animal studies have reached a positive change in the following parameters: weight, fat percentage, blood glucose, insulin and ALT and AST levels [4].

The study herein has found that a 5 times/week endurance exercise program decreases hepatic fat content and serum fibroblast growth factor 21 (FGF21) levels, without weight loss in elderly Japanese men. These evidences suggest that endurance physical exercise improves the resistance of FGF21 without weight loss and provides further clarification that exercise alone, without weight loss, has beneficial effects prevention of age-related cardio metabolic diseases [26].

Triglycerides, visceral fat and circulating HTGC in adults with NASH decreased with physical exercise, regardless of weight loss. Therefore, the intensity of the exercise must be considered in the prescription of exercise and intervention programs for NASH treatment for being effective in reducing the content of hepatic triglycerides (HTGC). Although physical exercise has a beneficial effect on HTGC and visceral fat for people with NASH, caloric restriction plays an essential role in the disease [10].

The exercise intervention in nondiabetic men with NAFLD has significantly improved both fitness and cardiometabolic health. In addition, they have shown a significant reduction

in intra hepatocellular fat (IHCL), however, their liver continued to export excessive amounts of triglycerides (TGs) in very-low-density lipoprotein (VLDL). This might have been due to a failure in normalizing IHCL and restoring hepatic insulin sensitivity. Hence, a longer duration or higher-intensity exercise intervention, or an approach combined with calorie restriction might be required to achieve lower amounts of TG and VLDL exportation and to lower plasma TG and VLDL production rates [24].

4. Conclusion

In conclusion, physical exercise improves non-alcoholic fat liver disease (NAFLD), however an optimum approach is still unclear. Regardless, a regular and long-term systematic practice of exercise leads to a better general health, body weight control, and life quality improvement. Interestingly, findings of the reviewed authors are quite varied and of the low significance level. However, 90% of them have shown a positive and/or beneficial impact on the subjects, without necessarily reducing the fat content specifically of the liver though. Therefore, NAFLD and NASH require pharmacological intervention, and physical exercise is an adjunctive therapy to enhance the effect of medication. Further studies should be carried out in humans under drug therapy to assess the impact of exercise on the effect of drugs. Furthermore, more specific studies involving animals are needed in order to reach accurate guidelines for future human applications, considering that although studies based on anthropometric and histochemical measurements have not shown significant impact on the measured parameters, most of them have, indeed, shown, in practice, to varied extents, improvements in several aspects analyzed, molecular research, both in animals and in humans, should be done seeking for more visible and measurable beneficial changes.

List of Abbreviations

NAFLD: Non-alcoholic fat liver disease; MRI: Image Resonance Magnetic; ALT: Alanine transaminase; AST: Aspartate transaminase; TG: Triglycerides; NASH: Non-alcoholic steatohepatitis; HCC: Hepatic cellular carcinoma; FDA: Food and Drug Administration; EX: exercise; CR: Caloric restriction; AMPK: AMP-activated protein kinase; FGF21: Fibroblast growth factor 21; VLDL: very-low-density lipoprotein; HR: heart rate; H. R Max: Heart rate maximum; mTORC1: mammalian target of rapamycin; TNF- α : Factor de Necrose Tumoral Alfa; IL-6: interleucina 6; AIT: aerobic interval training; HTGC: hepatic triglyceride content; IHCL: intrahepatocellular lipids.

Availability of Data and Material

All available data's and material are found in the manuscript.

Competing Interests

The authors declare that they have no competing interests.

Contributions

MFLF and FPS: Conceived the design, searching the literature, drafting the manuscript; FPS, IPC, MAF and WFB: searching literature, supervising and critical review of the manuscript. All authors read and approved the final manuscript for publication.

References

- [1] Xiao J, Bei Y, Liu J, *et al.* miR-212 downregulation contributes to the protective effect of exercise against non-alcoholic fatty liver via targeting FGF-21. *J Cell Mol Med.* 2016; 20 (2): 204–16.
- [2] Cho J, Koh Y, Han J, *et al.* Adiponectin mediates the additive effects of combining daily exercise with caloric restriction for treatment of non-alcoholic fatty liver. *International Journal of Obesity.* 2016; 40: 1760.
- [3] Singh V, Luthra S, Elajami TK, *et al.* Resolution of NASH with weight loss documented by hepatic MRI. *BMJ Case Rep.* 2015, doi: 10.1136/bcr-2014-207107.
- [4] Pinto CG de S, Marega M, de Carvalho JAM, *et al.* Physical activity as a protective factor for development of non-alcoholic fatty liver in men. *Einstein (Sao Paulo).* 2015; 13 (1): 34–40.
- [5] Al-Jiffri O, Al-Sharif F, Abd El-Kader S, *et al.* Weight reduction improves markers of hepatic function and insulin resistance in type-2 diabetic patients with non-alcoholic fatty liver. *Afr Health Sci.* 2013; 13 (3): 667–72.
- [6] Takahashi A, Abe K, Usami K, *et al.* Simple Resistance Exercise helps Patients with Non-alcoholic Fatty Liver Disease. *Int J Sports Med.* 2015; 36 (10): 848–52.
- [7] Kapravelou G, Martínez R, Nebot E, *et al.* The Combined Intervention with Germinated Vigna radiata and Aerobic Interval Training Protocol Is an Effective Strategy for the Treatment of Non-Alcoholic Fatty Liver Disease (NAFLD) and Other Alterations Related to the Metabolic Syndrome in Zucker Rats. *Nutrients.* 2017; 9 (7).
- [8] Wang B, Zeng J, Gu Q. Exercise restores bioavailability of hydrogen sulfide and promotes autophagy influx in livers of mice fed with high-fat diet. *Can J Physiol Pharmacol.* 2017; 95 (6): 667–74.
- [9] Batatinha HAP, Lima EA, Teixeira AAS, *et al.* Association Between Aerobic Exercise and Rosiglitazone Avoided the NAFLD and Liver Inflammation Exacerbated in PPAR- α Knockout Mice. *J Cell Physiol.* 2017; 232 (5): 1008–19.
- [10] Gonçalves IO, Passos E, Rocha-Rodrigues S, *et al.* Physical exercise antagonizes clinical and anatomical features characterizing Lieber-DeCarli diet-induced obesity and related metabolic disorders. *Clin Nutr.* 2015; 34 (2): 241–7.
- [11] Kawanishi N, Yano H, Mizokami T, *et al.* Exercise training attenuates hepatic inflammation, fibrosis and macrophage infiltration during diet induced-obesity in mice. *Brain, Behavior, and Immunity.* 2012; 26 (6): 931–41.
- [12] Moon HY, Song P, Choi CS, *et al.* Involvement of exercise-induced macrophage migration inhibitory factor in the prevention of fatty liver disease. *Journal of Endocrinology.* 2013; 218 (3): 339–48.
- [13] Rosa-Caldwell ME, Lee DE, Brown JL, *et al.* Moderate physical activity promotes basal hepatic autophagy in diet-induced obese mice. *Appl Physiol Nutr Metab.* 2017; 42 (2): 148–56.
- [14] Marcinko K, Sikkema SR, Samaan MC, *et al.* High intensity interval training improves liver and adipose tissue insulin sensitivity. *Mol Metab.* 2015; 4 (12): 903–15.
- [15] Zelber-Sagi S, Buch A, Yeshua H, *et al.* Effect of resistance training on non-alcoholic fatty-liver disease a randomized-clinical trial. *WJG.* 2014; 20 (15): 4382.
- [16] Oh S, Tanaka K, Tsujimoto T, *et al.* Regular exercise coupled to diet regimen accelerates reduction of hepatic steatosis and associated pathological conditions in nonalcoholic fatty liver disease. *Metab Syndr Relat Disord.* 2014; 12 (5): 290–8.
- [17] Croci I, Byrne NM, Chachay VS, *et al.* Independent effects of diet and exercise training on fat oxidation in non-alcoholic fatty liver disease. *WJH.* 2016; 8 (27): 1137.
- [18] MacLean C, Dillon J, Babraj JA, *et al.* The effect of low volume sprint interval training in patients with non-alcoholic fatty liver disease. *Phys Sportsmed.* 2018; 46 (1): 87–92.
- [19] Malin SK, Mulya A, Fealy CE, *et al.* Fetuin-A is linked to improved glucose tolerance after short-term exercise training in nonalcoholic fatty liver disease. *Journal of Applied Physiology.* 2013, doi: org/10.1152/jappphysiol.00237.2013.
- [20] Taniguchi H, Tanisawa K, Sun X, *et al.* Endurance Exercise Reduces Hepatic Fat Content and Serum Fibroblast Growth Factor 21 Levels in Elderly Men. *The Journal of Clinical Endocrinology & Metabolism.* 2016; 101 (1): 191–8.
- [21] Houghton D, Thoma C, Hallsworth K, *et al.* Exercise Reduces Liver Lipids and Visceral Adiposity in Patients With Nonalcoholic Steatohepatitis in a Randomized Controlled Trial. *Clinical Gastroenterology and Hepatology.* 2017; 15 (1): 96-102. e3.
- [22] Shojaei-Moradie F, Cuthbertson DJ, Barrett M, *et al.* Exercise Training Reduces Liver Fat and Increases Rates of VLDL Clearance But Not VLDL Production in NAFLD. *J Clin Endocrinol Metab.* 2016; 10.
- [23] Amann L, Ochoa-Sanchez R, Oliveira M, *et al.* Progressive resistance training prevents loss of muscle mass and strength in bile duct-ligated rats. *Liver Int.* 2019; 39 (4): 676–83.
- [24] Brouwers B, Hesselink MKC, Schrauwen P, *et al.* Effects of exercise training on intrahepatic lipid content in humans. *Diabetologia.* 2016; 59 (10): 2068–79.
- [25] Gaspar RC, Kuga GK, Muñoz VR, *et al.* Physical training prevent and treat hepatic lipid accumulation induced by fructose-rich diet. *Biosci j (Online).* 2018; 34 (4): 1041–50.

- [26] Gonçalves IO, Passos E, Diogo CV, *et al.* Exercise mitigates mitochondrial permeability transition pore and quality control mechanisms alterations in nonalcoholic steatohepatitis. *Appl Physiol Nutr Metab.* 2016; 41 (3): 298–306.
- [27] Huber Y, Pfirrmann D, Gebhardt I, *et al.* Improvement of non-invasive markers of NAFLD from an individualised, web-based exercise program. *Alimentary Pharmacology & Therapeutics*, 2019, doi: [org/10.1111/apt.15427](https://doi.org/10.1111/apt.15427).
- [28] Kapravelou G, Martínez R, Andrade AM, *et al.* Aerobic interval exercise improves parameters of nonalcoholic fatty liver disease (NAFLD) and other alterations of metabolic syndrome in obese Zucker rats. *Appl Physiol Nutr Metab.* 2015; 40 (12): 1242–52.
- [29] Miotto PM, Horbatuk M, Proudfoot R, *et al.* α -Linolenic acid supplementation and exercise training reveal independent and additive responses on hepatic lipid accumulation in obese rats. *Am J Physiol Endocrinol Metab.* 2017; 312 (6): E461–70.
- [30] Motta VF, Aguila MB, Mandarim-de-Lacerda CA. High-intensity interval training beneficial effects in diet-induced obesity in mice: adipose tissue, liver structure, and pancreatic islets. *Int j morphol.* 2016; 34 (2): 684–91.
- [31] Passos E, Pereira CD, Gonçalves IO, *et al.* Role of physical exercise on hepatic insulin, glucocorticoid and inflammatory signaling pathways in an animal model of non-alcoholic steatohepatitis. *Life Sci.* 2015; 123: 51–60.
- [32] Tutino V, De Nunzio V, Caruso MG, Bonfiglio C, *et al.* Aerobic Physical Activity and a Low Glycemic Diet Reduce the AA/EPA Ratio in Red Blood Cell Membranes of Patients with NAFLD. *Nutrients*, 2018; 10 (9): 1299. doi: [10.3390/nu10091299](https://doi.org/10.3390/nu10091299)
- [33] Winn NC, Jurrissen TJ, Grunewald ZI, *et al.* Estrogen receptor- α signaling maintains immunometabolic function in males and is obligatory for exercise-induced amelioration of nonalcoholic fatty liver. *Am J Physiol Endocrinol Metab.* 2019; 316 (2): E156–67.
- [34] Yoshimura E, Kumahara H, Tobina T, *et al.* Lifestyle Intervention Involving Calorie Restriction with or without Aerobic Exercise Training Improves Liver Fat in Adults with Visceral Adiposity. *J Obes*, 2014, 197216. doi: [10.1155/2014/197216](https://doi.org/10.1155/2014/197216)
- [35] Yoshimura S, Nakashima S, Tomiga Y, *et al.* Short- and long-term effects of high-fat diet feeding and voluntary exercise on hepatic lipid metabolism in mice. *Biochem Biophys Res Commun.* 2018; 507 (1–4): 291–6.
- [36] Zhang H-J, He J, Pan L-L, *et al.* Effects of Moderate and Vigorous Exercise on Nonalcoholic Fatty Liver Disease: A Randomized Clinical Trial. *JAMA Internal Medicine.* 2016; 176 (8): 1074–82.
- [37] Gonçalves IO, Passos I, Rocha-Rodrigues I, *et al.* Physical exercise prevents and mitigates non-alcoholic steatohepatitis-induced liver mitochondrial structural and bioenergetics impairments. *Mitochondrion*, 2014, vol: 5, p: 40-51. doi: [org/10.1016/j.mito.2014.03.012](https://doi.org/10.1016/j.mito.2014.03.012).
- [38] Frantz EDC, Medeiros RF, Giori IG, *et al.* Exercise training modulates the hepatic renin-angiotensin system in fructose-fed rats. *Experimental Physiology.* 2017; 102 (9): 1208–20.
- [39] Tsuchida, T.; Lee, Y. A.; Fujiwara, N.; Ybanez, M.; Allen, B.; Martins, S.; Friedman, S. L. A simple diet- and chemical-induced murine NASH model with rapid progression of steatohepatitis, fibrosis and liver cancer. *Journal of Hepatology*, 2018. 69 (2), 385–395.
- [40] Marra F, Lotersztajn S. Pathophysiology of NASH: perspectives for a targeted treatment. *Current Pharmaceutical Design.* 2013.
- [41] Sebastiani, G., alshaalan, R., Wong, P., Rubino, M., Salman, A., Metrakos, P., ... Ghali, P. Prognostic Value of Non-Invasive Fibrosis and Steatosis Tools, Hepatic Venous Pressure Gradient (HVPg) and Histology in Nonalcoholic Steatohepatitis. *PLOS ONE*, 2015. 10 (6), e0128774.
- [42] Chu, Huikuan; Williams, Brandon; Schnabl, Bernd. Gut microbiota, fatty liver disease, and hepatocellular carcinoma. *Liver Research*, 2018. Volume 2, Issue 1, 43-51.
- [43] Younossi, Z.; Anstee, Q. M.; Marietti, M.; HARDY, T.; Henry, L.; Eslam, M.; George, J.; Bugianesi, E. Global burden of NAFLD and NASH: trends, predictions, risk factors and prevention. *Gastroenterology & hepatology*, 2017.
- [44] Takahashi Y, Fukusato T. Histopathology of nonalcoholic fatty liver disease/nonalcoholic steatohepatitis. *World Journal of Gastroenterology*, 2014.
- [45] Pugh, Christopher J. A., Victoria S. Sprung, Graham J. Kemp, Paul Richardson, Fariba Shojae-Moradie, A. Margot Umpleby, Daniel J. Green, N. Timothy Cable, Helen Jones, e Daniel J. Cuthbertson. “Exercise training reverses endothelial dysfunction in nonalcoholic fatty liver disease”. *American Journal of Physiology-Heart and Circulatory Physiology* 307, n° 9 (5 de setembro de 2014): H1298–1306.
- [46] Kullman, Emily L., Karen R. Kelly, Jacob M. Haus, Ciaran E. Fealy, Amanda R. Scelsi, Mangesh R. Pagadala, Chris A. Flask, Arthur J. McCullough, e John P. Kirwan. “Short-term aerobic exercise training improves gut peptide regulation in nonalcoholic fatty liver disease”. *Journal of Applied Physiology* 120, n° 10 (31 de março de 2016): 1159–64.
- [47] Linden, Melissa A., Justin A. Fletcher, E. Matthew Morris, Grace M. Meers, M. Harold Laughlin, Frank W. Booth, James R. Sowers, Jamal A. Ibdah, John P. Thyfault, e R. Scott Rector. “Treating NAFLD in OLETF Rats with Vigorous-Intensity Interval Exercise Training”. *Medicine and Science in Sports and Exercise* 47, n° 3 (março de 2015): 556–67.
- [48] Haczeyni, Fahrettin, Vanessa Barn, Auvro Mridha, Matthew Yeh, Emma Estevez, Mark Febbraio, Christopher Nolan, Kim Bell-Anderson, Narci Teoh, e Geoffrey Farrell. “Exercise improves adipose function and inflammation and ameliorates fatty liver disease in obese diabetic mice”. *Obesity (Silver Spring)*, 1º de agosto de 2015.