



Determination of the Elements of Soil Water Balance for Wheat (*Triticum aestivum* L.) Under Shallow Water Table

Salloom B. Salim¹, Luma S. Khudhair²

¹Department of Soil Sciences and Water Resources, College of Agriculture, Baghdad University, Baghdad, Iraq

²Extension Specialists, Ministry of Agriculture, Baghdad, Iraq

Email address:

salloom_s@yahoo.com (S. B. Salim)

To cite this article:

Salloom B. Salim, Luma S. Khudhair. Determination of the Elements of Soil Water Balance for Wheat (*Triticum aestivum* L.) Under Shallow Water Table. *International Journal of Applied Agricultural Sciences*. Vol. 1, No. 3, 2015, pp. 84-90. doi: 10.11648/j.ijaas.20150103.17

Abstract: To determine the elements of soil water balance equation during the growing season detailed description for calculating daily contribution rates to evapotranspiration of wheat (ET) from applied irrigation water (ET_i) and upward flux capillarity (ET_c), depth of applied irrigation water ($DAIW$), change in water storage (ΔS) and cumulative evapotranspiration (ET_{cum}) were algorithmed in this study. Irrigation water was applied to three different depths 30, 30-60 and 60 cm at three different depletion rates 50, 70 and 90% from plant available water. Wheat ET ranged from 428.49 to 522.12 mm. Contributions to ET from applied irrigation water ranged from 334.20 to 496.50 mm and increased with increasing irrigation depth. Contributions to ET from upward flux capillarity ranged from 25.61 to 96.59 mm and decreased with increasing irrigation depth. Contributions to ET from applied irrigation water decreased with increasing depletion rate whilst contributions to ET from upward flux capillarity increased with increasing depletion rates. Daily rate contribution to evapotranspiration from irrigation water ranged from 2.15 to 3.20 mm.d-1 and from capillary flux ranged from 0.16 to 0.61 mm.d-1.

Keywords: Shallow Water Table, Water Balance, Depletion Rate, Capillary Flux, Cumulative Evapotranspiration

1. Introduction

Surface water resources in Iraq of the Euphrates and the Tigris rivers declined sharply during the last two decades thus creating severe water scarcity in most irrigated areas and reducing Falkenmark water scarcity index from 2648 m³ per capita per year for the 20th century to 989 m³ per capita per year for the 21st century (Al-Shahrabli 2009, Saleh 2010). Limited water resources is the most effecting factor governing agriculture production and rising instability of food security. Understanding the magnitude and dynamics of applied irrigation water is crucial to development of technological options for sustainable management of available water resources (Strauss et al., 2010). The general soil water balance equation is given as:

$$I + C + P - \Delta S = ET + R + D \quad (1)$$

where I is irrigation, C is upward capillary flux, P is precipitation, ΔS is change in soil water storage, ET is evapotranspiration, R is runoff losses and D is deep drainage losses. Components ET , P , I and C are always positive, R and

D are always negative but ΔS can be positive or negative. Negative ΔS indicates depletion of soil-water storage. In situ assessment of the hydrological mass flux is essential in food security research (Owonubi et al, 1991) and irrigation water and land management (Brown et al., 2012). Methodologies, experimental techniques and modeling at the field scale facilitate applying the right amount of irrigation water at the right time that can based on soil, plant and/or atmospheric measurements and is commonly known as scientific irrigation scheduling (Lieb et al., 2002). Metering the irrigation system laterals at the field scale allows precise application of the deficit in plant available water between irrigations to the effective root zone (ERZ) and minimizing run off and deep drainage losses (Moiwo et al., 2011). In arid climates the contribution to plant needs from precipitation is negligible since rainfall intensity is low and mean annual precipitation is less than 100mm. Precipitation is always measured easily and precisely by rain gauge .

In most studies of soil water balance irrigation water is applied to replenish depleted plant available water to field capacity. Eiasu et al. (2006) applied four water depletion levels (20, 40, 60 and 80% of the plant available soil water) in the top 0.8 m root zone and noticed that plant roots

extracted most soil water from the top 0.4 m soil layer. The depletion rate at which water content is determined depends on plant and soil characteristics and its critical to water productivity (Jalota et al., 2006). The difference in stored water in ERZ requires the determination of initial water content before planting besides final water content after harvest. When initial water content in the ERZ exceeds assumed deficit then stored water is positive. Even gravimetric method for determining water content is basic but it represents the yard stick for other methods thus can be used to monitor change in water content during irrigation (Odhiambo and Bomke, 2007). When water is applied to satisfy plant evapotranspiration only then no percolation below the root zone occurs and deep drainage losses is set to zero. To insure zero deep drainage losses tensiometers can be installed at discrete depth intervals including at least the upper and the lower boundary of the ERZ and the vicinity of the capillary fringe of a shallow water table to determine soil water potential and flux direction (Saini and Ghildyal, 1977). Contribution from ground water to plant evapotranspiration can be determined according to Richard's equation (Richard's, 1931). In case of shallow water table the capillary flux is determined from the change in water content in the layer designated by tensiometer placement at the lower boundary of ERZ as its upper boundary and at the water table (capillary fringe) as its lower boundary. This study aimed to detailed description of the components of soil water balance equation for determining daily water use by growing crops under shallow water table and providing all equations necessary for calculations.

2. Materials and Methods

This study was performed in the experimental field of the College of Agriculture/ Baghdad University located at 33°20' longitude north, 44.°12' latitude east and 32m altitude during 2008-2009 wheat growing season in Iraq. Wheat (*Triticum aestivum* L.) seeds were planted at a rate of 130 kg. ha⁻¹ on 27/11/2008 at a depth of 5cm in grain drill rows and was harvested on 3/5/ 2009. Treatments included three levels of depletion rates 50, 70, and 90% from plant available water (PAW) and three irrigation depths [Depth of Effective Root Zone (ERZ)]; 30 cm from planting to harvest, 30 cm from planting to flowering then 60 cm from flowering to harvest and 60 cm from planting to harvest. The study was conducted in randomized complete block design with three replicates resulting in twenty seven 5m×5m experimental plots. Depth of applied irrigation water (DAIW) was calculated based on depletion rate from PAW in the ERZ. Water content changes was monitored gravimetrically using soil sampling tube (2.5 cm OD) extending to 90 cm depth. It is necessary to mention here that this study required predetermination of the soil-water retention relationship. Amount of irrigation water for the first irrigation was assumed equal to the volume of water required to increase water content for the plot from initial water content before first irrigation ($\theta_o = 0.205 \text{ cm}^3 \cdot \text{cm}^{-3}$) to field capacity ($\theta_{fc} = 0.40 \text{ cm}^3 \cdot \text{cm}^{-3}$) for 30 cm depth. During first irrigation all plots received the same amount of irrigation water at the same time. Amount of irrigation water for the following irrigations was set equal to the volume of water required to increase water content of the ERZ for the plot from initial water content to field capacity. Values of initial water content at which 50, 70, and 90% were depleted from plant available water (PAW) were calculated according to following formula:

$$\theta_o = \theta_{fc} - (\% \text{depletion}) / 100 \times PAW \quad (2)$$

or

$$\theta_o = \theta_{pwp} + (\% \text{available}) / 100 \times PAW \quad (3)$$

where θ_o , θ_{fc} , and θ_{pwp} are initial water content (volumetric water content measured at 24 hours before irrigation), volumetric water content at field capacity, volumetric water content at permanent wilting point ($0.181 \text{ cm}^3 \cdot \text{cm}^{-3}$) and $PAW = \theta_{fc} - \theta_o$ is the plant available water for the ERZ.

Irrigation water was pumped into PVC piping net (5 cm ID) with water meters measuring $\mp 0.001 \text{ m}^3$ volume difference fixed at the end of plot lateral. Each lateral equipped with control valve located directly before the water meter. Except first irrigation water was applied at different times depending on depletion rates and irrigation depths. Gauge tensiometers were installed at 30, 60 and 90 cm as shown in figure (1) for measuring matric potential (Ψ_m) which was calculated according to Hanks and Ashcroft formula (Hanks and Ashcroft, 1980) as follows:

$$\Psi_m = -10 \text{ cm} \times G_r + z_o \quad (4)$$

Where G_r is gauge reading and z_o is the vertical distance from the gauge to the center of ceramic cup of the tensiometer.

Total water potential was then calculated as the sum of matric and gravitational potentials to determining flow direction. Rate of ground water contribution to evapotranspiration was calculated for each irrigation according to Richard's equation from the change in soil water content in a layer designated by ERZ as its upper boundary (30 or 60 cm) and depth of tensiometer placement at 60 cm for 30 cm ERZ and at 90 cm for 60 cm ERZ.




Fig. 1. Schematic representation for tensiometer placement and theoretical basis for depleted water content difference ($\Delta\theta$) between two consecutive irrigations.

3. Results and Discussion

Values of the components of the soil water balance equation during the growing season of wheat are given in table (1) for all treatments. Different numbers of irrigations were obtained depending on irrigation depth and depletion rate. For the 30 cm irrigation depth and 50% depletion rate ten irrigations were applied which represents the highest irrigation frequency during 158 days of wheat growing season compared with other treatments. Initial water content before each irrigation was necessary to determine differences in stored water before irrigation and must be accounted for when calculating DAIW for each irrigation according to the following formula:

$$DAIW = (\theta_{fc} - \theta_i)z \quad (5)$$

where θ_{fc} and θ_i are volumetric water content at field capacity ($0.401 \text{ cm}^3.\text{cm}^{-3}$) and initial water content before irrigation respectively. Measured initial water content values for each irrigation are shown in column (5). According to equation (5), total DAIW during the growing season ($DAIW_t$) was calculated as follows:

$$DAIW_t = \sum_{i=1}^k DAIW_i \quad (6)$$

where i is a counter for irrigation number and k is the total number of irrigations. Laboratory determined values of θ_i from soil water retention.

Table 1. Measured and calculated values of the components of the soil water balance equation.

Depletion rate (percent from plant available water)	Irrigation depth, cm	Irrigation number	Day after planting	Initial water content $\text{cm}^3.\text{cm}^{-3}$	DAIW $\text{mm}.\text{irrigation}^{-1}$	Δs $\text{mm}.\text{irrigation}^{-1}$	ET_r $\text{mm}.\text{d}^{-1}$	ET_c $\text{mm}.\text{d}^{-1}$	ET_r $\text{mm}.\text{irrigation}^{-1}$	ET_c $\text{mm}.\text{irrigation}^{-1}$	ET_{in} $\text{mm}.\text{irrigation}^{-1}$
50%	0-30	1	0.00	0.205	58.80	25.50	2.22	0.43	33.30	6.45	39.75
		2	15.00	0.290	33.30	0.60	1.92	0.45	32.70	7.65	40.35
		3	32.00	0.292	32.70	-0.60	1.96	0.46	33.30	7.82	41.12
		4	49.00	0.290	33.30	-0.60	1.99	0.48	33.90	8.16	42.06
		5	66.00	0.288	33.90	0.60	2.08	0.48	33.30	7.68	40.98
		6	82.00	0.290	33.30	-1.20	2.16	0.50	34.50	8.00	42.50
		7	98.00	0.286	34.50	-0.60	2.34	0.55	35.10	8.25	43.35
		8	113.00	0.284	35.10	-1.20	2.79	0.65	36.30	8.45	44.75
		9	126.00	0.280	36.30	2.40	3.39	0.68	33.90	6.80	40.70
		10	136.00	0.288	33.90	-23.40	2.60	0.41	57.30	9.02	66.32
	Harvest	158.00	0.210	365.10	1.50 [†]	2.35 [†]	0.51 [†]	363.60 ^{††}	78.28 ^{††}	441.88 ^{††}	
	0-30 30 - 60	1	0.00	0.205	58.80	26.10	2.18	0.41	32.70	6.15	38.85
		2	15.00	0.292	32.70	-0.60	1.96	0.48	33.30	8.16	41.46
		3	32.00	0.290	33.30	0.00	1.96	0.48	33.30	8.16	41.46
		4	49.00	0.290	33.30	-0.60	1.99	0.53	33.90	9.01	42.91
		5	66.00	0.288	33.90	-2.40	2.27	0.58	36.30	9.28	45.58
		6	82.00	0.280	36.30	0.60	2.23	0.62	35.70	9.92	45.62
		7	98.00	0.282	35.70	-1.20	2.46	0.68	36.90	10.20	47.10
		8	113.00	0.278	73.80	2.40	3.40	0.23	71.40	4.83	76.23
		9	134.00	0.282	71.40	-31.20	4.28	0.18	102.60	11.28	113.88
		Harvest	158.00	0.230	409.20	-6.90 [†]	2.53 [†]	0.47 [†]	416.10 ^{††}	76.99 ^{††}	493.09 ^{††}
	0 - 60	1	0.00	0.205	58.80	23.70	1.76	0.09	35.10	1.80	36.90
		2	20.00	0.284	70.20	5.40	3.41	0.12	64.80	2.28	67.08
		3	39.00	0.302	59.40	-0.60	2.73	0.15	60.00	3.30	63.30
		4	61.00	0.300	60.60	-1.80	2.97	0.19	62.40	3.99	66.39
		5	82.00	0.294	64.20	1.80	3.12	0.20	62.40	4.00	66.41
		6	102.00	0.300	60.60	-3.00	3.53	0.21	63.60	3.78	67.38
7		120.00	0.290	66.60	3.00	3.35	0.21	63.60	3.99	67.59	
8		139.00	0.300	60.60	-24.00	4.45	0.13	84.60	2.47	87.07	
Harvest		158.00	0.220	501.00	4.50 [†]	3.16 [†]	0.16 [†]	496.50 ^{††}	25.61 ^{††}	522.12 ^{††}	
70%		0-30	1	0.00	0.205	58.80	12.60	1.93	0.49	46.20	11.76
	2		24.00	0.247	46.20	-2.10	2.68	0.53	48.30	9.54	57.84
	3		42.00	0.240	48.30	-1.20	1.98	0.56	49.50	14.00	63.50
	4		67.00	0.236	49.50	1.80	2.17	0.57	47.70	12.54	60.24
	5		89.00	0.242	47.70	0.90	2.23	0.61	46.80	12.81	59.61
	6		110.00	0.245	46.80	1.50	2.38	0.63	45.30	11.97	57.27
	7		129.00	0.250	45.30	-12.00	1.98	0.58	57.30	16.82	74.12
	Harvest	158.00	0.210	342.60	1.50 [†]	2.19 [†]	0.57 [†]	341.10 ^{††}	89.44 ^{††}	430.54 ^{††}	
	0.30 0-60	1	0.00	0.205	58.80	14.10	2.13	0.49	44.70	10.29	54.99

Depletion rate (percent from plant available water)	Irrigation depth, cm	Irrigation number	Day after planting	Initial water content $\text{cm}^3.\text{cm}^{-3}$	DAIW $\text{mm}.\text{irrigation}^{-1}$	ΔS $\text{mm}.\text{irrigation}^{-1}$	ET_r $\text{mm}.\text{d}^{-1}$	ET_c $\text{mm}.\text{d}^{-1}$	ET_{ir} $\text{mm}.\text{irrigation}^{-1}$	ET_{is} $\text{mm}.\text{Irrigation}^{-1}$	ET_{ih} $\text{mm}.\text{irrigation}^{-1}$
90%	0-60	2	21.00	0.252	44.70	-1.80	1.72	0.53	46.50	14.31	60.81
		3	48.00	0.246	46.50	-3.60	1.79	0.56	50.10	15.68	65.78
		4	76.00	0.234	50.10	1.20	1.81	0.57	48.90	15.39	64.29
		5	103.00	0.238	97.80	1.20	3.22	0.24	96.60	7.20	103.80
		6	133.00	0.240	96.60	-18.00	4.58	0.17	114.60	4.25	118.85
		Harvest	158.00	0.210	394.50	-6.90 [†]	2.54 [†]	0.43 [†]	401.40 ^{††}	67.12 ^{††}	468.52 ^{††}
		1	0.00	0.205	58.80	32.40	1.10	0.09	26.40	2.16	28.56
	2	24.00	0.259	85.20	-0.60	3.30	0.15	85.80	3.90	89.70	
	3	50.00	0.258	85.80	0.60	3.16	0.23	85.20	6.21	91.41	
	4	77.00	0.259	85.20	-1.20	3.20	0.28	86.40	7.56	93.96	
	5	104.00	0.257	86.40	-1.80	3.27	0.28	88.20	7.56	95.76	
	6	131.00	0.254	88.20	-24.60	4.18	0.19	112.80	5.13	117.93	
	Harvest	158.00	0.213	489.60	4.80 [†]	3.03 [†]	0.20 [†]	484.80 ^{††}	32.52 ^{††}	517.32 ^{††}	
	1	0.00	0.205	58.80	0.30	2.34	0.55	58.50	13.75	72.25	
	2	25.00	0.206	58.50	1.80	2.18	0.57	56.70	14.82	71.52	
	3	51.00	0.212	56.70	-0.90	2.30	0.62	57.60	15.38	72.98	
	4	76.00	0.209	57.60	0.90	1.62	0.67	56.70	23.45	77.85	
	5	111.00	0.212	56.70	-1.80	2.44	0.68	58.50	16.32	74.82	
	6	135.00	0.206	58.50	12.30	2.01	0.56	46.20	12.88	59.08	
	Harvest	158.00	0.247	346.80	12.60 [†]	2.15 [†]	0.61 [†]	334.20 ^{††}	96.59 ^{††}	428.49 ^{††}	
	1	0.00	0.205	58.80	0.30	2.44	0.53	58.50	12.72	71.22	
	2	24.00	0.206	58.50	0.30	2.16	0.62	58.20	16.74	74.94	
	3	51.00	0.207	58.20	-1.20	2.20	0.75	59.40	20.25	79.65	
	4	78.00	0.203	59.40	0.30	2.69	0.85	59.10	18.70	77.80	
5	100.00	0.204	59.10	2.40	1.96	0.31	56.70	8.99	65.69		
6	129.00	0.212	113.40	-1.20	3.95	0.27	114.60	7.83	122.43		
Harvest	158.00	0.210	407.40	0.90 [†]	2.56 [†]	0.56 [†]	406.50 ^{††}	85.23 ^{††}	491.73 ^{††}		
1	0.00	0.205	58.80	1.20	1.99	0.15	57.60	4.35	61.95		
2	29.00	0.207	116.40	1.20	3.03	0.21	115.20	7.98	123.18		
3	67.00	0.209	115.20	-2.40	3.09	0.26	117.60	9.88	127.48		
4	105.00	0.205	117.60	1.20	3.06	0.26	116.40	9.88	126.28		
5	143.00	0.207	116.40	43.80	4.84	0.26	72.60	3.90	76.50		
Harvest	158.00	0.280	524.40	45.00 [†]	3.20 [†]	0.23 [†]	479.40 ^{††}	35.99 ^{††}	515.39 ^{††}		

† Average of data in column

†† Sum of data in column relationship

Relationships were 0.291, 0.247 and 0.203 $\text{cm}^3.\text{cm}^{-3}$ for the 50, 70 and 90% depletion rates during 158 days of wheat growing season. However actual θ_i values in column (5) does not necessarily equal laboratory estimated θ_i at the assumed depletion rates as shown in table (1) due to soil heterogeneity.

Value of θ_i before first irrigation was 0.205 $\text{cm}^3.\text{cm}^{-3}$ resulting in the highest DAIW value (58.8 mm) for the 30 cm depth and 50% depletion rate. DAIW for the first irrigation was assigned to be equal to required volume of applied irrigation water divided by plot area (25m²) to replenish initial water content of the 30 cm ERZ to the field capacity (0.401 $\text{cm}^3.\text{cm}^{-3}$). For the remaining irrigations DAIW was calculated on the same basis for different treatments based on irrigation depth and θ_i values. When θ_i value at irrigation was greater than water content at 50% depletion (0.291 $\text{cm}^3.\text{cm}^{-3}$) then water storage (ΔS) increases and ΔS was positive. High ΔS value (25.5 mm) was obtained during first

irrigation to compensate for the deficit in θ_i values between first and second irrigations (0.205 and 0.290 $\text{cm}^3.\text{cm}^{-3}$ respectively). The lowest ΔS values was obtained at harvest since soil profile was undergoing drying process under dry and hot conditions for 22 days after last irrigation. DAIW values were calculated according to depletion rate, irrigation depth and θ_i values.

The piped - delivery irrigation system was equipped with water meters at the end of plot laterals measuring $\pm 0.001 \text{ m}^3$ volume difference that facilitated applying exactly the right DAIW and eliminating run off losses during water application to experimental plots. Drainage losses were set equal to zero since total potential head at 30 cm was always lower than total potential head at 60 cm depth resulting in continuous upward water movement from the ground water (90 cm) towards ERZ (30 and or 60 cm). Daily contribution rate to plant evapotranspiration (ET_r) which was accounted for by irrigation water was calculated according to the following formula:

$$ET_r = (DAIW - \Delta S) / T \quad (7)$$

where T is time in days between two consecutive irrigations. Values of ET_r are given in column (8). Contribution from DAIW to plant evapotranspiration for each irrigation (ET_{ir}) was calculated according to the following formula:

$$ET_{ir} = ET_r \times T \quad (8)$$

and is given in column (10). Total contribution from DAIW to plant evapotranspiration (ET_{it}) is the sum of ET_{ir} 's for all irrigations and was calculated according to the following formula:

$$ET_{it} = \sum_{i=1}^k ET_{ir} \quad (9)$$

where i is a counter for irrigation number. ET_{it} value for this treatment was 363.60 mm during the growing season.

High ET_r rate, 2.22 mm.d⁻¹ during first irrigation is attributed to low canopy and high evaporation potential. Except last irrigation where soil profile was undergoing drying process for harvest a consistent increase in ET_r values occurred with increasing irrigation number during the growing season reflecting increasing plant demands of water and nutrients for growth development.

Rate of daily contribution to evapotranspiration from ground water expressed as upward flux (ET_c) for a soil layer designated by tensiometer placement was calculated according to Richards' equation as follows:

$$ET_c = \int_{z_n}^{z_{n+1}} \frac{\partial \theta}{\partial T} dz \quad (10)$$

where z_n and z_{n+1} were the upper and lower boundaries of the 30 cm layer below ERZ. Boundaries of this layer were determined by tensiometer placement either at 30 and 60 or at 60 and 90 cm for 30 and 60 cm ERZ respectively. $\partial \theta$ is the difference in volumetric water content which was calculated for the 30-60 cm layer according to following formula:

$$\partial \theta = (\theta_{60} - \theta_{30})|_{T_2} - (\theta_{60} - \theta_{30})|_{T_1} \quad (11)$$

where 30 and 60 represent the boundaries of the layer of the upward flux for the 30 cm ERZ. For the 60-90 cm layer $\partial \theta$ was calculated according to following formula:

$$\partial \theta = (\theta_{90} - \theta_{60})|_{T_2} - (\theta_{90} - \theta_{60})|_{T_1} \quad (12)$$

where 60 and 90 represent the boundaries of the layer of the upward flux for the 60 cm ERZ. ∂T is the time difference in days between two consecutive irrigations.

Daily ET_c values are given in column (9). Contribution from upward flux to evapotranspiration for each irrigation

(ET_{ic}) was calculated according to the following formula:

$$ET_{ic} = ET_c \times T \quad (13)$$

and is given in column (11). Total contribution from upward flux to plant evapotranspiration (ET_{ct}) is the sum of ET_{ic} 's for all irrigations and was calculated according to the following formula:

$$ET_{ct} = \sum_{i=1}^k ET_{ic} \quad (14)$$

ET_{ct} value for this treatment was 78.28 mm during the growing season which represents 21.5% of the ET_{it} .

Actual rate of evapotranspiration for each irrigation, ET_{ia} mm.d⁻¹, is defined to be the sum of ET_{ir} and ET_{ic} and was calculated according to the following formula:

$$ET_{ia} = ET_{ir} + ET_{ic} \quad (15)$$

and is given in column (12). ET_{ia} value for the first irrigation was 39.75 mm. Cumulative evapotranspiration (ET_{cum}) which represents actual evapotranspiration for the whole season was defined as the sum of ET_{ia} 's according to following formula:

$$ET_{cum} = \sum_{k=1}^n ET_{ia} \quad (16)$$

Value of ET_{cum} for this treatment was 441.88 mm. Measured data are given in columns 2 thru 5 while calculated values of the components of the water balance equation are given in columns 6 thru 12 as shown in table (1).

Only nine irrigations were applied when irrigation depth was switched from 30 cm to 60 cm at flowering. Components of the water balance for this treatment is given in table (1). Increasing irrigation depth from 30 to 60 cm increased DAIW by two folds, increased irrigation intervals of the last two irrigations as a result of increasing depth of ERZ from 30 to 69 cm, decreased ET_c and increased ET_r . Compared with 30 cm irrigation depth almost the same values were obtained for θ_i , ET_c , ET_{ir} , ET_{ia} and ET_{cum} during the first seven irrigations. Increasing irrigation depth to 60 cm along the growing season decreased number of irrigations and ET_c and increased DAIW, ET_{ir} and ET_{cum} as shown in table (1).

Increasing depletion rate to 70% decreased number of irrigations to seven for the 30 cm depth and to six for the 30-60 and 60 cm depths and θ_i value to 0.247 cm³.cm⁻³ which in turn increased DAIW as shown in table (1). Switching irrigation depth from 30 to 60 cm increased ET_r , ET_{cum} , and DAIW but significantly decreased ET_c . Averages of the ET_c were 0.57, 0.43 and 0.20 mm.d⁻¹ for the 30, 30-60 and 60 cm depths respectively. Decreasing average ET_c values with

increasing irrigation depth is best explained according to the Darcy's law where flux term between two points is proportional to the difference in total water potential divided by the distance (vertical in this study) between the two points at given water content or suction head values. Values of total potential head for 30, 60 and 90 cm after 24 hours following irrigation ranged from -105 to -75 cm H₂O as shown in fig. (2).




Fig. 2. Total potential head for the 30, 60 and 90 cm measured at 70% depletion rate and 24 hours after irrigations for the 30 cm irrigation depth.

Slight differences in total potential head values between different depths resulted in small hydraulic head gradient values. Fig. (2) also shows that the total potential head at the 90 cm depth is higher than total potential at the 60 cm depth for all irrigations. As a result the gradient values were around unity and the upward flux towards 60 cm decreased significantly. Pronounced differences in total potential values was developed between 30 and 60 cm depths as time proceeded after irrigation due to drying of the surface layer (0-30 cm) by evaporation from the soil surface and plant transpiration as shown in fig. (3).




Fig. 3. Total potential head for the 30, 60 and 90 cm measured at 70% depletion rate and 24 hours before irrigations for the 30 cm irrigation depth.

Values of total water potential at the 60 cm depth were almost the same during the growing season since water table level ranged between 70 and 90 cm. These findings agree with the findings of Salim and Rasheed (2012) who studied the movement of zero flux plane during irrigation under shallow water table condition. Compared with the distribution of total potential head shown in figure (3) different distribution patterns were obtained when the water table was deep enough from ERZ (Magdi et al. 2003, Joshi 1997).

Due to differences in ΔS values between irrigations ET_{it} value (401.40 mm) exceeded $DAIW_i$ (394.50 mm) for the 30-60 cm depth by 6.90 mm which means that the depletion rate exceeded 70% from plant available water (see table 1) even average ET_c value of the 30 cm depth (0.57 mm.d^{-1}) was greater than ET_c value at the 30-60 cm depths (0.43 mm.d^{-1}). Increasing irrigation depth increased ET_{it} to 341.40 mm, 401.40 mm and 484.80 mm for the 30 cm, 30-60 cm and from 30-60 cm depths respectively.

Depleting 90% from available water content decreased number of irrigations and increased $DAIW$. While ET_c values decreased, ET_i values increased with increasing irrigation depth. Highest ET_c contribution to actual evapotranspiration, 96.59 mm, occurred under 30 cm depth and decreased by 62.7% for the 60 cm depth treatments. Decreasing average ET_c values with increasing irrigation depth is attributed to decreasing hydraulic head gradient between 60 and 90 cm compared with the gradient values between 30 and 60 as shown in fig. (4) for the 90% depletion rate.




Fig. 4. Hydraulic head gradient of the layers 30 cm, 30-60 cm and 60-90 cm for 30, 30-60, and 60 cm irrigation depths respectively at 90% depletion rate.

The highest $DAIW$, 524.40 mm, was applied for the 60 cm depth resulting in the highest positive ΔS value (45.00 mm). It seems that the fifth irrigation was not necessary for plant growth since it was applied before 15 days from harvest where the grains were at the drying stage.

The highest ET_i contribution to actual evapotranspiration, 496.50mm occurred under 50% depletion and 60 cm depth

and decreased by 27.2% for the 30 cm depth. The highest ET_{cum} value (512.22 mm) obtained under 50% depletion rate and 60 cm irrigation depth. However differences in evapotranspiration values doesn't necessarily respond to differences in growth factors and yield of wheat.

4. Conclusions

Under water scarcity circumstances and shallow water table condition, a precise assessment of flow processes occurring during the water distribution and redistribution in the vadose zone is required to determine daily contribution rates to plant evapotranspiration from the soil water balance equation and to minimize water losses and maximize capillary flux contributions which in turns increase water productivity. Results of this study can be used as a field practice for maximizing contribution from shallow water table to crop evapotranspiration when considering depletion rate and irrigation depth.

References

- [1] Al-shahrabli. Q. 2009. Surface Water Resources in Iraq: Current and Future Scenarios. Iraqi Soil Salinity and Water Management Conference. July 15th- 17th 2009, Alrashid Hotel. Baghdad, Iraq.
- [2] Brown, I., Dunn, S., Matthews, K., Poggio, L., Sample, J. and Miller, D. 2012. Mapping of water supply-demand deficits with climate change in Scotland: land use implications, CREW report 2011/CRW006. Available online at: www.crew.ac.uk/publications/water-supply-demand-balance-and-climate-change.
- [3] Eiasu, B. K., J.M. Steyn and P. Soundy. 2006. Rose-scented geranium (*Pelargonium capitatum* × *P. radens*) growth and essential oil yield response to different soil water depletion regimes. *Agricultural Water Management*, 96(6): 991-1000.
- [4] Hanks, R. J. and G. L. Ashcroft. 1980. *Applied Soil Physics*. Advanced series in Agricultural Sciences8. Sprinkler-Verlag. Berlin Heidelberg New York Tokyo.
- [5] Jalota, S. K., A. Sood, G.B.S. Chahal and B.U. Choudhury. 2006. Crop water productivity of cotton (*Gossypium hirsutum* L.)–wheat (*Triticum aestivum* L.) system as influenced by deficit irrigation, soil texture and precipitation . *Agricultural Water Management*, 84(2): 137-146.
- [6] Joshi, B. 1997. Estimation of diffuse vadose zone soil-water flux in a semi- arid region. PhD thesis, Department of Agriculture and Bioresources Engineering . University of Saskatchewan, Canada.
- [7] Khalil, M., M. Sakai, M. Mizoguchi and T. Miyazaki. 2003. Current and Prospective Applications of Zero Flux Plane (ZFP) Method. *J. Jpn. Soc. Soil Phys. J. Jpn . Soc. Soil Phys.* 95: 75-90.
- [8] Leib, B.G., M. Hattendorf, T. Elliott and G. Matthews. 2002. Adoption and adaptation of scientific irrigation scheduling: trends from Washington, USA as of 1998. *Agricultural Water Management* 55: 105-120.
- [9] Moiwo J. P., F. Tao and W. lub. 2011. Estimating soil moisture storage change using quasi-terrestrial water. *Agriculture Water management*. 102(1)25-34.
- [10] Odhiambo, J. J. O. and A.A. Bomke. 2007. implications for cover crop management in south coastal British Columbia *Agricultural Water Management*, Volume 88, Issues 1–3, 16 March 2007, Pages 92-98
- [11] OJuana Paul Moiwo J. P.and, F.Tao and W. Lu. 2011. Estimating soil moisture storage change using quasi-terrestrialwater balance method. *Agricultural Water Management* 102(1): 25-34.
- [12] Owonubi , J. J. and S. Abduimumin. 1991. Review of soil water balance studies in the Sudano-Sahelian zone of Nigeria. *Proceedings of the Niamey Workshop, February 1991 . IAHS* 1991.
- [13] Quality 40:1652-1660.
- [14] Richards, L.A. 1931. Capillary conduction of liquids through porous mediums. *Physics* 1(5):318-333.
- [15] Saini, B. C. and B.P. Ghildyal. 1977. Seasonal water use by winter wheat grown under shallow water table conditions *Original Research Article Agricultural Water Management*, Volume 1, Issue 3, November 1977, Pages 263-276.
- [16] Saleh, D.K. 2010, Stream gage descriptions and stream flow statistics for sites in the Tigris River and Euphrates River Basins, Iraq: U.S. Geological Survey Data Series 540, 146 p.
- [17] Salim S. B. and T. L. Rasheed. 2013. Water balance in cultivated and uncultivated soil. *J. Sci. Tecnology*. 4(3):85-93.
- [18] Strauss, C., T. Harter and M. Radke. 2011. Effects of pH and Manure on Transport of Sulfonamide Antibiotics in Soil. *Journal of Environmental Quality* 40:1652-1660.