

Research Article

New Theorems Solving Fifth Degree Polynomial Equation in Complete Forms by Proposing New Five Roots Composed of Radical Expressions

Yassine Larbaoui* 

Department of Electrical Engineering, University Hassan 1er, Settati, Morocco

Abstract

This paper is presenting new theorems and formulas to solve fifth degree polynomial equation in general forms by proposing five formulary solutions that we can calculate nearly simultaneously. The proposed roots for fifth degree polynomial in this paper are composed of radical expressions distributed in a specific architecture that allows them to neutralize each other during multiplication by eliminating radicality and reducing degrees of terms. The proposed architecture for each solution of quantic polynomial is composed of at least six distributed terms, where each term is a multiplication of a pair of radical expressions, whereas multiplications among proposed terms is allowing the reduction of degrees in order to obtain a quartic form at maximum. Furthermore, the proposed architectures of radical roots in this paper are allowing also to solve polynomial equations with degrees higher than five by reducing their expressions. As a result, this paper is presenting two new theorems to solve quantic equation in general forms, where one theorem is eliminating the expression of fourth degree then reducing the whole fifth degree equation into a simplified form, whereas the other new theorem is solving the quantic equation without eliminating the expression of fourth degree. All proposed theorems in this paper are built on a scaling logic concretized by engineering the structure of solutions then calculating the precise formulas of roots for equations, which allow to calculate all roots nearly simultaneously and forward the used engineering methodology to solve polynomial equations with degrees higher than five.

Keywords

Engineered Root Architectures, New Five Solutions, New Theorems, Roots Nearly Simultaneous Calculation, Solving Fifth Degree Polynomial

1. Introduction

Developing the expressions of formulary solutions to solve n^{th} degree polynomial equations has been an enigmatic problem over hundreds of years to mathematicians, where many research attempts concluded to the impossibility of elaborating unified solution formulas for equations with degrees equal or

higher than fifth degree by using radicals. However, finding breakthroughs to structure universal radical solutions is a living hope to solve quantic equations and above.

Solving quantic polynomial equations in general forms has been center of focus over centuries, because they presented a

*Corresponding author: y.larbaoui@uhp.ac.ma (Yassine Larbaoui), yassine.larbaoui.uh1@gmail.com (Yassine Larbaoui)

Received: 20 February 2024; **Accepted:** 6 March 2024; **Published:** 2 April 2024



Copyright: © The Author(s), 2024. Published by Science Publishing Group. This is an **Open Access** article, distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

main problematic axe in many fields. Many adopted approaches to solve quantic equations majorly concluded to the impossibility of elaborating universal formulary expressions by using radicals as solutions, whereas others relied on treating resolvable quantic equations or reducing specific forms of fifth degree polynomials to other expressions with inferior degrees. However, developing new approaches where the scientific methodology is the principal goal, rather than aiming the discovery of specific expressions as solutions, may bring new results.

Even though solving quantic equations by using unified expressions is the center of focus in our research, developing new formulary solutions for fourth degree polynomial equations in general forms was always appealing to attention, because discovering new solution expressions for quartic equations might bring a breakthrough to express universal solutions for fifth degree polynomials by using radical expressions.

This article presents specific results of a large research in mathematic science by proposing five new formulary solutions for any fifth-degree polynomial equation in general form with real coefficients. The developed formulary solutions for fifth degree polynomials are based on new formulary expressions as solutions for fourth degree equations, which are also proposed in this paper within a new theorem.

This paper attacks the problem of solving polynomial equations of n^{th} degree, where $n=4$ and $n=5$, with a detailed manner basing on extendable logic, which enables to structure new theorems by scaling from solving quartic equations upward solving quantic equations by using radicals.

Lodovico Ferrari discovered a principal solution for quartic polynomial equations in the sixteen-century (1540), but since his discovery required having a breakthrough to solve cubic equations, which was not published yet by Gerolamo Cardano, Lodovico Ferrari could not publish his elaborated solution for fourth degree polynomials immediately. In the book *Ars Magna* [1], which was published by Ferrari's mentor (Gerolamo Cardano), the discovered quartic solution by Lodovico Ferrari was published along with the cubic solution.

The proposed cubic solution by Cardano [2] for third degree polynomials under the form $x^3 + px + q = 0$ helps Ferrari's quartic solution to solve fourth degree polynomial equations by reducing them to the second degree. However, the elaborated solution by Ferrari does not directly help to, properly; define the four roots of any fourth degree polynomial simultaneously. The proposed method by Cardano for cubic equations was also the base to solve particular forms of n^{th} degree polynomials; such as by giving radical expressions under the form $\sqrt{na} + \sqrt{b} + \sqrt{na - \sqrt{b}}$ for $n=2,3,4, \dots$, etc. [3].

There are also other proposed roots and elaborated methods to solve quartic equations such as Galois's method [4], Lagrange's method [5], algebraic geometry [6], Descartes's method [7] and Euler's solution [8].

Through the history of mathematics, the proposed solution

by Cardano for cubic equations was always essential as base ground for further research about solving polynomial equations of fourth degree and above [9, 10], such as trying to prove that quantic equations do not accept quadratic expressions as potential solutions, which is further discussed in [11-14]. Nevertheless, in the results of this paper, the proposed five solutions for fifth degree polynomial equations are structured by using the proposed four roots for quartic equations in Theorem 1, where we include the expressions of quadratic roots as subparts of each solution.

There are also other recent researches dedicated to solve fourth degree polynomials where the used methods are based on reducing the expressions of equations [15-17], whereas other researches are relying on using algorithms and numerical analysis to find the roots of polynomial equations with degrees higher than four [17, 18].

In the published paper by Tschirnhaus [19], he proposed an innovative method to solve polynomial equation $P_n(x)$ of n^{th} degree by transforming it into a reduced expression $Q_n(y)$ with fewer terms by extending the idea of Descartes; in which a polynomial of n^{th} degree is reducible by removing its term in degree $(n - 1)$. The projection of this method on quantic equations is more detailed in [20].

There are other papers treating resolvable quantic polynomials basing on radical expressions [21, 22], where the description of specific characteristics which determine whether a fifth-degree polynomial may accept roots with radical expressions or not [24, 25].

The advantage of this paper is proposing five formulary solutions for any form of fifth degree polynomial equations with real coefficients, and thereby having the possibility to calculate the five roots of any quantic equation nearly simultaneously. The developed solutions for fifth degree polynomial equations in this paper are based on new solutions for fourth degree polynomials in general forms, where we propose four formulary expressions as roots for any quartic equation.

As a result, this paper is presenting two new theorems to solve fifth degree polynomial equations in general forms where the possibility of calculating all five roots nearly simultaneously. The first new theorem is solving quantic equation after eliminating the quartic term, whereas the second theorem is solving the quantic equation directly without eliminating the quartic term.

The two new theorems in this paper are based on an engineered methodology to structure radical solutions for polynomial equations of n^{th} degree. We developed this structuring methodology to solve quartic equations [23], then we forwarded the methodology in this paper by expanding the engineered structures of roots to solve fifth degree polynomial equations in complete forms.

Because the presented content in this paper is original, and there are many new proposed mathematical expressions and new formulas related in a scaling manner basing on extendable logic; every expressed formula will be proved mathe-

mathematically and used to build the rest of content, and we will go through them by logical analysis and deduction basing on structured development.

This paper is structured as follow: section 2 where we propose new solutions for fourth degree polynomial equation in general form, section 3 where we propose new five formulary solutions for fifth degree polynomial equation in general form, and section 4 for conclusion.

2. New Four Solutions for Fourth Degree Polynomial Equation in General Form

This section presents new unified formulary solutions for fourth degree polynomial equations in general form (Eq. 1) along with their proof. The used logic and expressions in the proof of this section is essential to prove the proposed theorems in section 3.

2.1. Introduction on First Proposed Theorem

In this subsection, we propose a new theorem to resolve fourth degree polynomial equations that may be presented as shown in (Eq. 1).

The expressions of proposed solutions are dependent on the value of $\left(8\frac{b^3}{a^3} - 32\frac{cb}{a^2} + 64\frac{d}{a}\right)$.

We are proposing four solutions for $\left(8\frac{b^3}{a^3} - 32\frac{cb}{a^2} + 64\frac{d}{a}\right) < 0$, four solutions for $\left(8\frac{b^3}{a^3} - 32\frac{cb}{a^2} + 64\frac{d}{a}\right) > 0$ and four solutions for $\left(8\frac{b^3}{a^3} - 32\frac{cb}{a^2} + 64\frac{d}{a}\right) = 0$.

The proposed solutions for $\left(8\frac{b^3}{a^3} - 32\frac{cb}{a^2} + 64\frac{d}{a}\right) < 0$, $\left(8\frac{b^3}{a^3} - 32\frac{cb}{a^2} + 64\frac{d}{a}\right) > 0$ and for $\left(8\frac{b^3}{a^3} - 32\frac{cb}{a^2} + 64\frac{d}{a}\right) = 0$ are expressed by using $y_{0,1}$ in (Eq. 17), P in (Eq. 6) and Q in (Eq. 6).

The proof of this theorem is presented in an independent subsection, because it integrates long mathematical expressions.

2.2. Theorem 1

A fourth degree polynomial equation under the expressed form in (Eq. 1), where coefficients belong to the group of

$$y^4 + y^2 \left[-6\left(\frac{b}{a}\right)^2 + 16\frac{c}{a} \right] + y \left[8\left(\frac{b}{a}\right)^3 - 32\frac{cb}{a^2} + 64\frac{d}{a} \right] - 3\left(\frac{b}{a}\right)^4 + 16\frac{cb^2}{a^3} - 64\frac{db}{a^2} + 256\frac{e}{a} = 0 \quad (4)$$

To simplify the expression of polynomial equation in (Eq. 4), we replace the expressions of used coefficients as shown in (Eq. 5) where the values of those coefficients are defined in (Eq. 6).

$$y^4 + Py^2 + Qy + R = 0 \quad (5)$$

$$P = -6\left(\frac{b}{a}\right)^2 + 16\frac{c}{a}; Q = 8\left(\frac{b}{a}\right)^3 - 32\frac{cb}{a^2} + 64\frac{d}{a}; R = -3\left(\frac{b}{a}\right)^4 + 16\frac{cb^2}{a^3} - 64\frac{db}{a^2} + 256\frac{e}{a} \quad (6)$$

numbers \mathbb{R} and $\neq 0$, has four solutions.

$$ax^4 + bx^3 + cx^2 + dx + e = 0 \text{ with } a \neq 0 \quad (1)$$

If $\left(8\frac{b^3}{a^3} - 32\frac{cb}{a^2} + 64\frac{d}{a}\right) < 0$, and by using the expressions of $y_{0,1}$ in (Eq. 17), P in (Eq. 6) and Q in (Eq. 6):

1. Solution 1: $S_{1,1}$ in expression (Eq. 35);
2. Solution 2: $S_{1,2}$ in expression (Eq. 36);
3. Solution 3: $S_{1,3}$ in expression (Eq. 37);
4. Solution 4: $S_{1,4}$ in expression (Eq. 38).

If $\left(8\frac{b^3}{a^3} - 32\frac{cb}{a^2} + 64\frac{d}{a}\right) > 0$, and by using the expressions of $y_{0,1}$ in (Eq. 17), P in (Eq. 6) and Q in (Eq. 6):

1. Solution 1: $S_{2,1}$ in expression (Eq. 39);
2. Solution 2: $S_{2,2}$ in expression (Eq. 40);
3. Solution 3: $S_{2,3}$ in expression (Eq. 41);
4. Solution 4: $S_{2,4}$ in expression (Eq. 42).

If $\left(8\frac{b^3}{a^3} - 32\frac{cb}{a^2} + 64\frac{d}{a}\right) = 0$, and by using the expressions of $y_{0,1}$ in (Eq. 28), P in (Eq. 6) and Q in (Eq. 6):

1. Solution 1: $S_{3,1}$ in expression (Eq. 43);
2. Solution 2: $S_{3,2}$ in expression (Eq. 44);
3. Solution 3: $S_{3,3}$ in expression (Eq. 45);
4. Solution 4: $S_{3,4}$ in expression (Eq. 46).

2.3. Proof of Theorem 1

By dividing the polynomial (Eq. 1) on the coefficient a , we have the next form:

$$x^4 + \frac{b}{a}x^3 + \frac{c}{a}x^2 + \frac{d}{a}x + \frac{e}{a} = 0 \text{ with } a \neq 0 \quad (2)$$

We suppose that x is expressed as shown in (Eq. 3):

$$x = \frac{\left(\frac{-b}{a} + y\right)}{4} \quad (3)$$

We replace x with supposed expression in (Eq. 3) to reduce the form of presented polynomial in (Eq. 2). Thereby, we have the presented expression in (Eq. 4).

To solve the shown polynomial equation in (Eq. 5), we propose new expressions for the variable y ; expressions (Eq. 7) and (Eq. 8):

$$\text{for } Q \leq 0: y = \sqrt{y_0} + \sqrt{y_1} + \sqrt{y_2} \quad (7)$$

$$\text{for } Q \geq 0: y = -\sqrt{y_0} - \sqrt{y_1} - \sqrt{y_2} \quad (8)$$

We propose the expressions (Eq. 9) and (Eq. 10) for y_1 and y_2 successively, in condition of $y_0 \neq 0$. These expressions of y_1 and y_2 are based on quadratic solutions.

$$y_1 = -\frac{\frac{P}{2} + y_0}{2} + \sqrt{\left(\frac{\frac{P}{2} + y_0}{2}\right)^2 - \frac{Q^2}{64y_0}} \quad (9)$$

$$y_2 = -\frac{\frac{P}{2} + y_0}{2} - \sqrt{\left(\frac{\frac{P}{2} + y_0}{2}\right)^2 - \frac{Q^2}{64y_0}} \quad (10)$$

To reduce the expression of equation (Eq. 5) and find a way to solve it, we propose the following expressions for the coefficients P and Q :

$$-2[y_0^2 + y_1^2 + y_2^2] = P \quad (11)$$

$$\text{For } Q \leq 0: -8\sqrt{y_0}\sqrt{y_1}\sqrt{y_2} = Q \quad (12)$$

$$\text{For } Q \geq 0: 8\sqrt{y_0}\sqrt{y_1}\sqrt{y_2} = Q \quad (13)$$

In the following calculation, we replace the variable y with the expression (Eq. 7) where we suppose $Q < 0$, and we replace P and Q with their shown expressions in (Eq. 11) and (Eq. 12):

$$\begin{aligned} y^4 + Py^2 + Qy + R &= -[\sqrt{y_0^4} + \sqrt{y_1^4} + \sqrt{y_2^4}] + 2[\sqrt{y_0^2}\sqrt{y_1^2} + \sqrt{y_0^2}\sqrt{y_2^2} + \sqrt{y_1^2}\sqrt{y_2^2}] + R \\ &= -[\sqrt{y_0^2} + \sqrt{y_1^2} + \sqrt{y_2^2}]^2 + 4[\sqrt{y_0^2}\sqrt{y_1^2} + \sqrt{y_0^2}\sqrt{y_2^2} + \sqrt{y_1^2}\sqrt{y_2^2}] + R \\ &= 4\left[-y_0\left(\frac{P}{2} + y_0\right) + \frac{Q^2}{64y_0}\right] + R - \frac{P^2}{4} = 0 \\ -[\sqrt{y_0^2} + \sqrt{y_1^2} + \sqrt{y_2^2}]^2 + 4[\sqrt{y_0^2}\sqrt{y_1^2} + \sqrt{y_0^2}\sqrt{y_2^2} + \sqrt{y_1^2}\sqrt{y_2^2}] + R &= 0 \Rightarrow y_0^3 + \frac{P}{2}y_0^2 + \frac{P^2-4R}{16}y_0 - \frac{Q^2}{64} = 0 \quad (14) \end{aligned}$$

To solve the resulted expression in (Eq. 14), we use Cardan's solution for third degree polynomial equations.

$$\text{For } w^3 + cw + d = 0: w = \sqrt[3]{-\frac{d}{2} + \sqrt{\left(\frac{d}{2}\right)^2 + \left(\frac{c}{3}\right)^3}} + \sqrt[3]{-\frac{d}{2} - \sqrt{\left(\frac{d}{2}\right)^2 + \left(\frac{c}{3}\right)^3}} \quad (15)$$

For $y^3 + by^2 + cy + d = 0$, we use the form $y = \frac{-b+w}{3}$, and we suppose $D = 27d + 2b^3 - 9cb$ and $C = 9c - 3b^2$ to express the cubic solution as follow:

$$y = \frac{-b}{3} + \frac{1}{3}\sqrt[3]{-\frac{D}{2} + \sqrt{\left(\frac{D}{2}\right)^2 + \left(\frac{C}{3}\right)^3}} + \frac{1}{3}\sqrt[3]{-\frac{D}{2} - \sqrt{\left(\frac{D}{2}\right)^2 + \left(\frac{C}{3}\right)^3}} \quad (16)$$

By using the expression (Eq. 16), the solutions of third degree polynomial equation shown in (Eq. 14) are $y_{0,1}$ in (Eq. 17), $y_{0,2}$ in (Eq. 18) and $y_{0,3}$ in (Eq. 19), where $P = \frac{P}{2}$, $R = \frac{-27Q^2-2P^3+72PR}{64}$ and $Q = -\frac{3P^2+36R}{16}$.

$$y_{0,1} = -\frac{P}{3} + \frac{1}{3} \sqrt{-\frac{R}{2} + \sqrt{\left(\frac{R}{2}\right)^2 + \left(\frac{Q}{3}\right)^3}} + \frac{1}{3} \sqrt{-\frac{R}{2} - \sqrt{\left(\frac{R}{2}\right)^2 + \left(\frac{Q}{3}\right)^3}} \quad (17)$$

In condition of $y_{0,1} \neq 0$, $y_{0,2}$ and $y_{0,3}$ are as follow:

$$y_{0,2} = -\frac{\frac{P}{2} + y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}} \quad (18)$$

$$y_{0,3} = -\frac{\frac{P}{2} + y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}} \quad (19)$$

We deduce that when y_0 takes the value $y_{0,1}$, the value of $y_{0,2}$ is equal to the shown value of y_1 in (Eq. 9), and the value of $y_{0,3}$ is equal to the shown value of y_2 in (Eq. 10).

There are three other possible expressions for y which

respect the proposition $-8\sqrt{y_0}\sqrt{y_1}\sqrt{y_2} = Q$ when $Q \leq 0$, and they give the same results of calculations toward having the shown third degree polynomial in (Eq. 14). Thereby, they give the same values for roots $y_{0,1}$, $y_{0,2}$ and $y_{0,3}$. These three expressions are $y = -\sqrt{y_0} - \sqrt{y_1} + \sqrt{y_2}$, $y = -\sqrt{y_0} + \sqrt{y_1} - \sqrt{y_2}$ and $y = \sqrt{y_0} - \sqrt{y_1} - \sqrt{y_2}$.

By using the shown expressions in (Eq. 6) and (Eq. 17), the solutions of presented fourth degree polynomial equation in (Eq. 5) when $\left(8\left(\frac{b}{a}\right)^3 - 32\frac{cb}{a^2} + 64\frac{d}{a}\right) < 0$ are as shown in (Eq. 20), (Eq. 21), (Eq. 22) and (Eq. 23).

$$\text{Solution 1: } s_{1,1} = \sqrt{y_{0,1}} + \sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} + \sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} \quad (20)$$

$$\text{Solution 2: } s_{1,2} = -\sqrt{y_{0,1}} - \sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} + \sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} \quad (21)$$

$$\text{Solution 3: } s_{1,3} = -\sqrt{y_{0,1}} + \sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} - \sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} \quad (22)$$

$$\text{Solution 4: } s_{1,4} = \sqrt{y_{0,1}} - \sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} - \sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} \quad (23)$$

There are three other possible expressions for y which respect the proposition $8\sqrt{y_0}\sqrt{y_1}\sqrt{y_2} = Q$ when $Q \geq 0$, and they give the same third degree polynomial shown in (Eq. 14) after calculations. These three expressions are $y = -\sqrt{y_0} + \sqrt{y_1} + \sqrt{y_2}$, $y = \sqrt{y_0} - \sqrt{y_1} + \sqrt{y_2}$ and $y = \sqrt{y_0} + \sqrt{y_1} - \sqrt{y_2}$.

By using the shown expressions in (Eq. 6) and (Eq. 17), the solutions of presented fourth degree polynomial equation in (Eq. 5) when $\left(8\left(\frac{b}{a}\right)^3 - 32\frac{cb}{a^2} + 64\frac{d}{a}\right) > 0$ are as shown in (Eq. 24), (Eq. 25), (Eq. 26) and (Eq. 27).

$$\text{Solution 1: } s_{2,1} = -\sqrt{y_{0,1}} - \sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} - \sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} \quad (24)$$

$$\text{Solution 2: } s_{2,2} = -\sqrt{y_{0,1}} + \sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} + \sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} \quad (25)$$

$$\text{Solution 3: } s_{2,3} = \sqrt{y_{0,1}} - \sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} + \sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} \quad (26)$$

$$\text{Solution 4: } s_{2,4} = \sqrt{y_{0,1}} + \sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} - \sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} \quad (27)$$

Concerning $\left(8\left(\frac{b}{a}\right)^3 - 32\frac{cb}{a^2} + 64\frac{d}{a}\right) = 0$:

The expression of $y_{0,1}$ is as shown in (Eq. 28) where $P = \frac{P}{2}$, $R = \frac{-2P^3 + 72PR}{64}$ and $Q = -\frac{3P^2 + 36R}{16}$, whereas $y_{0,2}$ and $y_{0,3}$ are as shown in (Eq. 29) and (Eq. 30).

$$y_{0,1} = \frac{-P}{3} + \frac{1}{3} \sqrt[3]{-\frac{R}{2} + \sqrt{\left(\frac{R}{2}\right)^2 + \left(\frac{Q}{3}\right)^3}} + \frac{1}{3} \sqrt[3]{-\frac{R}{2} - \sqrt{\left(\frac{R}{2}\right)^2 + \left(\frac{Q}{3}\right)^3}} \quad (28)$$

$$y_{0,2} = -\frac{\frac{P}{2} + y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}} = 0 \text{ or } y_{0,2} = -\left(\frac{P}{2} + y_{0,1}\right) \quad (29)$$

$$y_{0,3} = -\frac{\frac{P}{2} + y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}} = -\left(\frac{P}{2} + y_{0,1}\right) \text{ or } y_{0,3} = 0 \quad (30)$$

Because of having either of the expressions ($y_{0,2} = 0$) or ($y_{0,3} = 0$), and having the intersection between the forms (Eq. 7) and (Eq. 8) for $Q=0$ ($y = \sqrt{y_0} + \sqrt{y_1} + \sqrt{y_2}$ for $Q \leq 0$ and $y = -\sqrt{y_0} - \sqrt{y_1} - \sqrt{y_2}$ for $Q \geq 0$), there are four solutions for the polynomial equation shown in (Eq. 5) and they are as shown in (Eq. 31), (Eq. 32), (Eq. 33) and (Eq. 34).

$$\text{Solution 1: } s_{3,1} = \sqrt{y_{0,1}} + \sqrt{-\left(\frac{P}{2} + y_{0,1}\right)} \quad (31)$$

$$\text{Solution 2: } s_{3,2} = -\sqrt{y_{0,1}} - \sqrt{-\left(\frac{P}{2} + y_{0,1}\right)} \quad (32)$$

$$\text{Solution 3: } s_{3,3} = -\sqrt{y_{0,1}} + \sqrt{-\left(\frac{P}{2} + y_{0,1}\right)} \quad (33)$$

$$\text{Solution 4: } s_{3,4} = \sqrt{y_{0,1}} - \sqrt{-\left(\frac{P}{2} + y_{0,1}\right)} \quad (34)$$

When we give the value $y_{0,1}$ in (Eq. 17) to y_0 , the values of $y_{0,2}$ in (Eq. 18) and $y_{0,3}$ in (Eq. 19) are equal to the shown values of y_1 in (Eq. 9) and y_3 in (Eq. 10) respectively. Thereby, even when we replace the value of $y_{0,1}$ in the expressions of proposed solutions by the values of $y_{0,2}$ or $y_{0,3}$, the results are only redundancies of proposed solutions, because the value of P in the precedent expressions and in the proposed solutions is as follow:

$$\frac{P}{2} = -(\sqrt{y_0^2} + \sqrt{y_1^2} + \sqrt{y_2^2}) = -(\sqrt{y_{0,1}^2} + \sqrt{y_{0,2}^2} + \sqrt{y_{0,3}^2}).$$

In order to solve the polynomial equation shown in (Eq. 2), we use the expression $x = \frac{-b+y}{4}$ where y is the unknown variable in polynomial equation (Eq. 5). By using expressions (Eq. 6) and (Eq. 17) for $\left(8\left(\frac{b}{a}\right)^3 - 32\frac{cb}{a^2} + 64\frac{d}{a}\right) < 0$, the solutions for equation (Eq. 1) are as shown in (Eq. 35), (Eq. 36), (Eq. 37) and (Eq. 38).

$$\text{Solution 1: } S_{1,1} = -\frac{b}{4a} + \frac{1}{4}\sqrt{y_{0,1}} + \frac{1}{4}\sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} + \frac{1}{4}\sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} \quad (35)$$

$$\text{Solution 2: } S_{1,2} = -\frac{b}{4a} - \frac{1}{4}\sqrt{y_{0,1}} - \frac{1}{4}\sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} + \frac{1}{4}\sqrt{-\frac{\frac{P}{2} + y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2} + y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} \quad (36)$$

$$\text{Solution 3: } S_{1,3} = -\frac{b}{4a} - \frac{1}{4}\sqrt{y_{0,1}} + \frac{1}{4}\sqrt{-\frac{\frac{P}{2}+y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2}+y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} - \frac{1}{4}\sqrt{-\frac{\frac{P}{2}+y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2}+y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} \quad (37)$$

$$\text{Solution 4: } S_{1,4} = -\frac{b}{4a} + \frac{1}{4}\sqrt{y_{0,1}} - \frac{1}{4}\sqrt{-\frac{\frac{P}{2}+y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2}+y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} - \frac{1}{4}\sqrt{-\frac{\frac{P}{2}+y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2}+y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} \quad (38)$$

By using the expression $x = \frac{\frac{b}{a}+y}{4}$ while relying on shown expressions in (Eq. 6) and (Eq. 17) for $\left(8\left(\frac{b}{a}\right)^3 - \frac{32cb}{a^2} + \frac{64d}{a}\right) > 0$, the proposed solutions for equation (Eq. 1) are as shown in (Eq. 39), (Eq. 40), (Eq. 41) and (Eq. 42).

$$\text{Solution 1: } S_{2,1} = -\frac{b}{4a} - \frac{1}{4}\sqrt{y_{0,1}} - \frac{1}{4}\sqrt{-\frac{\frac{P}{2}+y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2}+y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} - \frac{1}{4}\sqrt{-\frac{\frac{P}{2}+y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2}+y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} \quad (39)$$

$$\text{Solution 2: } S_{2,2} = -\frac{b}{4a} - \frac{1}{4}\sqrt{y_{0,1}} + \frac{1}{4}\sqrt{-\frac{\frac{P}{2}+y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2}+y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} + \frac{1}{4}\sqrt{-\frac{\frac{P}{2}+y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2}+y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} \quad (40)$$

$$\text{Solution 3: } S_{2,3} = -\frac{b}{4a} + \frac{1}{4}\sqrt{y_{0,1}} - \frac{1}{4}\sqrt{-\frac{\frac{P}{2}+y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2}+y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} + \frac{1}{4}\sqrt{-\frac{\frac{P}{2}+y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2}+y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} \quad (41)$$

$$\text{Solution 4: } S_{2,4} = -\frac{b}{4a} + \frac{1}{4}\sqrt{y_{0,1}} + \frac{1}{4}\sqrt{-\frac{\frac{P}{2}+y_{0,1}}{2} + \sqrt{\left(\frac{\frac{P}{2}+y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} - \frac{1}{4}\sqrt{-\frac{\frac{P}{2}+y_{0,1}}{2} - \sqrt{\left(\frac{\frac{P}{2}+y_{0,1}}{2}\right)^2 - \frac{Q^2}{64y_{0,1}}}} \quad (42)$$

By using the expression $x = \frac{\frac{b}{a}+y}{4}$ while relying on expressions (Eq. 6) and (Eq. 28) for $\left(8\left(\frac{b}{a}\right)^3 - \frac{32cb}{a^2} + \frac{64d}{a}\right) = 0$, the proposed solutions for equation (Eq. 1) are as shown in (Eq. 43), (Eq. 44), (Eq. 45) and (Eq. 46).

$$\text{Solution 1: } S_{3,1} = -\frac{b}{4a} + \frac{1}{4}\sqrt{y_{0,1}} + \frac{1}{4}\sqrt{-\left(\frac{P}{2} + y_{0,1}\right)} \quad (43)$$

$$\text{Solution 2: } S_{3,2} = -\frac{b}{4a} - \frac{1}{4}\sqrt{y_{0,1}} - \frac{1}{4}\sqrt{-\left(\frac{P}{2} + y_{0,1}\right)} \quad (44)$$

$$\text{Solution 3: } S_{3,3} = -\frac{b}{4a} - \frac{1}{4}\sqrt{y_{0,1}} + \frac{1}{4}\sqrt{-\left(\frac{P}{2} + y_{0,1}\right)} \quad (45)$$

$$\text{Solution 4: } S_{3,4} = -\frac{b}{4a} + \frac{1}{4}\sqrt{y_{0,1}} - \frac{1}{4}\sqrt{-\left(\frac{P}{2} + y_{0,1}\right)} \quad (46)$$

3. New Five Formulary Solutions for Quintic Polynomial Equation in General Form

In this section, we propose five new formulary solutions for

fifth degree polynomial equation in general form shown in (Eq. 47), where we rely on our proposed solutions for fourth degree polynomial equations to develop the structure of proposed solutions for quintic equation. We rely on used logic in precedent theorem (Theorem 1) by projection on quintic equations to prove the expressions of developed roots for fifth degree polynomials.

3.1. Introduction on Second Proposed Theorem

In this subsection, we present our second proposed theorem to introduce new five formulary solutions for fifth degree polynomial equation in general form shown in (Eq. 47). First, we divide the polynomial (Eq. 47) on A and then we replace w with $\frac{-B+A^2x}{5}$ to reduce the polynomial expression to the simplified form shown in (Eq. 48) where coefficients are expressed as shown in (Eq. 49), (Eq. 50), (Eq. 51) and (Eq. 52).

$$Aw^5 + Bw^4 + Cw^3 + Dw^2 + Ew + F = 0 \text{ with } A \neq 0 \quad (47)$$

$$x^5 + cx^3 + dx^2 + ex + f = 0 \quad (48)$$

$$c = -10\frac{B^2}{A^2} + 25\frac{C}{A} \quad (49)$$

$$d = 20\frac{B^3}{A^3} - 75\frac{CB}{A^2} + 125\frac{D}{A} \quad (50)$$

$$e = -15\frac{B^4}{A^4} + 75\frac{CB^2}{A^3} - 250\frac{DB}{A^2} + 625\frac{E}{A} \quad (51)$$

$$f = 4\frac{B^5}{A^5} - 25\frac{CB^3}{A^4} + 125\frac{DB^2}{A^3} - 625\frac{EB}{A^2} + 3125\frac{F}{A} \quad (52)$$

$$z^4 + \Gamma_3 z^3 + \Gamma_2 z^2 + \Gamma_1 z + \Gamma_0 = 0 \quad (53)$$

3.2. Theorem 2

After reducing the form of fifth degree polynomial shown in (Eq. 47) to the presented form in (Eq. 48) where coefficients are expressed in (Eq. 49), (Eq. 50), (Eq. 51) and (Eq. 52); the fifth degree polynomial equation shown in (Eq. 48), where coefficients belong to the group of numbers \mathbb{R} , can be reduced to a fourth degree polynomial equation, which may be expressed as shown in (Eq. 53). The reduction from quantic polynomial to quartic polynomial is conducted by supposing $= x_0x_1 + x_0x_2 + x_0x_3 + x_1x_2 + x_1x_3 + x_2x_3$, whereas supposing $z = (x_0 + x_1 + x_2 + x_3)$ is the solution for fourth degree polynomial equation in (Eq. 53) by using Theorem 1 and relying on the expression $x_3 = -\frac{\Gamma_3}{4}$. The variable Γ_3 is the solution for the polynomial equation shown in (Eq. 54), whereas the coefficients of this equation are shown in (Eq. 55), (Eq. 56) and (Eq. 57). The coefficients Γ_2 , Γ_1 and Γ_0 of quartic equation (Eq. 53) are determined by using calculated values of Γ_3 and using the shown expressions in (Eq. 58), (Eq. 59) and (Eq. 60). As a result, we have eight calculated values as potential solutions for fifth degree polynomial equation shown in (Eq. 48), where many of them are only redundancies of others, because there are only five official solutions to determine.

The eight solutions to calculate for quantic equation (Eq. 48) are as shown in the groups (Eq. 89) and (Eq. 90). The proposed five values as official solutions for fifth degree poly-

nomial equation shown in (Eq. 48) are as presented in (Eq. 91), (Eq. 92), (Eq. 93), (Eq. 94) and (Eq. 95). The proposed five values as official solutions for fifth degree polynomial equation shown in (Eq. 47) are as presented in (Eq. 96), (Eq. 97), (Eq. 98), (Eq. 99) and (Eq. 100).

$$\lambda_2 \Gamma_3^4 + \lambda_1 \Gamma_3^2 + \lambda_0 = 0 \quad (54)$$

$$\lambda_2 = 1024 \frac{(e - \frac{c^2}{4})}{16d}; \quad (55)$$

$$\lambda_1 = 512c - 40 \frac{(16d)^2}{e - \frac{c^2}{4}}; \quad (56)$$

$$\lambda_0 = -128 \frac{(16d)^2}{(e - \frac{c^2}{4})^2} [f - \frac{cd}{2}]; \quad (57)$$

$$\Gamma_2 = \frac{(16d)^2}{2\Gamma_3^2(e - \frac{c^2}{4})} \quad (58)$$

$$\Gamma_1 = -\frac{1}{4}\Gamma_3^3 + \frac{3(16d)^2}{16(e - \frac{c^2}{4})\Gamma_3} \quad (59)$$

$$\Gamma_0 = -\frac{1}{16}\Gamma_3^4 - \frac{(16d)^2}{32(e - \frac{c^2}{4})} + \frac{(f - \frac{cd}{2})(16d)^2}{2\Gamma_3^2(e - \frac{c^2}{4})^2} + \frac{(16d)^4}{16\Gamma_3^4(e - \frac{c^2}{4})^2} \quad (60)$$

3.3. Proof of Second Proposed Theorem

Considering the quantic equation shown in (Eq. 48), we propose the expression (Eq. 61) in order to reduce the form of quantic equation to a fourth degree polynomial equation. We also propose the expression (Eq. 62), which presents the solution form of quartic equation by extending the used logic and presented solutions in Theorem 1.

$$x = x_0x_1 + x_0x_2 + x_0x_3 + x_1x_2 + x_1x_3 + x_2x_3 \quad (61)$$

$$z = x_0 + x_1 + x_2 + x_3 \quad (62)$$

We replace x with its proposed value in (Eq. 61), in order to end by calculations to the reduced form shown in (Eq. 53).

In the shown expressions in (Eq. 63), we rely on the use of x_i , x_j and x_k where $\{x_i, x_j, x_k\} \in \{x_0, x_1, x_2, x_3\}$ and $i \neq j \neq k$.

$$\alpha_1 = \sum_{i=0}^3 x_i^2; \alpha_2 = \sum_{i \neq j} x_i^2 x_j^2; \alpha_3 = \sum_{i \neq j \neq k} x_i x_j x_k; \alpha_4 = x_0 x_1 x_2 x_3 \quad (63)$$

$$x = \frac{[z^2 - \alpha_1]}{2} \quad (64)$$

$$x^2 = \alpha_2 + 2\alpha_3 z + 6\alpha_4 \quad (65)$$

$$x^3 = z^3 \alpha_3 + \frac{1}{2} z^2 [\alpha_2 + 6\alpha_4] - z \alpha_1 \alpha_3 - \frac{1}{2} [\alpha_2 + 6\alpha_4] \alpha_1 \quad (66)$$

$$x^5 = 2z^4 \alpha_3^2 + 2z^3 [\alpha_2 + 6\alpha_4] \alpha_3 + \frac{1}{2} z^2 [(\alpha_2 + 6\alpha_4)^2 - 4\alpha_3^2 \alpha_1] - 2z \alpha_3 \alpha_1 [\alpha_2 + 6\alpha_4] - \frac{1}{2} \alpha_1 [\alpha_2 + 6\alpha_4]^2 \quad (67)$$

$$\gamma_4 z^4 + \gamma_3 z^3 + \gamma_2 z^2 + \gamma_1 z + \gamma_0 = 0 \quad (68)$$

We use the expressions of $\{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$ in (Eq. 63), x in (Eq. 64), x^2 in (Eq. 65), x^3 in (Eq. 66) and x^5 in (Eq. 67), to have the fourth degree polynomial shown in (Eq. 68) where the values of coefficients are as follow:

$$\gamma_4 = 2\alpha_3^2$$

$$\gamma_3 = 2[\alpha_2 + 6\alpha_4] \alpha_3 + c\alpha_3$$

$$\gamma_2 = \frac{1}{2} [(\alpha_2 + 6\alpha_4)^2 - 4\alpha_3^2 \alpha_1] + \frac{1}{2} c[\alpha_2 + 6\alpha_4] + \frac{e}{2}$$

$$\gamma_1 = -2\alpha_3 \alpha_1 [\alpha_2 + 6\alpha_4] - c\alpha_1 \alpha_3 + 2d\alpha_3$$

$$\gamma_0 = -\frac{1}{2} \alpha_1 [\alpha_2 + 6\alpha_4]^2 - \frac{1}{2} c[\alpha_2 + 6\alpha_4] \alpha_1 + d[\alpha_2 + 6\alpha_4] - \frac{e\alpha_1}{2} + f$$

We divide the polynomial equation shown in (Eq. 68) on γ_4 to simplify its expression. As a result, we have the shown expression in (Eq. 53) where the values of coefficients are as follow:

$$\Gamma_3 = \frac{[\alpha_2 + 6\alpha_4] + \frac{c}{2}}{\alpha_3} \Rightarrow \alpha_2 = \alpha_3 \Gamma_3 - \frac{c}{2} - 6\alpha_4$$

$$\Gamma_2 = \frac{\Gamma_3^2}{4} - \alpha_1 + \frac{e - \frac{c^2}{4}}{4\alpha_3^2}$$

$$\Gamma_1 = -\Gamma_3 \alpha_1 + \frac{d}{\alpha_3}$$

$$\Gamma_0 = -\frac{1}{4} \alpha_1 \Gamma_3^2 + \frac{d[\alpha_3 \Gamma_3 - \frac{c}{2}]}{2\alpha_3^2} + \frac{f - \frac{e\alpha_1}{2} + \frac{c^2 \alpha_1}{8}}{2\alpha_3^2}$$

$$R = \Gamma_3^4 - 16\Gamma_3^2 \alpha_1 + 4\Gamma_3^2 \frac{e - \frac{c^2}{4}}{\alpha_3^2} + 64d \frac{\Gamma_3}{\alpha_3} + 256 \frac{f - \frac{e\alpha_1}{2} + \frac{cd}{2} + \frac{c^2 \alpha_1}{8}}{2\alpha_3^2} \quad (73)$$

By using the shown expressions in (Eq. 11), (Eq. 12), (Eq. 13) and (Eq. 14) from the proof of first proposed theorem (Theorem 1), the values of P , Q and R will be as shown in (eq74), (Eq. 75) and (Eq. 76) successively.

$$P = -2[(4x_0)^2 + (4x_1)^2 + (4x_2)^2] \Rightarrow P = -32[\alpha_1 - \frac{\Gamma_3^2}{16}] \quad (74)$$

$$Q = -8(4x_0)(4x_1)(4x_2) \Rightarrow Q = -\frac{8(4x_0)(4x_1)(4x_2)(4x_3)}{4x_3} = 2048 \frac{\alpha_4}{\Gamma_3} \quad (75)$$

$$R = [(4x_0)^2 + (4x_1)^2 + (4x_2)^2]^2 - 4[(4x_0)^2(4x_1)^2 + (4x_0)^2(4x_2)^2 + (4x_1)^2(4x_2)^2] \\ \Rightarrow R = 256 \left[\alpha_1 - \frac{\Gamma_3^2}{16} \right]^2 - 1024 \left[\alpha_2 - \frac{\Gamma_3^2}{16} \left(\alpha_1 - \frac{\Gamma_3^2}{16} \right) \right] \quad (76)$$

From precedent section (section 2), we ended with solutions expressed as $y = y_0 + y_1 + y_2$ for fourth degree polynomial equation in simple form shown in (Eq. 5), whereas the solution for fourth degree polynomial equation in complete form is expressed as $z = -\frac{b}{4a} + \frac{1}{4}y_0 + \frac{1}{4}y_1 + \frac{1}{4}y_2$.

We replace z with $\frac{-\Gamma_3 + y}{4}$, in order to reduce the form of quartic equation from expression (Eq. 53) to expression (Eq. 69), where the values of coefficients are as shown in (Eq. 70).

$$y^4 + Py^2 + Qy + R = 0 \quad (69)$$

$$P = -6\Gamma_3^2 + 16\Gamma_2; Q = 8\Gamma_3^3 - 32\Gamma_2 \Gamma_3 + 64\Gamma_1; R = -3\Gamma_3^4 + 16\Gamma_2 \Gamma_3^2 - 64\Gamma_1 \Gamma_3 + 256\Gamma_0 \quad (70)$$

Concerning the fourth degree polynomial equation in (Eq. 53), where z is as shown in (Eq. 62), the principal proposed expressions for the solutions are $z = -\frac{\Gamma_3}{4} + \frac{1}{4}\sqrt{y_0} + \frac{1}{4}\sqrt{y_1} + \frac{1}{4}\sqrt{y_2}$ when $Q \leq 0$ and $z = -\frac{\Gamma_3}{4} - \frac{1}{4}\sqrt{y_0} - \frac{1}{4}\sqrt{y_1} - \frac{1}{4}\sqrt{y_2}$ when $Q \geq 0$; where $x_3 = -\frac{\Gamma_3}{4}$, $x_0 = \pm \frac{1}{4}\sqrt{y_0}$, $x_1 = \pm \frac{1}{4}\sqrt{y_1}$ and $x_2 = \pm \frac{1}{4}\sqrt{y_2}$. These two principal expressions are sufficient to conduct the calculations of proof, and then generalize the results by using the other expressed forms of solutions in Theorem 1.

We replace Γ_2 , Γ_1 and Γ_0 with their values in function of $\{\Gamma_3, \alpha_1, \alpha_3\}$, in order to have the expressions of P in (Eq. 71), Q in (Eq. 72) and R in (Eq. 73).

$$P = -2\Gamma_3^2 - 16\alpha_1 + 4 \frac{(e - \frac{c^2}{4})}{\alpha_3^2} \quad (71)$$

$$Q = -32\Gamma_3 \alpha_1 - 8\Gamma_3 \frac{(e - \frac{c^2}{4})}{\alpha_3^2} + 64 \frac{d}{\alpha_3} \quad (72)$$

We have a group of four variables $\{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$, whereas we have a group of only three equations to solve $\{P, Q, R\}$ where all of them are dependent on the value of Γ_3 . Thereby, the next step is about using the appropriate logic of analysis and calculation to find the value of Γ_3 while taking advantage of the fact that having a group of four variables enables us to solve four equations.

In order to reduce the expression of R in (Eq. 73), and find a way to determine the value of Γ_3 , we suppose that $\left(4\frac{e-\frac{c^2}{4}}{\alpha_3^2}\Gamma_3^2 + 64\frac{d}{\alpha_3}\right) = 0$ where $\frac{\Gamma_3}{\alpha_3} \neq 0$. As a result, we have the shown expression in (Eq. 77).

$$\frac{\Gamma_3}{\alpha_3} = -16\frac{d}{e-\frac{c^2}{4}} \quad (77)$$

From precedent calculations of dividing the polynomial equation shown in (Eq. 68) on γ_4 to simplify its expression, along with using the expression of Γ_3 , we have $\alpha_2 = (\alpha_3\Gamma_3 - \frac{c}{2} - 6\alpha_4)$.

We use the expression (Eq. 77) and we replace α_2 with $(\alpha_3\Gamma_3 - \frac{c}{2} - 6\alpha_4)$, in order to pass from expression (Eq. 76) to expression (Eq. 78).

$$R = -3\Gamma_3^4 + 32\Gamma_3^2\alpha_1 - 1024\left[-\frac{\Gamma_3^2}{16d}\left(e - \frac{c^2}{4}\right) - \frac{c}{2} - 6\alpha_4\right] + 256\alpha_1^2 \quad (78)$$

We have the resulted equation in (Eq. 79) by using the expressions of P in (Eq. 71) and (Eq. 74).

$$-4\Gamma_3^2 + 16\alpha_1 + 4\frac{e-\frac{c^2}{4}}{\alpha_3^2} = 0 \quad (79)$$

$$\alpha_1 = \frac{\Gamma_3^4 - \frac{(16d)^2}{e-\frac{c^2}{4}}}{4\Gamma_3^2} \quad (80)$$

We have the presented polynomial equation in (Eq. 81) by using the expressions of R in (Eq. 73) and (Eq. 78).

$$-4\Gamma_3^4 + 48\Gamma_3^2\alpha_1 - 1024\left[-\frac{\Gamma_3^2}{16d}\left(e - \frac{c^2}{4}\right) - \frac{c}{2} - 6\alpha_4\right] - 256\frac{f - \frac{e\alpha_1}{2} - \frac{cd}{2} + \frac{c^2\alpha_1}{8}}{2\alpha_3^2} + 256\alpha_1^2 = 0 \quad (81)$$

We replace α_1 with its value in (Eq. 80) and we use the deduced value of α_4 from (Eq. 75) and (Eq. 82), in order to pass from the expression of polynomial equation shown in (Eq. 81) to the polynomial equation shown in (Eq. 83) where the coefficients are structured only by using c , d , e and f .

$$\alpha_4 = x_1x_2x_3x_4 \Rightarrow 2048\alpha_4 = -32\Gamma_3^2\alpha_1 - 8\Gamma_3^2\frac{e-\frac{c^2}{4}}{\alpha_3^2} + 64\Gamma_3\frac{d}{\alpha_3} \quad (82)$$

$$\lambda_2(\Gamma_3^2)^3 + \lambda_1(\Gamma_3^2)^2 + \lambda_0(\Gamma_3^2) = 0 \quad (83)$$

The values of coefficients for polynomial equation shown in (Eq. 83) are as shown in (Eq. 55), (Eq. 56) and (Eq. 57).

Since we supposed that $(4\Gamma_3^2\frac{e-\frac{c^2}{4}}{\alpha_3^2} + 64d\frac{\Gamma_3}{\alpha_3}) = 0$, where we adopted the shown value in (Eq. 77) for $\frac{\Gamma_3}{\alpha_3}$, we eliminate the root zero as solution for polynomial equation (Eq. 83) and we use the quadratic solution to solve the polynomial equation shown in (Eq. 54), because all coefficients are expressed only in function of c, d, e and f . As a result, we have four possible values for Γ_3 as solutions for polynomial equation shown in (Eq. 54).

We suppose that $G_{\{\Gamma_3\}}$ is the group of solutions for shown equation in (Eq. 54), where these solutions are expressed as $\Gamma_{3,i}$ and $-\Gamma_{3,i}$ with $1 \leq i \leq 2$. The group of solutions $G_{\{\Gamma_3\}}$ is determined by relying on quadratic roots.

$$G_{\{\Gamma_3\}} = \{\Gamma_{3,1}, \Gamma_{3,2}, -\Gamma_{3,1}, -\Gamma_{3,2}\} \quad (84)$$

Supposing that $P_i = \frac{\lambda_1}{\lambda_2}$ and $Q_i = \frac{\lambda_0}{\lambda_2}$, whereas using the expressions (Eq. 55), (Eq. 56) and (Eq. 57); the solutions $\Gamma_{3,1}$ and $\Gamma_{3,2}$ for shown equation in (Eq. 54) are as follow:

$$\Gamma_{3,1}^2 = -\frac{P_i}{2} + \sqrt{\left(\frac{P_i}{2}\right)^2 - 4Q_i}$$

$$\Gamma_{3,2}^2 = -\frac{P_i}{2} - \sqrt{\left(\frac{P_i}{2}\right)^2 - 4Q_i}$$

After determining the values of Γ_3 by using quadratic solutions, the following step consists of solving the polynomial equation shown in (Eq. 69).

The coefficients P in (Eq. 71) and R in (Eq. 73) are dependent on Γ_3^2 and α_1 , whereas α_1 in (Eq. 80) is dependent on Γ_3^2 and the coefficient Q shown in (Eq. 72) is dependent on Γ_3 and α_1 . Therefore, we are going to use only $\Gamma_{3,1}$ and $\Gamma_{3,2}$ to calculate the potential values of z , because $\{-\Gamma_{3,1}, -\Gamma_{3,2}\}$ are going only to inverse the sign of coefficient Q and thereby inverting the signs of potential values of z as solutions for polynomial equation shown in (Eq. 53), which will not influence the potential values of x as solutions for fifth degree polynomial equation shown in (Eq. 48) because $x = \frac{1}{2}(z^2 - \alpha_1)$.

We use the first proposed theorem (Theorem 1) to calculate the four solutions for fourth degree polynomial equation

shown in (Eq. 69) after calculating the coefficients P , Q and R for each value of Γ_3 from the group $\{\Gamma_{3,1}, \Gamma_{3,2}\}$. Thereby, we have eight values to calculate as potential solutions for the polynomial equation shown in (Eq. 69).

After using first proposed theorem to solve the equation (Eq. 69) for each value of Γ_3 from the group $\{\Gamma_{3,1}, \Gamma_{3,2}\}$, we have two groups of potential solutions for polynomial equation shown in (Eq. 69), where each group is dependent on different value of Γ_3 . We express these groups of solutions as follow: $K_{\{\Gamma_{3,1}\}}, K_{\{\Gamma_{3,2}\}}$.

$$K_{\{\Gamma_{3,1}\}} = \{S_{(\Gamma_{3,1,1})}, S_{(\Gamma_{3,1,2})}, S_{(\Gamma_{3,1,3})}, S_{(\Gamma_{3,1,4})}\} \quad (85)$$

$$K_{\{\Gamma_{3,2}\}} = \{S_{(\Gamma_{3,2,1})}, S_{(\Gamma_{3,2,2})}, S_{(\Gamma_{3,2,3})}, S_{(\Gamma_{3,2,4})}\} \quad (86)$$

Concerning the fourth degree polynomial equation shown in (Eq. 53), we have two groups of solutions where each group is dependent on different value of Γ_3 ; as shown in (Eq. 87) and (Eq. 88). The values of $S_{(\Gamma_{3,i,j})}$, where $1 \leq i \leq 2$ and $1 \leq j \leq 4$, are from the expressed solutions in the groups (Eq. 85) and (Eq. 86).

$$M_{\{\Gamma_{3,1}\}} = \left\{ -\frac{1}{4}\Gamma_{3,1} + \frac{1}{4}S_{(\Gamma_{3,1,1})}, -\frac{1}{4}\Gamma_{3,1} + \frac{1}{4}S_{(\Gamma_{3,1,2})}, -\frac{1}{4}\Gamma_{3,1} + \frac{1}{4}S_{(\Gamma_{3,1,3})}, -\frac{1}{4}\Gamma_{3,1} + \frac{1}{4}S_{(\Gamma_{3,1,4})} \right\} \quad (87)$$

$$M_{\{\Gamma_{3,2}\}} = \left\{ -\frac{1}{4}\Gamma_{3,2} + \frac{1}{4}S_{(\Gamma_{3,2,1})}, -\frac{1}{4}\Gamma_{3,2} + \frac{1}{4}S_{(\Gamma_{3,2,2})}, -\frac{1}{4}\Gamma_{3,2} + \frac{1}{4}S_{(\Gamma_{3,2,3})}, -\frac{1}{4}\Gamma_{3,2} + \frac{1}{4}S_{(\Gamma_{3,2,4})} \right\} \quad (88)$$

We suppose that $S_{(\Gamma_{3,i,j})} = (-\frac{1}{4}\Gamma_{3,i} + \frac{1}{4}S_{(\Gamma_{3,i,j})})$ where $1 \leq i \leq 2$ and $1 \leq j \leq 4$, in order to simplify the expressed values in (Eq. 87) and (Eq. 88). Thereby, we have two groups of values as potential solutions for fifth degree polynomial equation shown in (Eq. 48). These two groups are as shown in (Eq. 89) and (Eq. 90) where $\alpha_{(1,\Gamma_{3,i})} = \frac{\Gamma_{3,i}^4 - (16d)^2/(e-c^2/4)}{4\Gamma_{3,i}^2}$. The expression of $\alpha_{(1,\Gamma_{3,i})}$ is an extending of the shown expression of α_1 in (Eq. 80) by changing the value of Γ_3 , where $\Gamma_{3,i}$ belong to the group $\{\Gamma_{3,1}, \Gamma_{3,2}\}$.

$$N_{\{\Gamma_{3,1}\}} = \left\{ \frac{1}{2}[S_{(\Gamma_{3,1,1})}^2 - \alpha_{(1,\Gamma_{3,1})}], \frac{1}{2}[S_{(\Gamma_{3,1,2})}^2 - \alpha_{(1,\Gamma_{3,1})}], \frac{1}{2}[S_{(\Gamma_{3,1,3})}^2 - \alpha_{(1,\Gamma_{3,1})}], \frac{1}{2}[S_{(\Gamma_{3,1,4})}^2 - \alpha_{(1,\Gamma_{3,1})}] \right\} \quad (89)$$

$$N_{\{\Gamma_{3,2}\}} = \left\{ \frac{1}{2}[S_{(\Gamma_{3,2,1})}^2 - \alpha_{(1,\Gamma_{3,2})}], \frac{1}{2}[S_{(\Gamma_{3,2,2})}^2 - \alpha_{(1,\Gamma_{3,2})}], \frac{1}{2}[S_{(\Gamma_{3,2,3})}^2 - \alpha_{(1,\Gamma_{3,2})}], \frac{1}{2}[S_{(\Gamma_{3,2,4})}^2 - \alpha_{(1,\Gamma_{3,2})}] \right\} \quad (90)$$

We have eight values as potential solutions for the fifth degree polynomial equation shown in (Eq. 48). However, many of them are only redundancies of others and there are only five official solutions to determine. The variables $\{\Gamma_{3,1}, \Gamma_{3,2}\}$ are the responsible of solution redundancies from one group to other.

In order to avoid the complications of calculating the eight values from the groups $N_{\{\Gamma_{3,1}\}}$ and $N_{\{\Gamma_{3,2}\}}$, and then determining the five solutions for quantic equation shown in (Eq. 48), we propose the five expressed values in (Eq. 91), (Eq. 92), (Eq. 93), (Eq. 94) and (Eq. 95) as the five official solutions for fifth degree polynomial equation shown in (Eq. 48).

The first four proposed values as solutions for fifth degree polynomial equation shown in (Eq. 48) are from the group $N_{\{\Gamma_{3,1}\}}$, whereas the fifth value is expressed by deduction. The useless redundancies of solutions are from one group to other; therefore, we choose the first four solutions from the same group $N_{\{\Gamma_{3,1}\}}$.

In our five proposed solutions, we use the values $S_{(\Gamma_{3,1,j})} = -\frac{1}{4}\Gamma_{3,1} + \frac{1}{4}S_{(\Gamma_{3,1,j})}$ where $1 \leq j \leq 4$ and $S_{(\Gamma_{3,1,j})}$ from $K_{\{\Gamma_{3,1}\}}$. The variable $\alpha_{(1,\Gamma_{3,1})}$ is expressed as $\alpha_{(1,\Gamma_{3,1})} = \left[\Gamma_{3,1}^4 - \frac{(16d)^2}{e-c^2/4} \right] / [4\Gamma_{3,1}^2]$, whereas $\Gamma_{3,1}$ is from the group $G_{\{\Gamma_3\}}$ shown in (Eq. 84), which contains the solutions

for polynomial equation (Eq. 54). The values of $S_{(\Gamma_{3,1,j})}$, where $1 \leq j \leq 4$, are the solutions for quartic equation (Eq. 53) and they are determined by using Theorem 1.

$$s_1 = \frac{1}{2}[S_{(\Gamma_{3,1,1})}^2 - \alpha_{(1,\Gamma_{3,1})}] \quad (91)$$

$$s_2 = \frac{1}{2}[S_{(\Gamma_{3,1,2})}^2 - \alpha_{(1,\Gamma_{3,1})}] \quad (92)$$

$$s_3 = \frac{1}{2}[S_{(\Gamma_{3,1,3})}^2 - \alpha_{(1,\Gamma_{3,1})}] \quad (93)$$

$$s_4 = \frac{1}{2}[S_{(\Gamma_{3,1,4})}^2 - \alpha_{(1,\Gamma_{3,1})}] \quad (94)$$

$$s_5 = \frac{f}{s_1 s_2 s_3 s_4} \quad (95)$$

The proposed five solutions for fifth degree polynomial equation in complete form shown in (Eq. 47) are as expressed in (Eq. 96), (Eq. 97), (Eq. 98), (Eq. 99) and (Eq. 100).

$$\text{Solution 1: } S_1 = -\frac{B}{5A} + \frac{1}{10}[S_{(\Gamma_{3,1,1})}^2 - \alpha_{(1,\Gamma_{3,1})}] \quad (96)$$

$$\text{Solution 2: } S_2 = -\frac{B}{5A} + \frac{1}{10}[S_{(\Gamma_{3,1,2})}^2 - \alpha_{(1,\Gamma_{3,1})}] \quad (97)$$

$$\text{Solution 3: } S_3 = -\frac{B}{5A} + \frac{1}{10} \left[S_{(\Gamma_{3,1,3})}^2 - \alpha_{(1,\Gamma_{3,1})} \right] \quad (98)$$

$$\text{Solution 4: } S_4 = -\frac{B}{5A} + \frac{1}{10} \left[S_{(\Gamma_{3,1,4})}^2 - \alpha_{(1,\Gamma_{3,1})} \right] \quad (99)$$

$$\text{Solution 5: } S_5 = \frac{F}{AS_1S_2S_3S_4} \quad (100)$$

3.4. Introduction on Third Proposed Theorem

In this subsection, we present a third theorem to propose new five formulary solutions for fifth degree polynomial equation in general form shown in (Eq. 101), where coefficients belong to the group of numbers \mathbb{R} , without simplifying the polynomial by eliminating the fourth degree part. This third proposed theorem is based on the same logic and calculations of Theorem 2. However, it is distinguished by not using the expression $x = \frac{-b+y}{5}$ which we used in Theorem 2 to eliminate the fourth degree part of quantic polynomial.

The shown equation in (Eq. 101) is resulted by dividing the polynomial $Ax^5 + Bx^4 + Cx^3 + Dx^2 + Ex + F$ on A whereas $A \neq 0$.

$$x^5 + bx^4 + cx^3 + dx^2 + ex + f = 0 \quad (101)$$

$$b = \frac{B}{A}; c = \frac{C}{A}; d = \frac{D}{A}; e = \frac{E}{A}; f = \frac{F}{A}.$$

3.5. Theorem 3

$$Y_0 = -\frac{1}{16}Y_3^4 + bY_3^2 - \frac{[16(d-bc)]^2}{32\left(e-\frac{c^2}{4}\right)} + \frac{\left(f-\frac{cd}{2}+\frac{bc^2}{4}\right)[16(d-bc)]^2}{2Y_3^2\left(e-\frac{c^2}{4}\right)^2} + \frac{b[16(d-bc)]^2}{2Y_3^2\left(e-\frac{c^2}{4}\right)} + \frac{[16(d-bc)]^4}{16Y_3^4\left(e-\frac{c^2}{4}\right)^2} \quad (105)$$

$$\beta_2 Y_3^4 + \beta_1 Y_3^2 + \beta_0 = 0 \quad (106)$$

$$\beta_2 = 1024 \frac{\left(e-\frac{c^2}{4}\right)}{16(d-bc)} \quad (107)$$

$$\beta_1 = 512c - 40 \frac{[16(d-bc)]^2}{e-\frac{c^2}{4}} + 1024b^2 \quad (108)$$

$$\beta_0 = -128 \frac{[16(d-bc)]^2}{\left(e-\frac{c^2}{4}\right)^2} \left[f - \frac{cd}{2} + \frac{bc^2}{4} \right] + 128b \frac{[16(d-bc)]^2}{\left(e-\frac{c^2}{4}\right)} \quad (109)$$

3.6. Proof of Theorem 3

Considering the fifth degree polynomial equation shown in (Eq. 101), we use the expression (Eq. 61) as solution without eliminating the fourth degree part by the supposition $x = \frac{-b+y}{5}$. By using the expression (Eq. 61), we reduce the fifth degree polynomial (Eq. 101) to the quartic polynomial shown in (Eq. 102).

The fifth degree polynomial equation shown in (Eq. 101) is reducible to the quartic equation shown in (Eq. 102), where coefficients belong to the group of numbers \mathbb{R} without the need to eliminate the fourth degree part. The reduction from fifth degree to fourth degree is conducted by supposing $x = x_0x_1 + x_0x_2 + x_0x_3 + x_1x_2 + x_1x_3 + x_2x_3$, whereas supposing $z = (x_0 + x_1 + x_2 + x_3)$ is the solution for fourth degree polynomial equation shown in (Eq. 102) by using Theorem 1. The variable Y_3 is the solution for the polynomial equation shown in (Eq. 106) by using the expression of quadratic solution, whereas $x_3 = -\frac{Y_3}{4}$. The coefficients of shown polynomial in (Eq. 106) are as expressed in (Eq. 107), (Eq. 108) and (Eq. 109). The value of Y_3 is equal to $Y_{3,1}$, which is presented in (Eq. 120). The coefficients Y_2 , Y_1 and Y_0 are determined by using calculated value of Y_3 in (Eq. 120) and using the shown expressions in (Eq. 103), (Eq. 104) and (Eq. 105). The five proposed solutions for polynomial equation (Eq. 101) are as shown in (Eq. 121), (Eq. 122), (Eq. 123), (Eq. 124) and (Eq. 125).

$$z^4 + Y_3z^3 + Y_2z^2 + Y_1z + Y_0 = 0 \quad (102)$$

$$Y_2 = \frac{[16(d-bc)]^2}{2Y_3^2\left(e-\frac{c^2}{4}\right)} + 4b \quad (103)$$

$$Y_1 = -\frac{1}{4}Y_3^3 + \frac{3[16(d-bc)]^2}{16\left(e-\frac{c^2}{4}\right)Y_3} + 4bY_3 \quad (104)$$

$$x^4 = 4z^2 \alpha_3^2 + 4\alpha_3z[\alpha_2 + 6\alpha_4] + [\alpha_2 + 6\alpha_4]^2 \quad (110)$$

$$v_4z^4 + v_3z^3 + v_2z^2 + v_1z + v_0 = 0 \quad (111)$$

We rely on the expressions of $\{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$ in (Eq. 63), x in (Eq. 64), x^2 in (Eq. 65), x^3 in (Eq. 66), x^4 in (Eq. 110) and x^5 in (Eq. 67), to express the fourth degree polynomial shown in (Eq. 111) where the values of coefficients are as follow:

$$v_4 = 2\alpha_3^2$$

$$v_3 = 2[\alpha_2 + 6\alpha_4]\alpha_3 + c\alpha_3$$

$$v_2 = \frac{1}{2}[(\alpha_2 + 6\alpha_4)^2 - 4\alpha_3^2\alpha_1] + 4b\alpha_3^2 + \frac{1}{2}c[\alpha_2 + 6\alpha_4] + \frac{e}{2}$$

$$v_1 = -2\alpha_3\alpha_1[\alpha_2 + 6\alpha_4] + 4b\alpha_3[\alpha_2 + 6\alpha_4] - c\alpha_1\alpha_3 + 2d\alpha_3$$

$$v_0 = -\frac{1}{2}\alpha_1[\alpha_2 + 6\alpha_4]^2 + b[\alpha_2 + 6\alpha_4]^2 - \frac{1}{2}c[\alpha_2 + 6\alpha_4]\alpha_1 + d[\alpha_2 + 6\alpha_4] - \frac{e\alpha_1}{2} + f$$

We have the shown fourth degree polynomial in (Eq. 102) after dividing the polynomial (Eq. 111) on v_4 . The coefficients of shown polynomial in (Eq. 102) are as follow:

$$Y_3 = \frac{[\alpha_2 + 6\alpha_4] + \frac{c}{2}}{\alpha_3}$$

$$Y_2 = \frac{Y_3^2}{4} - \alpha_1 + 2b + \frac{e - \frac{c^2}{4}}{4\alpha_3^2}$$

$$Y_1 = -Y_3\alpha_1 + \frac{2b[\alpha_3 Y_3 - \frac{c}{2}]}{\alpha_3} + \frac{d}{\alpha_3}$$

$$Y_0 = -\frac{1}{4}\alpha_1 Y_3^2 + \frac{b[\alpha_3 Y_3 - \frac{c}{2}]^2}{2\alpha_3^2} + \frac{d[\alpha_3 Y_3 - \frac{c}{2}]}{2\alpha_3^2} + \frac{f - \frac{e\alpha_1}{2} + \frac{c^2\alpha_1}{8}}{2\alpha_3^2}$$

We have the fourth degree polynomial $y^4 + My^2 + Ny + O = 0$ by replacing z with $\frac{-Y_3 + y}{4}$ in the polynomial (Eq. 102). The coefficients M , N and O are as expressed in (Eq. 112), (Eq. 113) and (Eq. 114).

112), (Eq. 113) and (Eq. 114).

$$M = -2Y_3^2 - 16\alpha_1 + 32b + 4\frac{(e - \frac{c^2}{4})}{\alpha_3^2} \quad (112)$$

$$N = -32Y_3\alpha_1 + 64bY_3 - 8Y_3\frac{e - \frac{c^2}{4}}{\alpha_3^2} + 64\frac{d - bc}{\alpha_3} \quad (113)$$

$$O = Y_3^4 - 16Y_3^2\alpha_1 + 32bY_3^2 + 4Y_3^2\frac{e - \frac{c^2}{4}}{\alpha_3^2} + 64(d - bc)\frac{Y_3}{\alpha_3} + 256\frac{f - \frac{e\alpha_1}{2} - \frac{cd}{2} + \frac{bc^2}{4} + \frac{c^2\alpha_1}{8}}{2\alpha_3^2} \quad (114)$$

In order to reduce the expression of O in (Eq. 114), and find a way to determine the value of Y_3 , we suppose that $\left(4\frac{e - \frac{c^2}{4}}{\alpha_3^2}Y_3^2 + 64(d - bc)\frac{Y_3}{\alpha_3}\right) = 0$ whereas $\frac{Y_3}{\alpha_3} \neq 0$. As a result, we have the shown expression in (Eq. 115).

$$\frac{Y_3}{\alpha_3} = -16\frac{d - bc}{e - \frac{c^2}{4}} \quad (115)$$

We rely on the same used logic and processes of calculation in the proof of Theorem 2, in order to continue the proof of third proposed theorem. Thereby, the variables α_1 , α_2 and α_4 are as expressed in (Eq. 116), (Eq. 117) and (Eq. 118) respectively, whereas the resulted polynomial equation to determine the value of Y_3 is as shown in (Eq. 119).

$$\alpha_1 = \frac{Y_3^4 - 8bY_3^2\frac{[16(d - bc)]^2}{e - \frac{c^2}{4}}}{4Y_3^2} \quad (116)$$

$$\alpha_2 = \alpha_3 Y_3 - \frac{c}{2} - 6\alpha_4 \quad (117)$$

$$2048\alpha_4 = -32Y_3^2\alpha_1 + 64bY_3^2 - 8Y_3^2\frac{e - \frac{c^2}{4}}{\alpha_3^2} + 64Y_3\frac{(d - bc)}{\alpha_3} \quad (118)$$

$$-4Y_3^4 + [48\alpha_1 - 32b]Y_3^2 - 1024\alpha_2 - 256\frac{f - \frac{e\alpha_1}{2} - \frac{cd}{2} + \frac{bc^2}{4} + \frac{c^2\alpha_1}{8}}{2\alpha_3^2} + 256\alpha_1^2 = 0 \quad (119)$$

We use the shown value of $\frac{Y_3}{\alpha_3}$ in (Eq. 115) and we replace α_1 , α_2 and α_4 with their shown expressions in (Eq. 116), (Eq. 117) and (Eq. 118), in order to pass from equation (Eq. 119) to polynomial expression $\beta_2(Y_3^2)^3 + \beta_1(Y_3^2)^2 + \beta_0(Y_3^2) = 0$ where coefficients are as presented in (Eq. 107), (Eq. 108) and (Eq. 109).

Relying on the proof of Theorem 2, we calculate only the expression $Y_{3,1}$ shown in (Eq. 120) as a root for expressed equation in (Eq. 106), and then we determine the roots of quartic equation shown in (eq102).

To determine the roots of shown equation in (Eq. 102), we start by calculating the values of α_1 , α_4 and α_2 by replacing

the variable Y_3 with the value of $Y_{3,1}$, then we calculate the values of M , N and O , and we finish by using Theorem 1.

Supposing that $M^i = \frac{\beta_1}{\beta_2}$ and $N^i = \frac{\beta_0}{\beta_2}$, whereas using the expressions (Eq. 107), (Eq. 108) and (Eq. 109); the solution $Y_{3,1}$ for expressed polynomial equation in (Eq. 106) is as follow:

$$Y_{3,1} = \sqrt{-\frac{M^i}{2} + \sqrt{\left(\frac{M^i}{2}\right)^2 - 4N^i}} \quad (120)$$

As we mentioned in the proof of Theorem 2, the use of other roots of polynomial (Eq. 106) in the quartic polynomial

$y^4 + My^2 + Ny + O = 0$ to calculate the values of M , N and O will generate redundancies of roots for the quantic equation shown in (Eq. 101).

We determine the group $K_{Y_{3,1}}^i$, which contains the four roots for quartic equation $y^4 + My^2 + Ny + O = 0$, by using Theorem 1.

$$K_{Y_{3,1}}^i = \{S_{(Y_{3,1},1)}, S_{(Y_{3,1},2)}, S_{(Y_{3,1},3)}, S_{(Y_{3,1},4)}\}$$

We present the group of roots for quartic equation shown in (Eq. 102) as $M_{\{Y_{3,1}\}}^i$, which is determined by relying on the group $K_{Y_{3,1}}^i$.

$$M_{\{Y_{3,1}\}}^i = \left\{ -\frac{1}{4}Y_{3,1} + \frac{1}{4}S_{(Y_{3,1},1)}, -\frac{1}{4}Y_{3,1} + \frac{1}{4}S_{(Y_{3,1},2)}, -\frac{1}{4}Y_{3,1} + \frac{1}{4}S_{(Y_{3,1},3)}, -\frac{1}{4}Y_{3,1} + \frac{1}{4}S_{(Y_{3,1},4)} \right\}$$

We present each root for quartic equation shown in (Eq. 102) as $\xi_{(Y_{3,1},i)} = -\frac{1}{4}Y_{3,1} + \frac{1}{4}S_{(Y_{3,1},i)}$, whereas $S_{(Y_{3,1},i)}$ is from the group $K_{Y_{3,1}}^i$.

The proposed five solutions for quantic equation shown in (Eq. 101) are as expressed in (Eq. 121), (Eq. 122), (Eq. 123), (Eq. 124) and (Eq. 125). The expressions $\xi_{(Y_{3,1},1)}$, $\xi_{(Y_{3,1},2)}$, $\xi_{(Y_{3,1},3)}$ and $\xi_{(Y_{3,1},4)}$ present the calculated roots for quartic equation (Eq. 102) by using Theorem 1. $Y_{3,1}$ is calculated by using the shown expression in (Eq. 120). We use the expression (Eq. 116) to calculate the value of $\alpha_{(1,Y_{3,1})}$; which has the following value:

$$\alpha_{(1,Y_{3,1})} = \frac{Y_{3,1}^4 - 8bY_{3,1}^2 - \frac{[16(d-bc)]^2}{e - \frac{c^2}{4}}}{4Y_{3,1}^2}$$

$$\text{Solution 1: } S_1 = \frac{1}{2} \left[\xi_{(Y_{3,1},1)}^2 - \alpha_{(1,Y_{3,1})} \right] \quad (121)$$

$$\text{Solution 2: } S_2 = \frac{1}{2} \left[\xi_{(Y_{3,1},2)}^2 - \alpha_{(1,Y_{3,1})} \right] \quad (122)$$

$$\text{Solution 3: } S_3 = \frac{1}{2} \left[\xi_{(Y_{3,1},3)}^2 - \alpha_{(1,Y_{3,1})} \right] \quad (123)$$

$$\text{Solution 4: } S_4 = \frac{1}{2} \left[\xi_{(Y_{3,1},4)}^2 - \alpha_{(1,Y_{3,1})} \right] \quad (124)$$

$$\text{Solution 5: } S_5 = \frac{f}{S_1 S_2 S_3 S_4} \quad (125)$$

4. Conclusion

In the first presented theorem, we propose new four formulary solutions for any quartic polynomial equation in general form, which enable us to develop the formulary structures of five roots for any fifth degree polynomial equation in general form.

The proposed expressions as solutions in first theorem

enable to calculate the four roots of any quartic polynomial equation nearly simultaneously, whereas the proposed solutions in second theorem enable to calculate the five roots of any fifth degree polynomial equation nearly simultaneously. The third proposed theorem is based on the same logic and calculations of Theorem 2 whereas proposing five roots for quantic equation; however, it is distinguished by not eliminating the fourth-degree part of fifth degree polynomial by avoiding the use of the expression $x = \frac{-b+y}{5}$.

The second and third proposed theorems are based on reducing any fifth degree polynomial to a fourth degree polynomial, and then using Theorem 1 to solve the resulted quartic equation.

The principal criteria of presented theorems is proposing new radical solutions for quartic equations and quantic equations to enable the calculation of all roots of these equations nearly simultaneously, whereas using the expressions of quadratic roots and cubic roots as subparts of each proposed solution.

Author Contributions

Yassine Larbaoui is the sole author. The author read and approved the final manuscript.

Conflicts of Interest

The author declares no conflicts of interest.

References

- [1] Cardano, G. 1545. *Artis Magnae, Sive de Regulis Algebraicis Liber Unus*. English transl.: *The Great Art, or The Rules of Algebra*. Translated and edited by Witmer, T. R. MIT Press, Cambridge, Mass. (1968).
- [2] Dumit, D. S. Foote, R. M. 2004. *Abstract algebra*. John Wiley, 606-616.
- [3] Osler, T. J. 2001. Cardan polynomials and the reduction of radicals, *Mathematics Magazine*, 47(1): 26-32.
- [4] Janson, S. 2010. Roots of polynomials of degrees 3 and 4.
- [5] Lagrange, J. L. 1869. *Réflexions sur la résolution algébrique des équations*, in: *oeuvres de Lagrange*. Vol. 3, 205-421. J. A. Serret ed & Gauthier-Villars.
- [6] Faucette, W. M. 1996. A geometric interpretation of the solution of the general quartic polynomial. *Amer. Math. Monthly* 103: 51-57.
- [7] René D. 2012. *The Geometry of Rene Descartes with a facsimile of the 1st edition*, Courier Corporation.
- [8] Euler, L. 1738. *De formis radicum aequationum cuiusque ordinis coniectatio*. English transl.: *A conjecture on the forms of the roots of equations*. Translated by Bell, J. Cornell University, NY, USA, 2008.

- [9] Bewersdor, J. 2004. Algebra fur Einsteiger, Friedr. Vieweg Sohn Verlag. English transl.: Galois Theory for Beginners. A Historical Perspective. Translated by Kramer, D. American Mathematical Society (AMS), Providence, R. I., USA, 2006.
- [10] Helfgott, H. Helfgott, M. A modern vision of the work of cardano and ferrari on quartics. Convergence (MAA).
- [11] Rosen, M. 1995. Niels hendrik abel and equations of the fifth degree. American Mathematical Monthly 102(6): 495-505.
- [12] Garling, D. J. H. 1968. Galois Theory. Cambridge Univ. Press, Mass., USA.
- [13] Grillet, P. 2007. Abstract Algebra. 2nd ed. Springer, New York, USA.
- [14] van der Waerden, B. L. 1996. Algebra, Vol. 1, 3rd ed., Springer-Verlag, Berlin. English transl.: Algebra. Translated by Schulenberg J. R. and Blum, F. Springer-Verlag, New York, 1991.
- [15] Shmakov, S. L. 2011. A universal method of solving quartic equations, International Journal of Pure and Applied Mathematics 71(2): 251-259.
- [16] Fathi, A. and Sharifan, N. 2013. A classic new method to solve quartic equations, Applied and Computational Mathematics 2(2): 24-27. <https://doi.org/10.11648/j.acm.20130202.11>
- [17] Tehrani, F. T. 2020. Solution to polynomial equations, a new approach. Applied Mathematics 11(2): 53-66. <https://doi.org/10.4236/am.2020.112006>
- [18] Nahon, Y. J. 2018. Method for solving polynomial equations. Journal of Applied & Computational Mathematics 7(3): 2-12. <https://doi.org/10.4172/2168-9679.1000409>
- [19] Tschirnhaus, E. A. 2003. Method for removing all intermediate terms from a given equation. THIS BULLETIN, 37(1): 1-3. Original: Methodus auferendi omnes terminos intermedios ex data equatione, Acta Eruditorum 2, 204-207(1683).
- [20] Adamchik, V. S. Jeffrey, D. J. 2003. Polynomial transformations of tschirnhaus, bring and jerrard. ACM SIGSAM Bulletin 37(3): 90-94. <https://doi.org/10.1145/1093528.1093530>
- [21] Lazard, D. Solving quintics in radicals. In: Olav Arnfinn Laudal, Ragni Piene, The Legacy of Niels Henrik Abel. 1869, 207-225. Springer, Berlin, Heidelberg.
- [22] Dummit, D. S. 1991. Solving solvable quintics. Mathematics of Computation 57(1): 387-401.
- [23] Larbaoui, Y. New Theorems and Formulas to Solve Fourth Degree Polynomial Equation in General Forms by Calculating the Four Roots Nearly Simultaneously. American Journal of Applied Mathematics. vol. 11(6): 95-105. <https://doi.org/10.11648/j.ajam.20231106.11>
- [24] Pesic, P. Abel's Proof: An Essay on the Sources and Meaning of Mathematical Unsolvability, MIT Press, 2004.
- [25] Wantzel, P. L. 1845. Démonstration de l'impossibilité de résoudre toutes les équations algébriques avec des radicaux, J. Math. Pures Appl., 1re série, vol. 4, 57-65.