

Research Article

Nematicidal Potential of Neem (*Azadirachta indica* A. Juss) Based Products Against Root-Knot Nematode (*Meloidogyne incognita*) and Yield Enhancement in Tomato

Mohammed Iqbal Faruk^{1, *}, Shamim Akhter², Rummana Islam², Ferdous-E-Elahi², Tasfia Hassan Tabassum², Alamgir Siddiky¹

¹Regional Agricultural Research Station, Bangladesh Agricultural Research Institute, Cumilla, Bangladesh

²Plant Pathology Division, Bangladesh Agricultural Research Institute, Gazipur, Bangladesh

Abstract

Root-knot nematode (*Meloidogyne incognita*) is one of the most destructive soil-borne pathogens that severely affect tomato (*Solanum lycopersicum* L.) production worldwide including Bangladesh, causing significant yield losses. The present study was conducted to evaluate the nematicidal potential of neem (*Azadirachta indica* A. Juss) based products for the management of *M. incognita* and their effects on tomato growth and yield under both pot culture and field conditions during 2017-18, 2018-19, 2019-20 and 2020-21 cropping seas. Different neem formulations, including neem seed extract, neem leaf extract, neem oil, and commercial neem cake, were applied artificial root knot nematode infested soil amendment. In pot house condition, all neem based product was effective against root knot nematode and increased plant growth parameters. In the field condition, results revealed that all neem products significantly reduced root-knot gall index, and increased plant growth parameters such as shoot and root growth as well as fruit number and yield of tomato compared to the untreated control. Among the tested treatments, neem oil cake and neem seed extract treatment exhibited the best nematicidal activity, with a reduction in gall index and remarkable improvement in plant height, shoot weight, root weight, fruit number, and total yield of tomato. Neem leaf extract, neem leaf powder and chemical nematicide Furan 5G treatments also performed better in reduction of gall index, improving plant growth and yield of tomato compared to untreated control. These findings suggest that neem-based products, particularly neem oil cake and neem seed extract, are effective, eco-friendly alternatives to synthetic nematicides for sustainable root-knot nematode management and yield improvement in tomato production systems.

Keywords

Neem, *Meloidogyne incognita*, Tomato, Nematicidal Activity, Neem Cake, Yield Enhancement

*Correspondence: Mohammed Iqbal Faruk (mifaruk2012@yahoo.com)

Received: 30 September 2025; Accepted: 18 November 2025; Published: 23 April 2026



Copyright: © The Author(s), 2026. Published by Science Publishing Group. This is an **Open Access** article, distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Tomato (*Solanumlycopersicum* L.) belongs to the family Solanaceae is one of the most important vegetable crops cultivated worldwide for its high nutritive and economic value. It ranks next to potato and sweet potato in respect of vegetable production in the world [1, 2]. However, its productivity is severely constrained by plant-parasitic nematodes, particularly root-knot nematodes (*Meloidogyne* spp.), among which *Meloidogyne incognita* is the most prevalent and destructive species [3, 4]. These obligate endoparasites induce characteristic root galls that interfere with the normal uptake of water and nutrients, leading to stunted growth, chlorosis, wilting, and substantial yield reduction [5, 6]. Plant-parasitic nematodes have been implicated as a major constraint to tomato production [7] Yield losses in tomato caused by *M. incognita* have been estimated to range between 30–100%, depending on the level of infestation and environmental conditions [8].

Chemical nematicides such as carbofuran, fenamiphos, and oxamyl were once widely used to suppress nematode populations; however, their extensive use has led to several adverse consequences, including environmental pollution, residue hazards to other beneficial soil micro flora and risks to human and animal health, high cost, and development of resistance [9–11]. In view of these limitations, there is a growing emphasis on identifying safer, biodegradable, and environmentally friendly alternatives for nematode management. Environmentally friendly alternatives management like various soil amendments by green and compost manures, organic and inorganic fertilizers, botanicals, bio-control agents like nematophagous fungi, soil bacteria are found to be more satisfactory to reduce the nematode population and increased crop yield [12]. Among such alternatives, botanicals have attracted considerable attention due to their bioactive secondary metabolites with nematicidal, repellent, or antifeedant properties [13, 14].

Plant-derived products, especially botanicals such as Azadirachta, Eucalyptus, Chrommelina, *Sidaacuta* and Targetis, have gained attention as sustainable substitutes due to their biodegradability, low toxicity, and multiple modes of action against pests and pathogens [9, 15]. Neem (*Azadirachta indica* A. Juss), belonging to the family Meliaceae, is one of the most promising plant sources of natural nematicidal compounds. Its bioactive compounds, particularly azadirachtin, salannin, nimbin, and related limonoids, exhibit strong antifeedant, repellent, growth-disrupting, and nematicidaleffects [16, 17]. Various neem-based products such as neem seed cake, neem oil, neem leaf extract, and azadirachtin formulations have been shown to possess strong nematicidal activity against a wide range of plant-parasitic nematodes [18–20]. Moreover, neem products improve soil fertility and stimulate beneficial microbial populations, thereby enhancing overall plant vigor and yield [21, 22]. In addition, neem products are compatible with integrated pest management (IPM) strategies and contribute to soil health improvement. However, in spite of the

wide distribution of root-knot nematode on many crops in Bangladesh, little work has been done on the management of root-knot nematode in the country and so far, no efforts have been made to exploit locally available botanicals especially neem based products for the control of root-knot nematode. Considering the increasing demand for environmentally safe management practices, neem-based products represent a promising approach for the control of *M. incognita* in tomato cultivation. The present study was undertaken to assess the nematicidal potential of various neem-based products against *M. incognita* and to evaluate their influence on growth and yield performance of tomato.

2. Materials and Methods

A. Pot house experiment

The study was conducted in the pothouse of the Plant Pathology Division, BARI, Gazipur, during the Rabi season of 2017–18. Seven treatments were evaluated: i) Furdan (F) 5G @ 15 g pot⁻¹, ii) soil application of neem leaf powder @ 15 g pot⁻¹, iii) soil application of neem leaf extract (1:10 w/v), iv) soil application of neem seed extract (1:10 w/v), v) soil application of neem oil (1:10 v/v), vi) soil application of neem oil cake @ 100 g pot⁻¹, and vii) untreated control. The experiment followed a completely randomized design (CRD) with four replications. Pots were filled with sterilized loamy-sand soil. The sterilized soil was inoculated with chopped, severely galled roots of Indian spinach at a rate of 10 g kg⁻¹ of soil. Neem leaf extract, neem seed extract, and neem oil were applied twice—once at the time of transplanting and again 40–45 days after transplanting. Neem leaf powder and Furdan were applied only at transplanting, while neem cake was thoroughly mixed with the soil 21 days before pot filling to allow decomposition. Twenty-five-day-old tomato seedlings (cv. BARI Tomato-15) raised in sterilized soil were transplanted, with five seedlings per pot. Standard cultural practices, including weeding, fertilization, and irrigation, were performed as required. After 65 days of transplanting, plants were carefully uprooted to avoid damaging the root system and washed with running tap water for data recording.

B. Field experiment

The field experiment was carried out at the Plant Pathology Division, Bangladesh Agricultural Research Institute (BARI), Gazipur, over three consecutive Rabi seasons viz. 2018–19, 2019–20, and 2020–21. The study was laid out in a randomized complete block design (RCBD) with seven treatments and three replications. The treatments included: i) Furdan (F) 5G @ 45 kg ha⁻¹, ii) soil application of neem leaf powder @ 2–3 g plant⁻¹, iii) soil application of neem leaf extract (1:10 w/v), iv) soil application of neem seed extract (1:10 w/v), v) soil application of neem oil (1:10 v/v), vi) soil application of neem oil cake @ 600 kg ha⁻¹, and vii) untreated control. Each

unit plot measured 3 m × 2.5 m, with a one-meter spacing between adjacent plots. A plant-to-plant distance of 50 cm and a line-to-line distance of 60 cm were maintained. During seedling transplanting, root-knot nematode-infected roots of Indian spinach were chopped and incorporated into the soil at the rate of 1 g per plant. Neem-based treatments namely neem leaf powder, neem leaf extract, neem seed extract, and neem oil were applied twice: first at the time of seedling transplanting and again 40–45 days after transplanting. The chemical nematicide Furanan 5G was applied only once, at the time of transplanting. Neem oil cake was thoroughly mixed with the soil and kept moist for 21 days before transplanting to ensure proper decomposition. Tomato (*Solanum lycopersicum*) cultivar BARI Tomato-15 seedlings were raised in sterilized potting mix and transplanted at the four-leaf stage (4 weeks old). Additionally, ten extra seedlings were transplanted between two rows in each plot. Necessary intercultural operations, including weeding and irrigation, were carried out following the recommended practices for tomato cultivation [23].

C. Data collection and Statistical analysis

After 65 days of transplanting, ten additional plants from each plot were carefully uprooted, and their root systems were thoroughly washed under running tap water. Data were collected on shoot length, shoot weight, root length, and root weight. The severity of root galling was assessed using a 0–10 gall index scale as described by Zeck [24]. In addition, the number of fruits per plant and fruit yield were recorded from five randomly selected plants in each plot, and the fruit yield was expressed in tons per hectare (t/ha). Data were subjected to analysis of variance (ANOVA) appropriate to the RCBD using statistical software MSTATC program. Treatment means were separated using least significant difference (LSD) at $P \leq 0.05$.

3. Results and Discussion

A. Efficacy of Neem Products and Nematicide on Root-Knot Nematode in the Pot Experiment

The effect of different treatments on the suppression of root-knot nematode (*Meloidogyne incognita*) and the enhancement of tomato plant growth under pot house conditions is presented in Table 1. The average shoot length of tomato plants in the control treatment was 46.12 cm plant⁻¹. Application of different neem-based products and Furanan 5G significantly increased the shoot length, ranging from 62.17 to 71.00 cm plant⁻¹. The highest shoot length was observed in plants treated with neem seed extract and neem oilcake, followed by Furanan 5G, neem leaf extract, and neem leaf powder. Similarly, the highest shoot weight (191.90 g plant⁻¹) was obtained with neem seed extract treatment, followed by neem oilcake, neem leaf extract, Furanan 5G, and neem leaf powder. The control treatment produced the lowest shoot weight (118.00 g plant⁻¹). The maximum root length and root weight were recorded in the neem oilcake treatment, followed by neem seed extract, neem leaf extract, Furanan 5G, and neem leaf powder,

while the control exhibited the lowest values. Treatment with neem products and Furanan 5G markedly reduced the severity of root gall formation compared to the control. The highest gall index (8.46) was observed in the control, which was reduced to values between 4.17 and 5.88 in the treated plants, indicating effective suppression of *M. incognita* by neem-based treatments and Furanan 5G.

B. Efficacy of neem products and nematicide on root knot nematode in the field experiment

1) Severity of root-knot disease

In all experimental years, the severity of root-knot nematode infestation in tomato was significantly reduced compared to the control as a result of soil treatments with neem-based products—namely neem leaf powder, neem leaf extract, neem seed extract, neem oil, and neem oil cake—as well as Furanan 5G (Table 2). In the first year, the control plot recorded the highest average gall index value of 5.27. This was reduced to a range of 2.00–2.80 following treatments with neem-based products, neem oil cake, and Furanan 5G. The lowest disease severity was observed in soil treated with neem oil cake, followed by neem seed extract, neem leaf extract, neem leaf powder, Furanan 5G, and neem oil. The highest reduction in root-knot severity (62.05%) was achieved with neem oil cake, while reductions of 59.58%, 56.93%, 54.46%, 53.13%, and 46.87% were obtained with neem seed extract, neem leaf extract, neem leaf powder, Furanan 5G, and neem oil, respectively. In the second year, the control plot exhibited a maximum gall index of 4.27, which decreased to 1.60–2.33 under various treatments. All treatments significantly reduced disease severity compared to the control. The greatest reduction (62.53%) was again observed with neem oil cake, followed by neem seed extract (60.89%), neem leaf powder (51.52%), and neem leaf extract (50.12%). The least effective treatments were neem oil and Furanan 5G, with reductions of 47.78% and 45.43%, respectively (Table 2). In the third year, all treatments significantly lowered root-knot disease severity relative to the control. The highest gall index (4.51) was observed in the untreated control, while treated plots showed values between 1.16 and 1.98. Neem oil cake remained the most effective treatment, providing a 74.28% reduction in disease severity, followed by neem seed extract (68.29%), neem leaf extract (67.85%), neem leaf powder (61.19%), Furanan 5G (58.98%), and neem oil (56.09%) (Table 2). The reduction in nematode population through neem treatments aligns with previous findings. Javed et al. [19] reported that neem extracts suppress nematode population density due to their nematicidal properties. Similarly, Singh and Hali [25] demonstrated that neem seed extract has strong potential to control root-knot nematodes in tomato. Ntalli et al. [26] also confirmed the efficacy of azadirachtin-based formulations in suppressing *M. incognita* under greenhouse conditions. These results are further supported by the findings of Sharma et al. [27], Rather and Siddiqui [28], Ganai et al. [29], Satyandra et al. [30], Babu and Rana [31], Archana and Prasad [32], Resha and Rani [33], and Khan et al. [34] who reported that organic soil amendments,

including dried poultry litter, municipal refuse, and various oil cakes (groundnut, mustard, and neem), effectively suppress *Meloidogyne incognita*. Likewise, Akhtar [18] and Chitwood [10] confirmed the strong nematicidal potential of neem-based products against *M. incognita* and other plant-parasitic nematodes.

2) Shoot growth

Average shoot length of tomato plants under the control condition was recorded as 54.87 cm plant⁻¹ in the first year, 71.00 cm plant⁻¹ in the second year, and 63.45 cm plant⁻¹ in the third year (Table 3). Application of different neem-based soil amendments—namely neem oil cake, neem seed extract, neem leaf extract, neem leaf powder, neem oil, and Furadan 5G—significantly enhanced shoot length, which ranged from 72.27 to 83.67 cm plant⁻¹ in the first year, 89.93 to 105.10 cm plant⁻¹ in the second year, and 70.40 to 82.40 cm plant⁻¹ in the third year. In the first year, the highest shoot height was obtained from soil treatments with neem oil cake and neem seed extract, followed by neem leaf extract, neem leaf powder, Furadan 5G, and neem oil. A similar trend was observed in the second year, where neem oil cake, neem seed extract, and neem leaf extract recorded the tallest plants. During the third year, all treatments produced a comparable positive effect on shoot growth, with neem oil cake again resulting in the highest shoot height, closely followed by neem seed extract and neem leaf extract. The least effective treatments across the years were neem oil and Furadan 5G (Table 3). In terms of shoot weight, tomato plants under control conditions produced 178.7 g plant⁻¹ in the first year, which increased to 221.2–320.1 g plant⁻¹ with neem based products and Furadan 5G treatments. The highest increase in shoot weight was achieved with neem seed extract and neem oil cake, followed by neem leaf extract, neem leaf powder, and Furadan 5G, while neem oil showed the least effect. In the second year, the lowest shoot weight (230.7 g plant⁻¹) was recorded in the control, whereas the application of neem oil cake yielded the highest shoot weight (408.0 g plant⁻¹), followed by neem seed extract (392.7 g plant⁻¹) and neem leaf extract (389.3 g plant⁻¹). Neem oil remained the least effective, followed by neem leaf powder and Furadan 5G. In the third year, neem oil cake and neem seed extract again produced the highest shoot weights, while control and neem oil treatments yielded the lowest shoot weight (Table 3). The superior performance of neem oil cake may be attributed to its dual role as a biofertilizer and disease suppressant, which enhances nutrient availability and suppresses nematode infestation [35]. The present findings are consistent with those of Sharma et al. [27], who reported that neem product applications increased root and shoot growth, and with Pandey et al. [36], who found that neem seed cake significantly improved plant height and leaf number. Similarly, Kumar and Khanna [37] observed enhanced plant growth with aqueous neem extracts, while Javed et al. [38] demonstrated that neem formulations, applied both as foliar sprays and soil treatments, provided systemic protection against *Meloidogyne incognita*, leading to greater vegetative growth and biomass

accumulation in tomato.

3) Root growth

Amendment of soil with neem oil cake, neem seed extract, neem leaf extract, neem leaf powder, neem oil, and the application of Furadan 5G showed positive effects on the root growth of tomato compared to the control (Table 4). In the first year, the minimum root length of 13.80 cm plant⁻¹ was recorded in the control, while the maximum root lengths of 21.93 cm plant⁻¹ and 21.33 cm plant⁻¹ were obtained from treatments with neem oil cake and neem seed extract, respectively. Neem leaf extract also increased root length to 19.20 cm plant⁻¹. Other treatments, such as neem leaf powder, neem oil, and Furadan 5G, enhanced root length within the range of 17.20–18.13 cm plant⁻¹ over the control. A similar trend was observed in the second year. The maximum root length (27.67 cm plant⁻¹) was recorded with neem oil cake, followed by neem seed extract (27.40 cm plant⁻¹) and neem leaf extract (25.67 cm plant⁻¹). The lowest value (20.27 cm plant⁻¹) was observed in the control. Neem leaf powder, neem oil, and Furadan 5G also increased root length compared to control, though their effects were significantly lower than the top-performing treatments. In the third year, root length under control was 17.80 cm plant⁻¹, which increased to 20.25–28.60 cm plant⁻¹ with different treatments (Table 4). Root weight under control was 17.60 g plant⁻¹ in the first year, 27.00 g plant⁻¹ in the second year, and 24.65 g plant⁻¹ in the third year. Soil amendment with neem-based products and Furadan 5G increased the root weight ranged from 22.76–28.39 g plant⁻¹, 31.87–39.13 g plant⁻¹, and 26.35–35.00 g plant⁻¹ in the first, second, and third years, respectively (Table 4). These findings are consistent with those of Sharma et al. [27], who reported that the application of neem products enhanced root length in tomato. Similar findings were also reported by several researchers. Akhtar and Alam [39] observed that neem products significantly reduced nematode populations and improved root and shoot growth in tomato. Khan et al. [40] also found neem oil cake to enhance root biomass and reduce root-knot severity in *M. incognita*-infested tomato plants. Furthermore, Gowda et al. [41] reported that neem leaf extract-treated tomato plants exhibited greater root length and dry weight compared to untreated controls, confirming the plant growth-promoting potential of neem-derived compounds.

4) Crop yield

Soil treatment with neem-based products namely neem oil cake, neem seed extract, neem leaf extract, neem leaf powder, and neem oil along with Furadan 5G, consistently enhanced fruit number per plant and fruit yield per hectare across all three years of study (Tables 5 and 6). Under control conditions, the fruit number per plant was 23.67 in the first year, 29.87 in the second, and 27.50 in the third year (Table 5). However, with neem-based products and Furadan treatments, fruit number increased remarkably to 31.33–37.67, 31.73–42.00, and 31.50–39.50 in the first year, second year and third year, respectively. In the first year, neem oil cake treatment produced the highest increased by 37.16% more fruits per plant than the

control followed by neem seed extract, neem leaf extract, and Furadan 5G. A similar pattern was observed in the subsequent years, where neem oil cake continued to outperform other treatments. In the second year, fruit number rose by 28.88% under neem oil cake treatment compared to control, while neem seed extract and neem leaf powder increased fruit number by 27.97% and 26.43%, respectively (Table 5).

Regarding fruit yield, control plots produced the lowest yields: 40.09 tha⁻¹ in the first year, 45.47 tha⁻¹ in the second year, and 43.33 tha⁻¹ in the third year (Table 6). Application of neem-based treatments and Furadan 5G increased yields substantially. In the first year, yields ranged from 50.00 to 68.53 t/ha, with neem oil cake achieving the highest yield (68.53 tha⁻¹), followed by neem seed extract (64.07 tha⁻¹). This represented a 41.50% yield increase over the control, with other treatments showing gains of 37.43% (neem seed extract), 26.84% (neem leaf powder), and 26.48% (neem leaf extract). Neem oil and Furadan 5G were less effective, with yield increases of 19.82% and 20.46%, respectively compared to control. In the second year, treated plots yielded 57.20–69.79 tha⁻¹, with neem oil cake again producing the highest yield, showing a 34.85% yield increase over control, followed closely by neem seed extract, neem leaf extract, neem leaf powder, and Furadan 5G with increases of 33.32%, 32.92%, 31.58%, and 28.63%, respectively over control. In the third year, the highest yield of 69.58 tha⁻¹ was achieved with neem oil cake, followed by neem seed extract (66.25 tha⁻¹), neem leaf extract (65.83 tha⁻¹), and neem leaf powder (61.25 tha⁻¹). The control plot produced only 43.33 t/ha. Once again, neem oil and Furadan 5G were the least effective producing 55.83 and 57.92 tha⁻¹, respectively, though they still increased yield by 21.44% and 25.19%, respectively (Table 6). The consistent improvement in fruit number and yield across years suggests that neem-based amendments effectively suppress root-knot nematodes, which are known to hinder plant growth and yield [19]. These findings align with earlier studies—Saravanapriya and Sivakumar [42] found that neem and other botanicals significantly boosted yield over untreated controls, while Khan et al. [40] reported that neem leaf application increased fruit number and weight in eggplant due to improved soil organic content. Similarly, Alam et al. [13] and Adegbite&Adesiyan [43] observed that neem cake treatments not only reduced

nematode infestation but also significantly enhanced tomato fruit weight and yield.

Correlation and regression analyses further confirmed the impact of neem treatments. The relationship between gall index and fruit yield, shoot weight, and root weight was linear and negative, with correlation coefficients (*r*) of 0.820, 0.673, and 0.755, respectively (Figure 1). The effects were significant, indicating that gall index accounted for 67.30% ($R^2 = 0.673$) of the variation in fruit yield, 45.30% ($R^2 = 0.453$) in shoot weight, and 57.00% ($R^2 = 0.570$) in root weight. These results demonstrate that neem-based soil amendments not only suppress nematode damage but also improve overall plant growth, likely due to the addition of essential nutrients and organic matter to the soil.

4. Conclusion

Although all treatments performed better than the control, neem oil cake and neem seed extract proved to be the most effective. These treatments significantly enhanced plant growth and yield parameters—including plant height, number of leaves, number of fruits, and fruit weight—while also reducing the soil nematode population and the number of root galls. Therefore, neem oil cake and neem seed extract can be recommended as suitable alternatives to synthetic nematicides for root knot nematode management.

Acknowledgments

The authors thankfully acknowledged to Bangladesh Agricultural Research Institute, Gazipur to provide financial support and logistic support for the pieces of research. Thanks go to Mr. Md. AbdurRazzak and Mr. ZamilAkter (Scientific Assistant) for their assistance for successfully complete the research in the field.

Conflicts of Interest

No potential conflict of interest was reported by the authors.

Appendix

Table 1. Effect of neem product and nematicide on the root knot disease incidence of tomato in pothouse.

Neem based products and Furadan 5G with dose	Shoot length (cm)	Shoot weight (gplant-1)	Root length(cm)	Root weight (gplant-1)	Gall index (0-10 scale)
Furadan 5G @15 gpot ⁻¹	68.42 ab	172.30 ab	22.00 d	33.50 bc	5.09 b
Neem leaf powder @ 15 gpot ⁻¹	68.06 ab	170.60 ab	26.46 bc	33.08 bc	4.33 c
Neem leaf extract (1:10 w/v)	68.38 ab	187.00 ab	27.25 bc	33.33 bc	4.17 d

Neem based products and Furadan 5G with dose	Shoot length (cm)	Shoot weight (gplant ⁻¹)	Root length(cm)	Root weight (gplant ⁻¹)	Gall index (0-10 scale)
Neem seed extract (1:10 w/v)	71.00 a	191.90 a	28.67 b	34.92 ab	4.29 d
Neem oil (1:10 v/v)	62.17 b	155.50 b	24.75 cd	31.25 bc	5.88 b
Neem oilcake @100gpot ⁻¹	69.33 a	175.80 ab	32.92 a	38.25 a	4.33 d
Control	46.12 c	118.00 c	17.25 e	30.25 c	8.46 a
LSD (P=0.05)	5.811	30.46	3.084	4.139	0.571

In a column, similar letter (s) do not differ significantly at 5% level of probability

Table 2. Efficacy of soil treatment with neem based products and nematicide on the severity of root -knot disease (*Meloidogyne incognita*) of tomato in three consecutive years.

Neem products and Furadan 5G with dose	Gall Index (0-10 scale)			Reduction of gall index over control (%)		
	2016-17	2017-18	2018-19	2016-17	2017-18	2018-19
Furadan 5G @45 kg ha ⁻¹	2.47 bc	2.23 b	1.85 bc	53.13	47.78	58.98
Neem leaf powder @ 2-3 g plant ⁻¹	2.40 bcd	2.07 bc	1.75 bc	54.46	51.52	61.19
Neem leaf extract (1:10 w/v)	2.27 cd	2.13 bc	1.45 cd	56.93	50.12	67.85
Neem seed extract (1:10 w/v)	2.13 cd	1.67 c	1.43 cd	59.58	60.89	68.29
Neem oil (1:10 v/v)	2.80 b	2.33 b	1.98 b	46.87	45.43	56.09
Neem oilcake @500 kg ha ⁻¹	2.00 d	1.60 c	1.16 d	62.05	62.53	74.28
Control	5.27 a	4.27 a	4.51 a	-	-	-
LSD (P=0.05)	0.389	0.530	0.501	-	-	-

Values within the same column with a common letter do not differ significantly (P=0.05)

Table 3. Effect of soil treatment with neem based products and a nematicide on shoot growth of tomato in three consecutive years.

Neem products and Furadan 5G with dose	Shoot height (cm plant ⁻¹)			Shoot weight (g plant ⁻¹)		
	2016-17	2017-18	2018-19	2016-17	2017-18	2018-19
Furadan 5G @45 kg ha ⁻¹	72.87 b	90.67 b	70.65 bc	248.9 c	349.3 c	201.4 b
Neem leaf powder @ 2-3 g plant ⁻¹	74.13 b	92.07 b	73.47 ab	277.5 b	356.3 c	207.2 b
Neem leaf extract (1:10 w/v)	76.80 b	101.70 a	75.70 ab	286.2 b	389.3 b	221.5 b
Neem seed extract (1:10 w/v)	83.33 a	102.80 a	78.05 ab	320.1 a	392.7 b	247.2 a
Neem oil (1:10 v/v)	72.27 b	89.93 b	70.40 bc	221.2 d	348.0 c	198.7 bc
Neem oilcake @500 kg ha ⁻¹	83.67 a	105.10 a	82.40 a	314.6 a	408.0 a	262.3 a
Control	54.87 c	71.00 c	63.45 c	178.7 e	230.7 d	173.8 c
LSD (P=0.05)	5.736	6.449	9.572	20.10	9.507	25.45

Values within the same column with a common letter do not differ significantly (P=0.05)

Table 4. Efficacy of soil treatment with neem based products and a nematicide on root growth of tomato in three consecutive years.

Neem products and Furadan 5G with dose	Root length (cmplant ⁻¹)			Root weight (gplant ⁻¹)		
	2016-17	2017-18	2018-19	2016-17	2017-18	2018-19
Furadan 5G @45 kg ha ⁻¹	18.07 c	23.33 c	23.35 bc	23.91 b	33.47 bc	30.10 abc
Neem leaf powder @ 2-3 gplant ⁻¹	18.13 c	23.67 c	24.75 abc	22.76 b	38.00 a	32.05 ab
Neem leaf extract (1:10 w/v)	19.20 b	25.67 b	25.15 ab	23.00 b	35.00 b	32.25 ab
Neem seed extract (1:10 w/v)	21.33 a	27.40 ab	26.55 ab	26.88 a	37.40 a	33.25 a
Neem oil (1:10 v/v)	17.20 c	23.53 c	20.25 cd	22.85 b	31.87 c	26.35 bc
Neem oilcake @500 kg ha ⁻¹	21.93 a	27.67 a	28.60 a	28.39 a	39.13 a	35.00 a
Control	13.80 d	20.27 d	17.80 d	17.60 c	27.00d	24.65c
LSD (P=0.05)	1.057	1.839	4.581	1.585	1.811	6.791

Values within the same column with a common letter do not differ significantly (P=0.05)

Table 5. Efficacy of soil treatment with neem based products and a nematicide on the fruit number per plant of tomato in soil inoculated with *Meloidogyne incognita*.

Neem products and Furadan 5G with dose	Number of fruits plant ⁻¹			Number of fruits increased over control (%)		
	2016-17	2017-18	2018-19	2016-17	2017-18	2018-19
Furadan 5G @45 kg ha ⁻¹	33.00 bc	35.93 c	36.25 ab	28.27	16.87	24.14
Neem leaf powder @ 2-3 gplant ⁻¹	32.67 bc	40.60 ab	37.25 a	27.55	26.43	26.17
Neem leaf extract (1:10 w/v)	33.00 bc	38.53 b	37.25 a	28.27	22.48	26.17
Neem seed extract (1:10 w/v)	36.33 ab	41.47 a	38.25 a	34.85	27.97	28.11
Neem oil (1:10 v/v)	31.33 c	31.73 d	31.50 bc	24.45	05.86	12.70
Neem oilcake @500 kg ha ⁻¹	37.67 a	42.00 a	39.50 a	37.16	28.88	30.38
Control	23.67 d	29.87 d	27.50 c	-	-	-
LSD (P=0.05)	3.41	2.072	5.505	-	-	-

Values within the same column with a common letter do not differ significantly (P=0.05)

Table 6. Efficacy of soil treatment with neem based products and a nematicide on the fruit yield of tomato in soil inoculated with *Meloidogyne incognita*.

Neem products and Furadan 5G with dose	Fruits yield (tha ⁻¹)			Yield increased over control (%)		
	2016-17	2017-18	2018-19	2016-17	2017-18	2018-19
Furadan 5G @45 kg ha ⁻¹	50.40 d	63.71 c	57.92 bc	20.46	28.63	25.19
Neem leaf powder @ 2-3 gplant ⁻¹	54.80 c	67.79ab	61.25abc	26.84	32.92	29.26
Neem leaf extract (1:10 w/v)	54.53 c	66.46b	65.83ab	26.48	31.58	34.18
Neem seed extract (1:10 w/v)	64.07 b	68.19 ab	66.25 ab	37.43	33.32	34.60
Neem oil (1:10 v/v)	50.00 d	57.20 d	55.83 c	19.82	20.50	21.44

Neem products and Furadan 5G with dose	Fruits yield (tha ⁻¹)			Yield increased over control (%)		
	2016-17	2017-18	2018-19	2016-17	2017-18	2018-19
Neem oilcake @500 kgha ⁻¹	68.53 a	69.79 a	69.58 a	41.50	34.85	37.73
Control	40.09 e	45.47 e	43.33 d	-	-	-
LSD (P=0.05)	3.84	2.359	8.921	-	-	-

Values within the same column with a common letter do not differ significantly (P=0.05)




Figure 1. Relationship of shoot weight, root weight and fruit yield with gall index of tomato grown in soil inoculated with *Meloidogyne incognita* and treated with poultry neem oil cake, neem seed extract, neem leaf extract, neem leaf powder, neem oil and Furadan 5G.

References

- [1] Food and Agriculture Organization. 1997. FAO production years books. Basis Data Unit. Statistics Div. FAO Rome Italy, 21: 125-127.
- [2] Hossain, M. E., M. J. Alam, M. A. Hakim, A. S. M. Amanullah and Ahsanulla. 2010. An assessment of Physicochemical Properties of Some Tomato Genotypes and Varieties Grown At Rangpur. Bangladesh Res. Pub. J. 4(3): 135-243.
- [3] Sasser, J. N. 1989. Plant Parasitic Nematodes: The Farmer's Hidden Enemy. North Carolina State University, Raleigh, North Carolina. 7-14.
- [4] Karssen, G. and M. Moens. 2006. Root-knot nematodes. In: Perry, R. N. and Moens, M. (eds.), *Plant Nematology*, CABI Publishing, Wallingford, pp. 59-90.
- [5] Dropkin, V. H. 1992. *Introduction to Plant Nematology*, 2nd ed. John Wiley & Sons, New York.
- [6] Moens, M., R. N. Perry and J. L. Starr. 2009. *Meloidogyne* species - a diverse group of novel and important plant parasites. In: Perry, R. N. et al. (eds.), *Root-Knot Nematodes*, CABI, Wallingford, pp. 1-17.
- [7] Luc, M., R. A. Sikora and J. Bridge. 2005. Plant-parasitic nematodes in subtropical and tropical agriculture - Second edition. CABI Publishing, Wallingford, UK: pp 331-871.
- [8] Anwar, S. A. and M. V. McKenry. 2010. Incidence and reproduction of *Meloidogyne incognita* on vegetable crop genotypes. *Pakistan Journal of Zoology*, 42(2), 135-141.
- [9] Akhtar, M. and A. Malik. 2000. Roles of organic soil amendments and soil organisms in the biological control of plant-parasitic nematodes: a review. *Bioresource Technology*, 74: 35-47.
- [10] Chitwood, D. J. 2002. Phytochemical-based strategies for nematode control. *Annual Review of Phytopathology*, 40: 221-249.
- [11] Wachira, P. M., J. W. Kimenju, S. A. Okoth and R. K. Mibey. 2009. Stimulation of Nematode-Destroying Fungi by Organic Amendments Applied in Management of Plant Parasitic Nematode. *Asian J. of Plant Sci.* 8: 153-159.
- [12] Goswami, B. K. and S. Neetu. 2012. Combat the Soil Borne Fungal and Root Knot nematode Diseases Infecting Horticultural Crops. *Amity CJBiopest*, 5: 255-258.

- [13] Adegbite, A. A. and S. O. Adesiyun. 2005. Root extracts of plants to control root-knot nematode on edible soybean. *World Journal of Agricultural Sciences*, 1(1), 18-21.
- [14] Ntalli, N. G. and P. Caboni. 2012. Botanical nematicides: A review. *Journal of Agricultural and Food Chemistry*, 60(40): 9929-9940.
- [15] Umar, I., Z. Muhammad and B. A. O Okusanya. 2010. Effect of organic amendments on the control of *Meloidogyne javanica* (Kofoid and White, 1999) Chitwood, 1949) on tomato (*Lycopersicon lycopersicum*, Mill). *Agric., Business Tech.* 8: 63-77.
- [16] Isman, M. B. 2006. Botanical insecticides, deterrents, and repellents in modern agriculture and an increasingly regulated world. *Annual Review of Entomology*, 51: 45-66.
- [17] Karyne, A., R. S. Braga and C. S. Cavalcante. 2013. Nematicidal activity of neem oil and neem cake against *Meloidogyne incognita*. *Journal of Medicinal Plants Research*, 7(22): 1626-1632.
- [18] Akhtar, M. 2000. Nematicidal potential of the neem tree (*Azadirachta indica*) and its products. *Integrated Pest Management Reviews*, 5, 57-66.
- [19] Javed, N., S. El-hassan, S. Gowen, B. Pemproke and M. Inam-Ul-haq. 2008. The potential of combining *Pasteuria penetrans* and neem (*Azadirachta indica*) formulations as a management system for root knot nematodes on tomato. *Eur J Plant Pathol.* 120: 53-60.
- [20] Oka, Y., S. Nacar, E. Putievsky, U. Ravid, Z. Yaniv and Y. Spiegel. 2000. *Nematicidal activity of essential oils and their components against the root-knot nematode*. *Phytopathology*, 90: 710-715.
- [21] Akhtar, M. and I. Mahmood. 1994. *Potentiality of phytochemicals in nematode control: A review*. *Bioresource Technology*, 48, 189-201.
- [22] Hussain, M. A., T. Mukhtar and M. Z. Kayani. 2016. "Management of Root-Knot Nematode (*Meloidogyne incognita*) in Tomato through the Use of Biofertilizers and Organic Amendments." *Pakistan Journal of Zoology* 48 (5): 1353-1358.
- [23] Anonymous. Research Report on Horticultural Crops (2006-07). Horticulture Research Centre, BARI, Joydebpur, Gazipu, 2007: 91p.
- [24] Zeck, M. W. 1971. A rating scheme for field evaluation of root knot nematode infestation. *Planzenschutz-Nacht*, 24: 141-144.
- [25] Singh, K. and S. Hali. 2017. Nematicidal Efficacy of Neem (*Azadirachta indica*, A. Juss) Seed Extract against *Meloidogyne incognita*. *International Journal of Plant & Soil Science*, 18(3): 1-6.
- [26] Ntalli, N. G., S. Vargiu, U. Menkissoglu-Spiroudi and P. Caboni. 2009. Nematicidal activity of natural compounds from neem tree (*Azadirachta indica* A. Juss) against *Meloidogyne incognita*. *Journal of Agricultural and Food Chemistry*, 57(3), 579-583.
- [27] Sharma, H. K., S. Singh and J. Pankaj. 2007. Management of *Meloidogyne incognita* with *Paecilomyces lilacinus* and neem cake on okra. *Pestic Res J.* 19: 166-168.
- [28] Rather, M. A. and M. A. Siddiqui. 2007. Neem for the control of root-knot nematode infecting tomato. 37(1): 81-110.
- [29] Ganai, A. M., B. Rehman, P. Kavita, M. Asi and M. A. Siddiqui. 2014. Phytotherapeutic approach for the management of *Meloidogyne incognita* affecting *Abelmoschus esculentus* (L.) Moench. *Archives of Phytopathology and Plant Protection* 47: 1797-1805.
- [30] Satyandra, S., A. B. Rai and R. K. Sing. 2011. Bio-management of root-knot disease of Chili (*Capsicum annum*) caused by *Meloidogyne incognita*. *Vegetable Science*, 38(1): 63-67.
- [31] Babu, and L. Rana. 2012. Efficacy of neem products as seed treatment against *Meloidogyne incognita* in Okra. *Annals of Plant Protection Sciences* Volume: 20: 1; 268- 269.
- [32] Archana, U. S. and D. Prasad. 2014. Management of Plant-parasitic Nematodes by the Use of Botanicals. *J Plant PhysiolPathol*, 2: 1.
- [33] Resha, M. and D. S. Rani. 2015. Management of root Knot Nematode (*Meloidogyne Incognita*) using Neem (*Azadirachta Indica*). *IOSR Journal of Environmental Science, Toxicology and Food Technology*, 9(3): 12-15.
- [34] Khan, F., N. Javed, and M. Kamran. 2019. Evaluation of the nematicidal potential of some botanicals against root-knot nematode (*Meloidogyne incognita*) infecting tomato. *Pakistan Journal of Nematology*, 37(1): 59-64.
- [35] Siddiqui, Z. A., M. S. Akhtar and K. Futai. 2007. Biological control of plant-parasitic nematodes by fungi: A review. *Bio-control Science and Technology*, 17(8): 837-852. <https://doi.org/10.1080/09583150701593930>
- [36] Pandey, G., R. K. Pandey and H. Pant. 2005. Influence of organic amendments on nematode fauna and microflora of chick-pea rhizosphere. *Indian J Pulses Res.* 18: 263-264.
- [37] Kumar, S. and A. S. Khanna. 2006. Efficacy of *Trichoderma viride* and neem cake against root-knot nematode, *Meloidogyne incognita* on tomato. *Pest Manag Econ Zool.* 14: 103-106.
- [38] Javed, N., S. R. Gowen, S. A. El-Hassan, M. Inam-ul-Haq and F. Shahina. 2007. Systemic and persistent effect of neem (*Azadirachta indica*) formulations against root-knot nematodes, *Meloidogyne javanica*, on tomato. *Crop Protection*, 26: 36-43.
- [39] Akhtar, M., and M. M. Alam. 1990. *Control of plant-parasitic nematodes with organic and inorganic soil amendments*. *Plant and Soil*, 132, 195-199.
- [40] Khan, M. R., R. D. Solanki, B. Bohra and B. N. Vyas. 2012. Evaluation of Ahook (Azadirachtin 1500ppm) against root knot nematode (*Meloidogyne incognita*) infecting okra. *South Asian Journal of Experimental Biology*, 2.
- [41] Gowda, D. N., M. N. Sreenivasa, and K. S. Jagadeesh. 2018. *Effect of neem-based products on suppression of root-knot nematode and growth promotion in tomato*. *Journal of Pharmacognosy and Phytochemistry*, 7(5): 1289-1294.

- [42] Saranavanapriya, B. and M. Sivakumar. 2005. Management of root-knot nematode, *Meloidogyne incognita* on tomato with botanicals. *Natural Product Radiance*. 4: 158-161.
- [43] Alam, M. M., M. Ahmad and A. M. Khan. 1979. Effect of organic amendments on the growth and chemical composition of tomato, eggplant and chilli and their susceptibility to attack by *Meloidogyne incognita*. *Plant and Soil*, 51: 231-236.