

Review Article

Influences of Salient Agronomic Practices for Mitigation of Biennial Bearing in Coffee (*Coffea arabica* L.)

Dawit Merga^{*}, Daba Etana

Ethiopian Institute of Agricultural Research, Jimma Agricultural Research Center, Jimma, Ethiopia

Email address:

dawitmerga@gmail.com (Dawit Merga)

^{*}Corresponding author**To cite this article:**Dawit Merga, Daba Etana. Influences of Salient Agronomic Practices for Mitigation of Biennial Bearing in Coffee (*Coffea arabica* L.).*Journal of Plant Sciences*. Vol. 11, No. 4, 2023, pp. 136-143. doi: 10.11648/j.jps.20231104.15**Received:** June 21, 2023; **Accepted:** July 10, 2023; **Published:** August 9, 2023

Abstract: *Coffea arabica* L. is a perennial cash crop which is highly affected by biennial bearing. Most of the time, agronomic practices are applied with the objectives of boosting coffee productivity and production. But very less emphasis on their role of mitigating coffee yield bienniality. Most literatures and scholars didn't clearly and directly indicate the significance of agronomic practices in diminishing oscillation productivity of Arabica coffee. Thus, this review was conducted to assess the importance of salient agronomic practices in alleviating biennial bearing of Arabica coffee and to identify the existing gap by generating biennial intensity analysis from different findings. Salient agronomic practices such as fertilizer application, cover crop, shade, plant density and pruning methods were articulated in this article. These practices contributed in increasing productivity and mitigating bienniality via improving physiological jeopardize agents such as branch die-back, profuse flower bud, plant nutrient, over bearing of berries and rapid exhaustion of productivity parts and environmental stress. It was elucidated that sole field management was less efficient for biennial bearing mitigation than integrated management practices. The integrated agronomic practices that boost coffee yield need to be supported by non-biennial bearing variety for complete abolish of bienniality. Currently, there is no non-biennial bearing Arabica coffee variety. Thus, coffee genetic improvement for bienniality is highly recommended in future coffee breeding program.

Keywords: Agronomic Practices, Bienniality, *Coffea arabica* L., Integrated and Productivity

1. Introduction

Perennial crops' production affected by different factors which are manageable and irrepressible. Once established, they persist for long years giving yield [1, 2]. Commencement from planting up to their senescence stage, their growth and performance in economic traits influenced by different scenarios of endogenic and extraneous factors such as climate change, disease, insect pest and edaphic problems [3]. These may cause genetic erosion, yield reduction or fluctuation [4] and quality lose which totally result food security problem in the world if not solved scientifically.

Alternate bearing is one of the most complex and principal problem in perennial crop production. Many fruit trees and other horticultural crops' production reduced due to biennial effects. For instance, Mango, Apple, Pear, Apricot and

Avocado are highly affected by bienniality [5-7]; but, Grape biennial bearing is negligible [8].

Arabica coffee is one of the most affected cash crop by bienniality in the world including Ethiopia. As a result of this, the *Coffea arabica* L. yield is fluctuating or higher one year and lower the next [9, 7]; this of course affects farmers' annual incomes [10]. The intermittent of harvesting coffee yield consequences food security problem especially in developing countries which relied on coffee production. Also, the producers' on and off harvesting affect total coffee production which disturb supply and demand in the world market [10]. Currently, the oscillation of Arabica coffee production became bottleneck for world's coffee industries.

In yield biennial pattern, over load of berries cropping year followed by very low yield bearing season. Over bearing of berries affects coffee quality via lessening fruit size and

disturbing physiological process [11] during fruit development and physiological maturity. Also, the break of branches and disease severity such as coffee leaf rust is serious during on year [5]; heavy crop results branch-dieback [12] which leads to biennial production cycle [11]. Additionally, by the depletion of the reserves of assimilation in on year [13], the coffee trees may be exposed to jeopardize such as complete death. Thus, focus need to be taken to apply appropriate agronomic practices and to develop null in yield biennial coffee variety for mitigation of biennial bearing and production.

Biennial bearing cycle in *Coffea arabica* L. can be emanates from growth and fruit habit itself, environmental factors and management practices (shade, stumping and fertilization) [14, 15]. Arabica coffee exhibits biennial growth and allocate photosynthetic product to fruit formation and growth during year of high production and vegetative growth during years of low production. Shade and pruning alleviate the biennial problem via avoiding overbearing [11] and encouraging renewing bearing surface or growing productive lateral branches that enable producing fruit respectively [16]. Montoya and Umana [17] found reduction of branch die-back to 66% using 200g N per plant in relation to coffee trees to which no N fertilizer applied; this treatment directly reduces yield bienniality. Thus, in addition to developing regular bearing variety, mitigating biennial bearing by management practices is highly momentous in order to frequently harvesting yield from high yielder variety that has bienniality nature. Sustainable coffee yield harvest using integrated technologies is decisive to generates continues annual income for producers which contributes in realizing food security; also, it solves unbalanced supply and demand problem in the world coffee industry. Therefore, in the current situation many agronomic practices are underutilization in huge coffee farming for increasing coffee productivity and production; but there is less understanding on their advantage for combating biennial bearing in Arabica coffee. Most findings and articles emphasis and reported the role of agronomic practices on increasing coffee productivity and production and climate change resilience than their direct and/or indirect contribution in biennial bearing mitigation. Their role in alleviating alternate bearing of Arabica coffee is not clearly and directly indicated in many literatures. Thus, the current review conducted with the intension to know the significance of salient agronomic practices in biennial bearing alleviation and to identify the gap for mitigating alternate bearing of Arabica coffee.

2. Agronomic Practices for Alleviating Bienniality in Arabica Coffee

2.1. Population Density

In Ethiopia, the recommended coffee population density and most widely practiced by farms or users is 2500 trees ha⁻¹; But most of the world's coffee producers plant less than 2000 tree ha⁻¹ [11]. Numerous findings authenticated that closely

planting coffee trees is more preferable than planting with high spacing [18, 19]; but the canopy nature has to be taken in to consideration. The densely planted coffee productivity is higher than that of traditional panting [11]. Also, Pereira et al. [20] and Silveira et al. [21] reported that closer planting increases Arabica coffee productivity.

Plantation of high population density is one of the most important agronomic practice which is highly applicable for compact growth habit coffee varieties/cultivars. This practice presents lower production per plant, but increase production per area [22]. It reduces coffee productivity fluctuation by lessening stress to coffee plant; it enhances more leaf growth, provides suitable microclimate via decreasing air temperature inside coffee trees relative to external environment [22]. Additionally, due to high mutual shading, dense planting decreases over floral initiation, heavy bearing and branch die-back [11] which are important in mitigation of biennial production cycle.

2.2. Integrated Spacing with Pruning Methods

Plant space directly related to population density; it increases or decreases the interior between coffee trees and between rows. Applying closer space planting coffee trees or high density can upsurge productivity per area [22- 24]. Integrating such agronomic practice with pruning methods boost coffee productivity and production [25]. Agronomic practice such as pruning encourage new branch growth and new flower bud formation on newly grown branches [25] which is important in alleviating alternate bearing; also, it helps continuous yield harvest by removing criss-cross and unproductive branch, lean, lanky and whippy wood. These activities are very important to generate uninterrupted income for smallholder farmers and other coffee producers. Coffee productivity and production fluctuation affect growers sustainable income and disturb world coffee industry. The integrated agronomic practices such as spacing and pruning methods improve on and off bearing of Arabica coffee. Coffee trees treated with treatments seven (T7) reduced biennial intensity below 0.1 (Figure 1B) and it gave the highest yield next to treatment three (T3).

Also, despite low yield recorded in treatments (T1, T4 and T5), the biennial bearing was managed to almost null or negligible level. The highest yield was harvested from plots of T3 that planted 62×62 consisting 2652 coffee plants and treated by training on single stem with multiple stems on middle plant (Figure 1A). In agreement, the closer spacing lead to high trees density which enhances production increment per unit areas [24, 26]. But high on and off in bearing or high yield fluctuation observed in plot treated by T3 (Figure 1B). This pointed out field management systems need to be integrated with the genetically improved technology viz non-biennial bearing coffee variety. Mansood et al. [27] confirmed the importance of regular bearing variety to overcome bienniality in Pistachio fruit.

Sole spacing is not effective in mitigation of biennial bearing in Arabica coffee (Figure 2). From pooled yield of over five seasons, the biennial intensity recorded from 0.23 to

0.46. Likewise, Luiz and Aledir [28] reported that high yield fluctuation across years using four different spacing without integrate use of other field practices. Thus, the highest biennial intensity (0.46) observed from closer spacing (1m×0.5m); but the highest yield was recorded across harvesting and over five years mean yield for this treatment. In

contrast, Gokavi et al. [25] obtained biennial intensity less than 0.1 using combined spacing with four pruning methods. Hence, using integrated agronomic practices are more efficient in managing or minimizing yield biennial intensity relative to sole agronomic practice.




Figure 1. Influence of agronomic practices (Spacing and Pruning methods) on Yield and Biennial bearing.

Source: Gokavi et al. [25]

T1 -Square system of planting (62×62) + training on single stem + regular light pruning (Control) (1210 plants acre-1), T2 -Square system at close spacing (52×52) + training on single stem + Rock-n-roll pruning of alternate rows once 3-4 crops (1742 plants acre-1), T3- Square system of planting (62×62 Quincunx) + training on single stem + multiple stem on middle plant (2652 plants acre-1), T4 -Square system at close spacing (52×52) + training on multiple stem without topping + cyclic pruning after each harvest (1742 plants acre-1), T5- Hedge row system on single stem (62×32) + Rock-n-roll pruning of alternate rows (2420 plants acre-1), T6- Hedge row system on multiple stem without topping (62×32) + Cyclic pruning after each harvest (2420 plants acre-1), T7- Paired row system 62×32×72 for tall Arabica + single stem training + Rock-n-roll pruning alternate rows (1452 plants acre-1)




Figure 2. Role of spacing alone in alleviation of biennial bearing.

S- Represent spacing, S1- 3m×1m, S2- 2m×1m, S3- 2m×0.5m, S4- 1m×0.5m, I1-I4 –represent biennial intensity over two years, three years, four years and five years respectively; Source: Luiz and Aledir [28]

2.3. Integrated Number of Stems Per Plant with Pruning Methods

Science the commencement of research on increasing coffee yield, agronomist and breeders have been doing many field management practices and genetic improvement

respectively. Fostering the growth of one or more number of stem per plant is among the prominent agronomic practices for coffee productivity increment. Integrated pruning with number of stem per plant is practiced for increasing coffee yield [29] like other field management activities. Across five harvesting seasons, yield discrepancy revealed although

different from joined treatments to treatments (Figure 3A). From the joint use of pruning with stem number, biennial intensity could be reduced to 0.07 and 0.15 for T2 and T3 respectively (Figure 3B). The highest yield was recorded by T4 and T5 next to T6 respectively. However, the biennial

bearing ranged from 0.26 to 0.33 observed in these treatments. For efficiently diminishes of bienniality problem, applying the joint compatible coffee management practices with regular bearing coffee variety is crucial.




Figure 3. Combined Pruning methods with number of stems for biennial bearing improvement.

T1- Traditional pruning without removal of reproductive branches with 2 stems number per plant, T2- CPPAC with removal of 70% of reproductive branches that produced in the season with 2 stems number per plant, T3- CPPAC with removal of 50% of reproductive branches that produced in the season with 2 stems number per plant, T4- Traditional pruning without removal of reproductive branches with 3 stems number per plant, T5- CPPAC with removal of 70% of reproductive branches that produced in the season with 3 stems number per plant, T6- CPPAC with removal of 50% of reproductive branches that produced in the season with 3 stems number per plant. Source: Verdin Filho et al. [29]




Figure 4. Coffee productivity (60kg ha⁻¹) and biennial bearing intensity from 2004 to 2007.

Unsh: Unshaded, Sh: Shaded, I1-I3: Biennial intensity of two years, three years and four years; shade materials: Banana (*Musa sp.*) and coral bean plants (*Erythrina verna*), Source: Marta et al. [36]

2.4. Shade and Unshaded in Mitigation of Bienniality

Coffee production under shade has great importance for biodiversity conservation and buffering climate change [30-32]. In addition to this, shade increases the longevity

productivity of Arabica coffee and improve climate condition for coffee plant, improve soil condition and lower nutrient demand [33-35]. Hence, shade affects in multi-direction on mitigation of oscillation production. The significant of shade over unshaded in reducing bienniality is not revealed during

early seasons (I1) (Figure 4); but, the difference between management systems in reducing the level of bienniality observed in late harvesting seasons [36]. In agreement, Jaramillo-Botero et al. [1] confirmed that the effects of shade on yield fluctuation reduction is after the age of high yielding. Thus, importance of shade over unshaded in mitigation of biennial bearing was manifested from over three (I2) and four (I3) harvesting seasons (Figure 4). This may be due to shade gradual contributions in edaphic characteristics improvement [36-40] and its positive impacts in counterbalance the fruit bearing potential during on and off seasons [41].

The highest influence of yield bienniality revealed on unshaded coffee trees [1]. Shade enhances coffee vegetative growth aspect such as increase number of primary and secondary branches [9] and reduces branch die-back and drying of coffee branches [11, 42] which increase the potential productivity of coffee trees [36]. These phenomena have role in mitigation of biennial bearing by fostering new vegetative growth and generating flower bud of fruit on newly grown branches. Over shade decrease node number and coffee yield per area due to photosynthetically active radiation limitation [43-45]; but lessen yield biennial pattern relative to unshaded [1, 46] and has positive contribution in lengthen the age of productivity. Very low bienniality doesn't mean high yield [1]; but there is steady yield harvesting by producers when

compared with complete biennial bearing.

2.5. Fertilizer Application for Biennial Bearing Reduction

Applying fertilizer is another significant practices for sustainable coffee yield harvesting. Especially during on year/over bearing seasons high sink than source may be expected; this may expose coffee trees to different stress such as disease and dieback if not treated well. In such condition, following over bearing season, coffee tree's yield extremely decreased and it allows nutrient recovery and necessary growth causing biennial pattern [1]. Also, high load of berries causes for coffee trees exhaustion which may results complete drying of the trees. Thus, application of required level of fertilize protect such coffee tree from stress damage and exhaustion. Similarly, coffee tree requires high fertilizer after the beginning of high yield or high load berries stage [11]. Appropriate management of coffee trees using fertilizer reduces threat of highly necking and other physiological jeopardize, and it enhances yield bearing with less fluctuations. The use of 40% recommended fertilization for 48% shaded or photosynthetically active radiation blockage has better contributions in alternate bearing mitigation relative to 0% or unshaded coffee plots [1, 9].




Figure 5. Role of Integrated weed management on coffee yield.

T1-One slash in May + one slash in August, T2- One slash in May followed by one slash in July and one slash in August, T3- Herbicide in may followed one slash in July followed by mulch in September, T3- Herbicide in May soybean intercropping in June followed by mulch in September, T4- Herbicide in May followed by common bean intercropping in June followed by mulching in September, T5- Desmodium cover crop, T6- Vetch cover crop in June followed with one hand weeding, T7- Weed free. I1-I3 –represent biennial intensity of over two, three and four years respectively, Source: JARC [49]

2.6. Effects of Cover Crop in Alleviation of Biennial Bearing

Desmodium is one of the most important cover crop that has multi-function. It is highly important in coffee field/farm management practice for weed suppression, improve soil structure and increase soil fertility via fixing nitrogen. Also, it is one of the momentous practices for preventing infestation of coffee wilt disease (CWD) from coffee farm via avoiding the

agent such as slashing for weed control. Coffee is a perennial crop and grow slowly; the space between coffee trees and rows is wide and remain open for quit long seasons [47]; this encourages frequent growth of weed which seriously strive coffee for resource and leads decrease in productivity and production. Competitive weed can cause yield loss which reach up 65% to complete failure of coffee trees depending on types of weeds and coffee growth stage [48]. JARC [49] reported weed control efficiency of desmodium is 91.1%,

87.5% and 100% for second, third and fourth coffee harvesting seasons respectively. This may impose great contributions in coffee production increment and reduces high fluctuation of production due to reduced weed rivalry problem. Treatment 7 (Desmodium) and Treatment 4 (Herbicide+ common bean + mulching) gave the highest yield across seasons and in pooled over four seasons (Figure 5A). Both treatments had reduced biennial bearing intensity of the combined over four seasons (I3) to 0.21 and 0.27 respectively (Figure 5B). Despite their high yielding advantage and lessening bienniality to less than 0.3 (<0.3), still yield fluctuation observed when compared with the other low yielder treatments. Hence, one has to be conscious to use the desmodium with regular bearing cultivar during production for efficient biennial control. Scholars reported that the importance of integrating non-biennial cultivar with agronomic practices to reduce bienniality while increasing productivity [27, 50].

3. Conclusion and Recommendation

Agricultural practices play vital role in resilience abiotic and biotic factors which are hindrance for crop production. Arabica coffee is a perennial crop that needs intensive field managements. Agricultural practices buffer environmental stress and mitigate physiological jeopardize from coffee tree; these have great attribute in boosting coffee productivity and production. In addition to upsurge productivity, they alleviate yield fluctuation from year to year which is essential in realizing food security. Field managements viz shade, population density, fertilizer application and cover crop were discussed in lessening biennial bearing. These practices reduce bienniality by eliminating branch drying, branch die-back, improve soil structure, profuse flower initiation, over bearing and increase longevity of productivity of coffee tree.

The integrated agronomic practices mitigate biennial bearing than sole agronomic practice. Most practices that enhanced high yielding exhibited higher biennial bearing intensity which point out the importance of regular bearing cultivars involvement in mitigation of biennial intensity. For complete bienniality control and yield increment, compatible agronomic practices need to be integrated with non-biennial bearing variety. Thus, it is highly recommended to focus on developing regular yielding variety in the future *Coffea arabica* L. improvement program.

Competing Interests

The authors have declared that no competing interests exist among them.

References

- [1] Jaramillo-Botero C, Ricardo HSS, Herminia E P, Paulo RC, Merci PF (2010) Production and vegetative growth of coffee trees under fertilization and shade levels, Sci. Agric. (Piracicaba, Braz.) 67: 639-645.
- [2] Liliane T. N, Charles M S (2020) Factors Affecting Yield of Crops. In (Ed.), Agronomy - Climate Change & Food Security. Intech Open. <https://doi.org/10.5772/intechopen.90672>
- [3] Hatfield JL, Walthall CL (2014) Climate Change: Cropping System Changes and Adaptations. Encyclopedia of Agriculture and Food Systems 2: 256-264.
- [4] Guitton B, Kelner JJ, Velasco R, Gardiner SE, Chagné D, Costes E (2012) Genetic control of biennial bearing in apple. Journal of Experimental Botany 63: 131-149.
- [5] Jonkers H (1979) Biennial Bearing in Apple and Pear: A Literature Survey. Scientia Horticulturae 11, 303-317.
- [6] Nimisha S, Sanjay K S, Jai P, Manish S, Ajay K M, Nagendra K S (2017) Understand Bearing Habit in Mango. Curr Trends Biomedical Eng & Biosci 7: 30-32.
- [7] João C, Ana B (2018) Biennial bearing in apple cultivars. Rev. Ceres, Viçosa, 65: 144-149.
- [8] Monselise SP, Goldschmidt EE (1982) Alternate bearing in fruit trees. Horticultural Reviews 4: 128-173.
- [9] Jaramillo-Botero C, Santos RHS, Martinez HEP, Cecon PR, Fardin MP (2009) Production and vegetative development of coffee trees grown under solar radiation and fertilization levels, during years of high and low yield. American-Eurasian Journal of Agricultural and Environmental Science 6: 143-151.
- [10] Fichtner EJ, Chao YY, Ferguson L, Verreyne JS, Tang L, Lovatt CJ (2021) Repeating cycles of ON and OFF yields in alternate bearing olive, pistachio and citrus trees - different mechanisms, common solutions. Acta Horticulturae 1315: 1-10.
- [11] Da Matta FM (2004) Eco-physiological constraints on the production of shaded and unshaded coffee: a review. Field Crops Research 86: 99-114.
- [12] DaMatta FM, Rena AB, (2002) Ecofisiologia de cafezais sombreados e a pleno sol. In: Zambolim, L. (Ed.), O Estado da Arte de Tecnologias na Produção de Café. Universidade Federal de Viçosa, Viçosa, pp. 93-135.
- [13] Williams RR, Arnold GM, Flook VA, Jefferies CJ (1980) The effect of picking date on blossoming and fruit set in the following year for the apple cv. 'Bramley's seedling'. Journal of Horticultural Science 55: 359-362.
- [14] Romero-Alvarado Y, Soto-Pinto L, Garcia-Barrios L, BarreraGaytán F (2002) Coffee yields and soil nutrients under the shades of Inga sp. vs multiple species in Chiapas, México. Agroforestry Systems 54: 215-224.
- [15] Jaramillo-Botero, C., Martinez, H. E. P., & Santos, R. H. S. (2006). Characteristics of shade-grown coffee (*Coffea arabica* L.) in North of Latin America and Brazil: a comparative analysis. Coffee Science 1: 94-102.
- [16] Beaumont JH, Fukunaga ET (1958) Factors affecting the growth and yield of coffee in Kona, Hawaii, p40.
- [17] Montoya LA, Umanã R (1961) Effect of three light intensities and three levels of nitrogen (urea) on the die-back incidence. Café 3, 1-8.
- [18] Nacif, A. P. (1997). Phenology and productivity of coffee (*Coffea arabica* L.) cv. Catuaí under different planting densities and fertilizer doses, in the Cerrado of Patrocínio MG. D. Sc. Thesis. Federal University of Viçosa, Brazil, p. 124.

- [19] Taye k, Tesfaye S, Alemseged Y, Anteneh N, Endale T (2001) The impact of close spacing on yield of arabica coffee under contrasting agro-ecologies of Ethiopia. *African Crop Science Journal* 9: 401-410.
- [20] Pereira SP, Bartholo GF, Baliza DP, Sobreira FM, Guimarães RJ (2011) Growth, productivity and bienniality of coffee plants according to cultivation spacing. *Pesquisa Agropecuária Brasília*, 46: 152-160.
- [21] Silveira JMC, Nasser MD, Mariano-Nasser FAC, Pagliarini MK, Giomo GS (2018) Population density of Arabica coffee cultivars for bean quality and yield. *Pesq. Agropec. Trop., Goiânia* 48: 358-36.
- [22] Camargo MBP (2010) The Impact of Climatic Variability and Climate Change on Arabic coffee crop in Brazil: Review. *Bragantia, Campinas* 69: 239-247.
- [23] Pratibha LS, Goswami AK (2013) Effect of pruning and planting systems on growth, flowering, fruiting and yield of guava cv. Sardar. *Indian journal of horticulture* 70: 496-500.
- [24] Pandey SD, Amrendra K, Patel RK, Rai RR, Vishalnath (2015) Influence of planting densities on plant growth, yield and quality of litchi cv. Shahi. *Ecoscan. Journal of Environmental Sciences* 7: 397-401.
- [25] Gokavi N, Mukharib DS, Mote K, Manjunatha AN, RaghuramuluY (2019) Studies on planting geometry and pruning methods for improving productivity and reduce labour drudgery in Arabica coffee cv. Chandragiri. *Journal of Crop and Weed* 15: 58-64.
- [26] Njoroge JM et al. (1992) The influence of tree training and plant density on growth, yield components and yield of Arabica coffee cv. Ruiru 11. *Journal of Horticultural Science* 67: 695-702.
- [27] Masood, K., Richard, H., Gurreet, B., & Louise, F. (2020). Alternate bearing in pistachio (*Pistacia vera* L.): a review. *Trees*, <https://doi.org/10.1007/s00468-020-01967-y>
- [28] Luiz CP, Aledir CR (2004) Coffee Nutrition as a Function of Plant Density and NPK Fertilization. *Bragantia, Campinas* 63: 239-251.
- [29] Verdin Filho C, Paulo SV, Maria AG, Romário GF, Aldo LM, Aymbiré F, Almeida da Fonseca F, Alixsandro T, Saul de Andrade J (2016) New Management Technology for Arabica Coffee: The Cyclic Pruning Program for Arabica coffee. *Coffee Science, Lavras* 11: 475 – 483.
- [30] Lonera SP, Ivette P, Juan CH, Javier CN (2000) Shade effect on coffee production at the northern Tzeltal zone of the state of Chiapas, Mexico. *Agriculture, Ecosystems and Environment* 80: 61-69.
- [31] Perfecto I, Vandermeer JH, Bautista GL, Nunez GI, Greenberg R, Bichier P, Langridge S (2004) Greater predation in shaded coffee farms: The role of resident neotropical birds. *Ecology* 85: 2677-2681.
- [32] Perfecto I, Vandermeer J (2010) The agro ecological matrix as alternative to the land-sparing/agriculture intensification model. *Proceedings of the National Academy of Sciences* 107: 5786-5791.
- [33] Valdir A, Fernando F, Goulart, Tamiel K, Jacobson B, Reinaldo J, de Miranda F, Clarilton E D, Cardoso R (2016) Shade's Benefit: Coffee Production under Shade and Full Sun. *Journal of Agricultural Science*, 8: 11-19.
- [34] Vaast P, Harmand JM, Rapidel B, Jagoret P, Deheuvels O (2016) Coffee and Cocoa Production in Agroforestry—A Climate-Smart Agriculture Model. In: Torquebiau, E. (eds). *Climate Change and Agriculture Worldwide* 209-224.
- [35] Sarmiento-Soler A, Vaast P, Hoffmann MP, Rötter RP, Jassogne L, van Asten PJ, Graefe S (2019) Water use of *Coffea arabica* L. in open versus shaded systems under smallholder's farm conditions in Eastern Uganda. *Agricultural and Forest Meteorology* 266–267: 231–242.
- [36] Marta SF, Janaina RC, Nelson GO, Marinete BR (2011) Vegetative and productive aspects of organically grown coffee cultivars under shaded and unshaded systems. *Sci. Agric. (Piracicaba, Braz.)* 68: 424-430.
- [37] Damatta FM, Ramalho JD (2006) Impacts of drought and temperature stress on coffee physiology and production: A review. *Brazilian Journal of Plant Physiology* 18: 55-81.
- [38] Souza H N, de Goede RGM, Brussaard L, Cardoso IM, Duarte EM, Fernandes RB, Gomes LC, Pulleman MM (2012) Protective shade, tree diversity and soil properties in coffee agroforestry systems in the Atlantic Rainforest biome. *Agriculture, Ecosystems & Environment* 146: 179–196.
- [39] Meylan L, Gary C, Allinne C, Ortiz J, Jackson L, Rapidel B (2017) Evaluating the effect of shade trees on provision of ecosystem services in intensively managed coffee plantations. *Agric. Ecosyst. Environ* 245: 32–42.
- [40] Sauvadet M, den Meersche KV, Allinne C, Gay F, de Melo Virginio F E, Chauvat M, Becquer T, Tixier P, Harmand JM (2019) Shade trees have higher impact on soil nutrient availability and food web in organic than conventional coffee agroforestry. *Science of the Total Environment*, 649: 1065–1074.
- [41] Molla M A (2015) Effect of Tree Shade on Coffee Crop Production. *Journal of Sustainable Development* 8: 66-70.
- [42] Vaast P, van Kanten R, Siles P, Angrand J, Aguilar A (2008) Biophysical Interactions Between Timber Trees and Arabica Coffee in Suboptimal Conditions of Central America. In: Jose, S., Gordon, A. M. (eds) *Toward Agroforestry Design. Advances in Agroforestry*, Springer, 4: 133-146.
- [43] Campanha MM, Silva Santos RH, De Freitas GB, Prieto Martinez HF, Rebeiro Garcia SL, Finger FI (2004) Growing and yield of coffee plant in Agroforestry and monoculture systems in Minas, Gerais, Brazil. *Agroforestry systems* 63: 75-82.
- [44] Franck N, Vaast P, Dauzat J (2005) Coffee a Shade-adapted Plant: Implications on Its Carbon Balance and Consequences on Coffee Yield and Quality in Agroforestry Systems. In: 21st International Conference on Coffee Science, Montpellier (France), 11th - 15th September 2006.
- [45] Vaast P, Bertrand B, Perriot JJ, Guyot B, Génard M (2006) Fruit thinning and shade improve bean characteristics and beverage quality of coffee (*Coffea arabica* L.) under optimal conditions. *Journal of Agricultural Science* 86: 197–204.
- [46] DaMatta FM, Avila RT, Cardoso AA, Martins SC, Ramalho JC (2018) Physiological and agronomic performance of the coffee crop in the context of climate change and global warming: a review. *Journal of Agricultural Food Chemistry* 66: 5264–5274.

- [47] Tadesse E, Tesfu K (2015) Effect of Weed Management Methods on Yield and Physical Quality of Coffee at Gera, Jimma zone, South West Ethiopia. *Journal of Resources Development and Management* 11: 82-89.
- [48] Tadesse E (1998) Weed control in the western coffee growing areas of Ethiopia. pp 22-27. In: Beyene, S., & Abera, D. (eds), Workshop, 12-14 November 1996, Nekemt, Ethiopia.
- [49] Jimma Agricultural Research Center (JARC) (2020) Research Progress Report for the period 2019/20. Habtamu S., Yechalew S., Efrem A., Admikew G., & Mikru T. (eds.), EIAR/JARC, Jimma, pp 346.
- [50] Kallsen C E, Parftt DE, Holtz BA (2014) Early differences in the intensity of alternate bearing among selected pistachio genotypes. *Horticultural Science* 42: 1740–1743.