

# A nonexistence of solutions to a supercritical problem

Kamal Ould Bouh

Department of Mathematics, Taibah University, Almadinah Almunawwarah, KSA

**Email addresses:**

hbouh@taibahu.edu.sa, kamal\_bouh@yahoo.fr

**To cite this article:**

Kamal Ould Bouh. A Nonexistence of Solutions to a Supercritical Problem. *Pure and Applied Mathematics Journal*.

Vol. 2, No. 6, 2013, pp. 184-190. doi: 10.11648/j.pamj.20130206.13

**Abstract:** In this paper, we study the nonlinear elliptic problem involving nearly critical exponent  $(P_\epsilon) : -\Delta u = K u^{\frac{n+2}{n-2}+\epsilon}$  in  $\Omega$ ;  $u > 0$  in  $\Omega$  and  $u = 0$  on  $\partial\Omega$  where  $\Omega$  is a smooth bounded domain in  $\mathbb{R}^n$ ,  $n \geq 3$ ,  $K$  is a  $C^3$  positive function and  $\epsilon$  is a small positive real parameter. We prove that, for  $\epsilon$  small,  $(P_\epsilon)$  has no positive solutions which blow up at one critical point of the function  $K$ .

**Keywords:** Nonlinear Elliptic Equations, Critical Exponent, Variational Problem

## 1. Introduction

Let  $\Omega$  be a smooth bounded domain in  $\mathbb{R}^n$ ,  $n \geq 3$ . We consider the following nonlinear elliptic problem

$$(P_\epsilon) \begin{cases} -\Delta u = K u^{p+\epsilon} & \text{in } \Omega \\ u > 0 & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

where  $K$  is a  $C^3$  positive function,  $p + 1 = 2n/n - 2$  is the critical Sobolev exponent and  $\epsilon$  is a small positive real parameter.

Problem  $(P_\epsilon)$  is in some sense related to the limiting problem (when  $\epsilon = 0$ ) and the interest to it comes from its resemblance to the scalar curvature problem in differential geometry, which consists in finding suitable conditions on a given function  $K$  defined on  $M$  such that  $K$  is the scalar curvature for a metric  $\bar{g}$  conformally equivalent to  $g$ , where  $(M, g)$  is a  $n$ -dimensional Riemannian manifold without boundary.

Note that the limiting problem has been

widely studied in various works see for example [1], [2], [7] and [10].

In another view point, it is interesting to study the problem  $(P_\epsilon)$  with  $\epsilon < 0$  and  $\epsilon > 0$  and to understand what happens to the solutions of  $(P_\epsilon)$  (if they exist) as  $\epsilon \rightarrow 0$  !!

When  $\epsilon \in (1 - p, 0)$ , the mountain pass lemma proves the existence of solutions of  $(P_\epsilon)$  (see [3]). Note that, many works have been devoted to the study of positive solutions of  $(P_\epsilon)$  with  $\epsilon < 0$ . In sharp contrast to this, very little study has been made concerning the sign-changing solutions of  $(P_\epsilon)$  with  $\epsilon < 0$  and even less for  $\epsilon > 0$ .

When  $\epsilon > 0$ , problem  $(P_\epsilon)$  becomes more delicate since we lose the Sobolev embedding which is an important difficulty to overcome.

Concerning the supercritical case,  $\epsilon > 0$  and  $K$  is a constant, it was proved in [4] that  $(P_\epsilon)$  has no positive solution which blows up at a single point. This result shows that the situation is different from the subcritical one. However, del Pino et al [6] gave an existence result for two blow up points, provided that  $\Omega$  satisfies some geometrical conditions. In sharp contrast to this, it proved in [5] for the case  $K$  is a constant and [8] for the case  $K$  is a non constant function that, for  $\epsilon$  small,  $(P_\epsilon)$  has no sign-changing solutions which blow up at two points.

In this paper, we consider the case  $K$  is a non constant function and we look to understand the influence of the function  $K$  in the study of the positive solutions of  $(P_\epsilon)$  which blows up at a single point.

It is well known that problem  $(P_\epsilon)$  has a variational structure. Setting

$$J(u) = \frac{\int_\Omega |\nabla u|^2}{(\int_\Omega K |u|^{p+1+\epsilon})^{\frac{p+1+\epsilon}{2}}}, u \in H_0^1(\Omega), u \neq 0,$$

the positive critical points of  $J$  are solutions to  $(P_\epsilon)$ , up to a multiplicative constant.  $J$  satisfies the Palais-Smale condition in the subcritical case, whereas this condition fails in the critical case. Such a failure is due to the function

$$\delta_{(a,\lambda)}(x) = C_0 \frac{\lambda^{\frac{n-2}{2}}}{(1+\lambda^2|x-a|^2)^{\frac{n-2}{2}}}, C_0 = (n(n-2))^{\frac{n-2}{4}}, \lambda > 0, a \in \mathbb{R}^n \quad (1)$$

which are the only solutions of

$$-\Delta u = u^{\frac{n+2}{n-2}}, u > 0 \text{ in } \mathbb{R}^n, \text{ with } u \in L^{p+1}(\mathbb{R}^n) \text{ and } \nabla u \in L^2(\mathbb{R}^n) \text{ and are also the only minimizers of the}$$

Sobolev inequality on the whole space, that is

$$S = \inf_{L^{n-2}(\mathbb{R}^n)} |u|^{-\frac{2n}{n-2}}, s.t \nabla u \in L^2, u \in L^{\frac{2n}{n-2}}(\mathbb{R}^n), u \neq 0 \quad (2)$$

We have the following nonexistence result for  $(P_\epsilon)$ :

**Theorem 1**

Let  $\Omega$  be any smooth bounded domain in  $\mathbb{R}^n, n \geq 3$ . Assume that  $a_0 \in \Omega$  is a critical point of  $K$  satisfying one of the following conditions:

- (i)  $n = 3$ ,
- (ii)  $n = 4$ , and  $c_1 H(a_0, a_0) - \frac{c_3 \Delta K(a_0)}{16K(a_0)} > 0$ ,
- (iii)  $n \geq 5$ , and  $-\Delta K(a_0) > 0$ .

Then the problem  $(P_\epsilon)$  has no solution  $u_\epsilon$  such that  $u_\epsilon = \alpha_\epsilon P\delta_{a_\epsilon, \lambda_\epsilon} + v_\epsilon$  with  $|u_\epsilon|^\epsilon$  is bounded and  $v_\epsilon \rightarrow 0$  in  $H_0^1(\Omega)$   $\alpha_\epsilon \rightarrow K(a)^{(2-n)/4}, a_\epsilon \in \Omega, a_\epsilon \rightarrow a_0$  and  $\lambda_\epsilon d(a_\epsilon, \partial\Omega) \rightarrow +\infty$  as  $\epsilon \rightarrow 0$ .

**2. Preliminary Results**

We need to introduce some notations:

$P\delta_{a, \lambda}$  is defined as the only function in  $H_0^1(\Omega)$  such that  $\Delta P\delta_{a, \lambda} = \Delta\delta_{a, \lambda}$ . Writing

$$P\delta_{a, \lambda} = \delta_{a, \lambda} - \theta_{a, \lambda} \quad (3)$$

we have

$$\Delta\theta_{a, \lambda} = 0 \text{ in } \Omega; \theta_{a, \lambda} = \delta_{a, \lambda} \text{ on } \partial\Omega \quad (4)$$

We note that projections  $P\delta_{a, \lambda}$  of  $\delta_{a, \lambda}$ 's on  $H_0^1(\Omega)$  are approximate solutions to the limiting problem as  $a_\epsilon \in \Omega$  and  $\lambda_\epsilon d(a_\epsilon, \partial\Omega)$  goes to infinity.

Let  $G$  be the Green's function for the Laplace operator with Dirichlet boundary conditions, that is, for any  $x \in \Omega$ .

$$\begin{cases} -\Delta G(x, \cdot) = c_n \delta_x \text{ in } \Omega \\ G(x, \cdot) = 0 \text{ on } \partial\Omega \end{cases}$$

with  $\delta_x$  the Dirac mass at  $x$  and  $c_n = (n-2)|S^{n-1}|$

We denote by  $H$  the regular part of  $G$ , i.e.

$$H(x_1, x_2) = |x_1 - x_2|^{2-n} - G(x_1, x_2) \text{ for } (x_1, x_2) \in \Omega \times \Omega$$

The maximum principle provides us with the uniform estimate

$$\theta_{a, \lambda}(x) = C_0 \frac{H(x, a)}{\lambda^{\frac{n-2}{2}}} + O\left(\frac{1}{\lambda^{\frac{n+2}{2}(d(a, \partial\Omega))^n}}\right) \text{ as } \lambda d(a, \partial\Omega) \rightarrow +\infty \quad (5)$$

Corresponding estimates hold for the derivatives of  $\theta_{a, \lambda}$  with respect to  $a, \lambda$  and  $x$ .

Note that  $H(x, x) = O(d(x, \partial\Omega)^{2-n})$  as  $d(x, \partial\Omega) \rightarrow 0$  [9]. From [9] we also know that

$$\int_\Omega |\nabla\theta_{a, \lambda}|^2 = O(\lambda d(a, \partial\Omega)^{2-n}) \text{ as } \lambda d(a, \partial\Omega) \rightarrow +\infty \quad (6)$$

Next, we recall that for  $u_\epsilon$  satisfying the assumption of the theorem, there is a unique way to choose  $a_\epsilon, \lambda_\epsilon$  and  $v_\epsilon$  such that

$$u_\epsilon = \alpha_\epsilon P\delta_{a_\epsilon, \lambda_\epsilon} + v_\epsilon \quad (7)$$

with

$$\begin{cases} \alpha_\epsilon \in \mathbb{R}, \alpha_\epsilon \rightarrow K(\alpha_\epsilon)^{(2-n)/4} \\ \alpha_\epsilon \in \Omega, \lambda_\epsilon \in \mathbb{R}_+^*, \lambda_\epsilon d(a_\epsilon, \partial\Omega) \rightarrow +\infty \\ v_\epsilon \rightarrow 0 \text{ in } H_0^1(\Omega), v_\epsilon \in E_{a_\epsilon, \lambda_\epsilon} \end{cases} \quad (8)$$

and for any  $(a, \lambda) \in \Omega \times \mathbb{R}_+^*, E_{(a, \lambda)}$  denotes the subspace of  $H_0^1(\Omega)$  defined by

$$E_{(a, \lambda)} = \left\{ w \in H_0^1(\Omega) / (w, P\delta_{(a, \lambda)})_{H_0^1} = \left( w, \frac{\partial P\delta_{(a, \lambda)}}{\partial \lambda} \right)_{H_0^1} = \left( w, \frac{\partial P\delta_{(a, \lambda)}}{\partial a_i} \right)_{H_0^1} = 0, 1 \leq i \leq n \right\}$$

For the proof of this fact, see [1], [9]. In the following, we always assume that  $u_\epsilon$ , satisfying the assumption of the theorem, is written as in (8). In order to simplify the notations, we set

$$\delta_{a_\epsilon, \lambda_\epsilon} = \delta_\epsilon, P\delta_{a_\epsilon, \lambda_\epsilon} = P\delta_\epsilon \text{ and } \theta_{a_\epsilon, \lambda_\epsilon} = \theta_\epsilon$$

**Lemma 2**

Let  $u_\epsilon$  satisfying the assumption of the theorem 1. Then

$$(i) \int_\Omega |\nabla u_\epsilon|^2 \rightarrow S^{n/2}; \quad (ii) \int_\Omega K u_\epsilon^{p+1+\epsilon} \rightarrow S^{n/2}$$

as  $\epsilon \rightarrow 0$ ,  $S, S$  denoting the Sobolev constant defined by (2).

**Proof.**

We have

$$\int_\Omega |\nabla u_\epsilon|^2 = \int_\Omega |\nabla(\alpha_\epsilon P\delta_\epsilon + v_\epsilon)|^2 = \alpha_\epsilon^2 \int_\Omega |\nabla P\delta_\epsilon|^2 + \int_\Omega |\nabla v_\epsilon|^2 \text{ since } v_\epsilon \in E_{a_\epsilon, \lambda_\epsilon}$$

From the fact that  $\delta_\epsilon$  satisfies  $-\Delta\delta_\epsilon = \delta_\epsilon^p$  in  $\mathbb{R}^n$  and is a minimizer for  $S$ , we deduce that  $\int_{\mathbb{R}^n} |\nabla \delta_\epsilon|^2 = S^{n/2}$

On the other hand, an explicit computation provides us with

$$\int_\Omega |\nabla\delta_{a, \lambda}|^2 = \int_{\mathbb{R}^n} |\nabla\delta_{a, \lambda}|^2 + O\left(\frac{1}{(\lambda d(a, \partial\Omega))^n}\right) \text{ as } \lambda d(a, \partial\Omega) \rightarrow +\infty.$$

Taking account of (6), claim (i) is a consequence of (8). Claim (ii) follows from the fact that  $u_\epsilon$  solves  $(P_\epsilon)$ .

**3. Estimating  $v_\epsilon$**

As usual in this type of problems, we first deal with the  $v$ -part of  $u$ , in order to show that it is negligible with respect to the concentration phenomenon.

**Lemma 3**

Let  $u_\epsilon$  satisfying the assumption of the theorem.  $\lambda_\epsilon$  occurring in (7) satisfies

$$\lambda_\epsilon^\epsilon \rightarrow 1, \text{ as } \epsilon \rightarrow 0 .$$

Proof.

According to Lemma 2, we have

$$\int_\Omega K u_\epsilon^{p+1+\epsilon} = S^{n/2} + o(1) \text{ as } \epsilon \rightarrow 0 \quad (9)$$

and

$$\begin{aligned} \int_\Omega K u_\epsilon^{p+1+\epsilon} &= \int_\Omega K (\alpha_\epsilon P \delta_\epsilon + v_\epsilon)^{p+\epsilon} \alpha_\epsilon P \delta_\epsilon \\ &+ \int_\Omega K u_\epsilon^{p+\epsilon} v_\epsilon \\ &= \alpha_\epsilon^{p+1+\epsilon} \int_\Omega K P \delta_\epsilon^{p+\epsilon+1} \\ &- \int_\Omega \Delta u_\epsilon v_\epsilon \\ &+ O\left(\int_\Omega P \delta_\epsilon^{p+\epsilon} |v_\epsilon| + \int_\Omega |v_\epsilon|^{p+\epsilon} P \delta_\epsilon\right) \\ &= \alpha_\epsilon^{p+1+\epsilon} \int_\Omega K P \delta_\epsilon^{p+\epsilon+1} \\ &+ O\left(\lambda_\epsilon^{\frac{\epsilon(n-2)}{2}} \int_\Omega P \delta_\epsilon^p |v_\epsilon| \right. \\ &\left. + \lambda_\epsilon^{\frac{\epsilon(n-2)}{2}} \int_\Omega |v_\epsilon|^{p+\epsilon} P \delta_\epsilon^{1-\epsilon} + |v_\epsilon|_{H_0^1}\right) \\ &= \alpha_\epsilon^{p+1+\epsilon} \int_\Omega K P \delta_\epsilon^{p+\epsilon+1} \\ &+ O\left(\lambda_\epsilon^{\epsilon(n-2)/2} |v_\epsilon|_{L^{p+1}} \right. \\ &\left. + \lambda_\epsilon^{\epsilon(n-2)/2} |v_\epsilon|_{L^{p+1}}^{p+\epsilon} + |v_\epsilon|_{H_0^1}\right) \end{aligned}$$

Thus

$$\int_\Omega K u_\epsilon^{p+1+\epsilon} = \alpha_\epsilon^{p+1+\epsilon} \int_\Omega K P \delta_\epsilon^{p+\epsilon+1} + o\left(\lambda_\epsilon^{\epsilon(n-2)/2} + 1\right) \quad (10)$$

We observe that

$$\begin{aligned} \int_\Omega K P \delta_\epsilon^{p+1+\epsilon} &= \int_\Omega K (\delta_\epsilon - \theta_\epsilon)^{p+\epsilon+1} = \int_\Omega K \delta_\epsilon^{p+1+\epsilon} + \\ &O\left(\int_\Omega \delta_\epsilon^{p+\epsilon} \theta_\epsilon\right) \\ &= C_0^{p+\epsilon+1} K(a_\epsilon) \int_B \left(\frac{\lambda_\epsilon}{1+\lambda_\epsilon^2|x-a_\epsilon|^2}\right)^{\frac{(p+1+\epsilon)(n-2)}{2}} + \\ &O\left(|\theta_\epsilon|_{L^\infty} \int_\Omega \left(\frac{\lambda_\epsilon}{1+\lambda_\epsilon^2|x-a_\epsilon|^2}\right)^{\frac{(p+1+\epsilon)(n-2)}{2}} + \frac{\lambda_\epsilon^{\frac{\epsilon(n-2)}{2}}}{(\lambda_\epsilon d_\epsilon)^n}\right) \end{aligned}$$

where  $B = B(a_\epsilon, d_\epsilon)$ . Using Proposition 1 of [9], we obtain

$$\begin{aligned} &\int_\Omega K P \delta_\epsilon^{p+1+\epsilon} \\ &= \lambda_\epsilon^{\frac{\epsilon(n-2)}{2}} K(a_\epsilon) \left( C_0^{p+\epsilon+1} \int_{\mathbb{R}^n} \frac{dx}{(1+|x|^2)^{\frac{(p+1+\epsilon)(n-2)}{2}}} \right. \\ &- C_0^{p+\epsilon+1} \int_{\mathbb{R}^n/B} \frac{dx}{(1+|x|^2)^{\frac{(p+1+\epsilon)(n-2)}{2}}} \left. \right) \\ &+ O\left(\frac{\lambda_\epsilon^{\frac{\epsilon(n-2)}{2}}}{(\lambda_\epsilon d_\epsilon)^{n-2}}\right) \\ &= \lambda_\epsilon^{\epsilon(n-2)/2} K(a_\epsilon) C_0^{p+\epsilon+1} \int_{\mathbb{R}^n} \frac{dx}{(1+|x|^2)^{(p+1+\epsilon)(n-2)/2}} \\ &+ O\left(\frac{\lambda_\epsilon^{\frac{\epsilon(n-2)}{2}}}{(\lambda_\epsilon d_\epsilon)^{n-2}}\right) \end{aligned}$$

We note that

$$\begin{aligned} C_0^{p+\epsilon+1} \int_{\mathbb{R}^n} \frac{dx}{(1+|x|^2)^{(p+1+\epsilon)(n-2)/2}} &= C_0^{p+1} \int_{\mathbb{R}^n} \frac{dx}{(1+|x|^2)^n} + \\ &O(\epsilon) = S^{n/2} + O(\epsilon). \end{aligned}$$

Therefore

$$\int_\Omega K P \delta_\epsilon^{p+1+\epsilon} = \lambda_\epsilon^{\epsilon(n-2)/2} K(a_\epsilon) (S^{n/2} + O(\epsilon) + o(1)) \quad (11)$$

so (10) and (11) provide us with

$$\int_\Omega K u_\epsilon^{p+1+\epsilon} = \alpha_\epsilon^{p+1+\epsilon} \lambda_\epsilon^{\epsilon(n-2)/2} K(a_\epsilon) (S^{n/2} + o(1)) + o(1) \quad (12)$$

Combination of (9) and (12) proves the lemma.

Next, we recall the following estimate [10] :

**Remark 4**

$\delta_\epsilon^\epsilon(x) - C_0^\epsilon \lambda_\epsilon^{\epsilon(n-2)/2} = O(\epsilon \log(1 + \lambda_\epsilon^2|x - a_\epsilon|^2))$  in  $\Omega$ . We are now able to study the  $v_\epsilon$  -part of  $u_\epsilon$ .

**Lemma 5**

Let  $u_\epsilon$  satisfying the assumption of the theorem.  $v_\epsilon$  occurring in (7) satisfies

$$\begin{aligned} &|v_\epsilon|_{H_0^1(\Omega)} \\ &\leq C + C \begin{cases} \frac{\nabla K(a_\epsilon)}{\lambda_\epsilon} + \frac{1}{(\lambda_\epsilon d_\epsilon)^{n-2}} & \text{if } n < 6 \\ \frac{\nabla K(a_\epsilon)}{\lambda_\epsilon} + \frac{\log(\lambda_\epsilon d_\epsilon)}{(\lambda_\epsilon d_\epsilon)^4} & \text{if } n = 6 \\ \frac{\nabla K(a_\epsilon)}{\lambda_\epsilon} + \frac{1}{(\lambda_\epsilon d_\epsilon)^{(n+2)/2}} & \text{if } n > 6 \end{cases} \end{aligned}$$

with  $C$  independent of  $\epsilon$ .

$$Q_0(v, v) \geq c|v|_{H_0^1}^2 \quad \forall v \in E_{(a_\epsilon, \lambda_\epsilon)}. \quad (14)$$

**Proof.**

Multiplying  $(P_\epsilon)$  by  $v_\epsilon$  and integrating on  $\Omega$ , we obtain

$$0 = \int_{\Omega} \nabla u_\epsilon \cdot \nabla v_\epsilon - \int_{\Omega} K u_\epsilon^{p+\epsilon} v_\epsilon$$

Thus

$$0 = \int_{\Omega} |\nabla v_\epsilon|^2 - \int_{\Omega} K [(\alpha_\epsilon P \delta_\epsilon)^{p+\epsilon} + (p + \epsilon)(\alpha_\epsilon P \delta_\epsilon)^{p-1+\epsilon} v_\epsilon + O(\delta_\epsilon^{p-2+\epsilon} v_\epsilon^2 \chi_{|v_\epsilon| < \delta_\epsilon} + |v_\epsilon|^{p+\epsilon})] v_\epsilon.$$

Using the assumption that  $|u_\epsilon|^\epsilon$  is bounded, we find

$$0 = Q_\epsilon(v_\epsilon, v_\epsilon) - f_\epsilon(v_\epsilon) + o(|v_\epsilon|_{H_0^1}^{\min(3, p+1)} + |v_\epsilon|_{H_0^1}^{p+1}) \quad (13)$$

with

$$Q_\epsilon(v, v) = |v_\epsilon|_{H_0^1}^2 - (p + \epsilon) \int_{\Omega} K (\alpha_\epsilon P \delta_\epsilon)^{p-1+\epsilon} v^2$$

and

$$f_\epsilon(v) = \int_{\Omega} K (\alpha_\epsilon P \delta_\epsilon)^{p+\epsilon} v.$$

We observe that

$$\begin{aligned} Q_\epsilon(v, v) &= |v_\epsilon|_{H_0^1}^2 - p \int_{\Omega} K (\alpha_\epsilon P \delta_\epsilon)^{p-1+\epsilon} v^2 \\ &\quad + o(\epsilon |v_\epsilon|_{H_0^1}^2) \\ &= |v_\epsilon|_{H_0^1}^2 - p \alpha_\epsilon^{p-1+\epsilon} K(a_\epsilon) \int_{\Omega} (\delta_\epsilon^{p-1+\epsilon} \\ &\quad + O(\delta_\epsilon^{p-1+\epsilon} \theta_\epsilon)) v^2 + o(|v|_{H_0^1}^2) \\ &= |v_\epsilon|_{H_0^1}^2 - p \alpha_\epsilon^{p-1+\epsilon} K(a_\epsilon) C_0^\epsilon \lambda_\epsilon^{\epsilon(n-2)/2} \int_{\Omega} \delta_\epsilon^{p-1} v^2 \\ &\quad + o\left(\int_{\Omega} (\delta_\epsilon^{p-1+\epsilon} - C_0^\epsilon \lambda_\epsilon^{\epsilon(n-2)/2} \delta_\epsilon^{p-1} |v|^2)\right) + o(|v|_{H_0^1}^2) \end{aligned}$$

Using Remark 4, we find

$$Q_\epsilon(v, v) = Q_0(v, v) + o(|v|_{H_0^1}^2) \quad \text{with}$$

$$Q_0(v, v) = |v|_{H_0^1}^2 - \int_{\Omega} \delta_\epsilon^{p-1} v^2.$$

According to [1],  $Q_0$  is coercive, that is, there exists some constant  $c > 0$  independent of  $\epsilon$ , for  $\square$  small enough, such that

We also observe that

$$\begin{aligned} f_\epsilon(v) &= \alpha_\epsilon^{p+\epsilon} \int_{\Omega} K (\delta_\epsilon^{p+\epsilon} + O(\delta_\epsilon^{p-1+\epsilon} \theta_\epsilon)) v \\ &= \alpha_\epsilon^{p+\epsilon} \left[ C_0^\epsilon \lambda_\epsilon^{\epsilon(n-2)/2} \int_{\Omega} K \delta_\epsilon^p v \right. \\ &\quad \left. + O\left(\epsilon \int_{\Omega} K \log(1 + \lambda_\epsilon^2 |x - a_\epsilon|^2) \delta_\epsilon^2 |v| \right. \right. \\ &\quad \left. \left. + \int_{\Omega} \delta_\epsilon^{p-1} \theta_\epsilon |v| \right) \right] \end{aligned}$$

The last equality follows from Remark 4. Therefore we can write,

$$\begin{aligned} &\text{with } B = B(a_\epsilon, d_\epsilon) \\ f_\epsilon(v) &= o\left(\epsilon |v|_{H_0^1} + \int_B \delta_\epsilon^{p-1} \theta_\epsilon |v_\epsilon| + \int_{\mathbb{R}^n \setminus B} \delta_\epsilon^p |v|\right) \\ f_\epsilon(v) &= o\left(\left(\epsilon + \frac{|\nabla K(a_\epsilon)|}{\lambda_\epsilon^2}\right) |v|_{H_0^1} \right. \\ &\quad \left. + |v|_{H_0^1} |\theta_\epsilon|_{L^\infty} \left(\int_B \delta_\epsilon^{\frac{8n}{n^2-4}}\right)^{\frac{n+2}{2n}} \right. \\ &\quad \left. + |v_\epsilon|_{H_0^1} \left(\int_{\mathbb{R}^n \setminus B} \delta_\epsilon^{\frac{2n}{n^2-2}}\right)^{\frac{n+2}{2n}}\right) \end{aligned}$$

We notice that

$$\int_{\mathbb{R}^n \setminus B} \delta_\epsilon^{2n/(n-2)} = o\left(\frac{1}{(\lambda_\epsilon d_\epsilon)^n}\right)$$

and

$$\left(\int_B \delta_\epsilon^{\frac{8n}{n^2-4}}\right)^{(n+2)/2n} \leq C \begin{cases} \frac{d_\epsilon^{(n-6)/2}}{\lambda_\epsilon^2} & \text{if } n > 6 \\ \frac{\log(\lambda_\epsilon d_\epsilon)}{\lambda_\epsilon^2} & \text{if } n = 6 \\ \frac{1}{\lambda_\epsilon^{(n-2)/2}} & \text{if } n < 6 \end{cases}$$

Using (5), we obtain

$$\begin{aligned} |f_\epsilon(v)| &\leq C |v|_{H_0^1} \\ &+ C \begin{cases} \left(\frac{|\nabla K(a_\epsilon)|}{\lambda_\epsilon} + \frac{1}{(\lambda_\epsilon d_\epsilon)^{n-2}}\right) |v|_{H_0^1} & \text{if } n < 6 \\ \left(\frac{|\nabla K(a_\epsilon)|}{\lambda_\epsilon} + \frac{\log(\lambda_\epsilon d_\epsilon)}{(\lambda_\epsilon d_\epsilon)^4}\right) |v|_{H_0^1} & \text{if } n = 6 \\ \left(\frac{|\nabla K(a_\epsilon)|}{\lambda_\epsilon} + \frac{1}{(\lambda_\epsilon d_\epsilon)^{(n+2)/2}}\right) |v|_{H_0^1} & \text{if } n > 6 \end{cases} \quad (15) \end{aligned}$$

Combining (13), (14) and (15), we obtain the desired estimate.

### 4. Proof of Theorem

Let us start by proving the following crucial result :

**Proposition 6**

Let  $u_\epsilon$  satisfying the assumption of the theorem. Then,

$$\left| \alpha_\epsilon c_1 \frac{H(a_\epsilon, a_\epsilon)}{\lambda_\epsilon^{n-2}} - \alpha_\epsilon \frac{c_3}{n^2} \frac{\Delta K(a_\epsilon)}{K(a_\epsilon) \lambda_\epsilon^2} + \alpha_\epsilon c_2 \varepsilon \right| \leq c \left( \epsilon^2 + \frac{1}{\lambda_\epsilon^3} + |v|_{H_0^1}^2 \right) + \begin{cases} \frac{\log(\lambda_\epsilon d_\epsilon)}{(\lambda_\epsilon d_\epsilon)^n} & (n \geq 4) \\ \frac{1}{(\lambda_\epsilon d_\epsilon)^2} & (n = 3) \end{cases} \quad (16)$$

where  $a_\epsilon, \lambda_\epsilon$  and  $d_\epsilon = (a_\epsilon, \partial\Omega)$  are given in (7) and  $c_1, c_2, c_3$  are positive constants defined by

$$c_1 = c_0^{\frac{2n}{n-2}} \int_{\mathbb{R}^n} \frac{dx}{(1+|x|^2)^{(n+2)/2}},$$

$$c_2 = \frac{n-2}{2} c_0^{\frac{2n}{n-2}} \int_{\mathbb{R}^n} \log(1 + |x|^2) \frac{|x|^2 - 1}{(1+|x|^2)^{n+1}} dx$$

and 
$$c_3 = c_0^{\frac{2n}{n-2}} \int_{\mathbb{R}^n} \frac{|x|^2}{(1+|x|^2)^n} dx.$$

**Proof.**

Multiplying  $(P_\epsilon)$  by  $\lambda_\epsilon \frac{\partial P \delta_\epsilon}{\partial \lambda}$  and integrating on  $\Omega$ ,

we obtain

$$\begin{aligned} 0 &= - \int_{\Omega} \Delta u_\epsilon \lambda_\epsilon \frac{\partial P \delta_\epsilon}{\partial \lambda} - \int_{\Omega} K u_\epsilon^{p+\epsilon} \lambda_\epsilon \frac{\partial P \delta_\epsilon}{\partial \lambda} \\ &= \int_{\Omega} \nabla(\alpha_\epsilon P \delta_\epsilon + v_\epsilon) \nabla \left( \lambda_\epsilon \frac{\partial P \delta_\epsilon}{\partial \lambda} \right) - \int_{\Omega} K(\alpha_\epsilon P \delta_\epsilon \\ &\quad + v_\epsilon)^{p+\epsilon} \lambda_\epsilon \frac{\partial P \delta_\epsilon}{\partial \lambda} \\ &= \alpha_\epsilon \int_{\Omega} \delta_\epsilon^p \lambda_\epsilon \frac{\partial P \delta_\epsilon}{\partial \lambda} - \int_{\Omega} K [(\alpha_\epsilon P \delta_\epsilon)^{p+\epsilon} \\ &\quad + (p + \epsilon)(\alpha_\epsilon P \delta_\epsilon)^{p-1+\epsilon} v \\ &\quad + O(\delta_\epsilon^{p-2+\epsilon} |v_\epsilon|^2 \\ &\quad + |v_\epsilon|^{p+\epsilon})] \lambda_\epsilon \frac{\partial P \delta_\epsilon}{\partial \lambda}. \end{aligned} \quad (17)$$

We estimate each term of the right hand side in (17). First, we have

$$\int_{\Omega} \delta_\epsilon^p \lambda_\epsilon \frac{\partial P \delta_\epsilon}{\partial \lambda} = \int_{\Omega} \delta_\epsilon^p \lambda_\epsilon \frac{\partial \delta_\epsilon}{\partial \lambda} - \int_{\Omega} \delta_\epsilon^p \lambda_\epsilon \frac{\partial \theta_\epsilon}{\partial \lambda}$$

whence

$$\begin{aligned} &\int_{\Omega} \delta_\epsilon^p \lambda_\epsilon \frac{\partial P \delta_\epsilon}{\partial \lambda} \\ &= \int_{\mathbb{R}^n} \delta_\epsilon^p \lambda_\epsilon \frac{\partial \delta_\epsilon}{\partial \lambda} - \int_{\mathbb{R}^n \setminus \Omega} \delta_\epsilon^p \lambda_\epsilon \frac{\partial \delta_\epsilon}{\partial \lambda} - \int_B \delta_\epsilon^p \lambda_\epsilon \frac{\partial \theta_\epsilon}{\partial \lambda} \\ &\quad - \int_{\Omega \setminus B} \delta_\epsilon^p \lambda_\epsilon \frac{\partial \theta_\epsilon}{\partial \lambda} \\ &= O\left(\frac{1}{(\lambda_\epsilon d_\epsilon)^n}\right) - \lambda_\epsilon \frac{\partial \theta_\epsilon}{\partial \lambda}(a_\epsilon) \int_B \delta_\epsilon^p \\ &\quad + O\left(\lambda_\epsilon \int_B \delta_\epsilon^p |x \right. \\ &\quad \left. - a_\epsilon|^2 \sup_B \left| D_x^2 \frac{\partial \theta_\epsilon}{\partial \lambda} \right| \right) \end{aligned}$$

with  $B = (a_\square, d_\square)$ . According to [9], we have

$$\begin{aligned} \lambda_\epsilon \frac{\partial \theta_\epsilon}{\partial \lambda}(a_\epsilon) &= -\frac{n-2}{2} \frac{c_0}{\lambda_\epsilon^{(n-2)/2}} H(a_\epsilon, a_\epsilon) \\ &\quad + O\left(\frac{1}{\lambda_\epsilon^{(n+2)/2} d_\epsilon^n}\right) \end{aligned}$$

and

$$\sup_B \left| D_x^2 \frac{\partial \theta_\epsilon}{\partial \lambda} \right| = O\left(\frac{1}{\lambda_\epsilon^{n/2} d_\epsilon^n}\right)$$

Therefore, estimating the integrals we obtain

$$\begin{aligned} \int_{\Omega} \delta_\epsilon^p \lambda_\epsilon \frac{\partial P \delta_\epsilon}{\partial \lambda} &= \frac{n-2}{2} c_1 \frac{H(a_\epsilon, a_\epsilon)}{\lambda_\epsilon^{n-2}} \\ &\quad + O\left(\frac{\log(\lambda_\epsilon d_\epsilon)}{(\lambda_\epsilon d_\epsilon)^n}\right) \end{aligned} \quad (18)$$

Secondly, we compute

$$\begin{aligned} &\int_{\Omega} K(P \delta_\epsilon)^{p+\epsilon} \lambda_\epsilon \frac{\partial P \delta_\epsilon}{\partial \lambda} \\ &= \int_{\Omega} K [\delta_\epsilon^{p+\epsilon} - (p + \epsilon) \delta_\epsilon^{p-1+\epsilon} \theta_\epsilon \\ &\quad + O(\theta_\epsilon^2 \delta_\epsilon^{p-2+\epsilon} + \theta_\epsilon^{p+\epsilon})] \lambda_\epsilon \frac{\partial P \delta_\epsilon}{\partial \lambda} \\ &= \int_B K \delta_\epsilon^{p+\epsilon} \lambda_\epsilon \frac{\partial \delta_\epsilon}{\partial \lambda} \\ &\quad - \int_B K \delta_\epsilon^{p+\epsilon} \lambda_\epsilon \frac{\partial \theta_\epsilon}{\partial \lambda} - (p \\ &\quad + \epsilon) \int_B K \delta_\epsilon^{p-1+\epsilon} \theta_\epsilon \lambda_\epsilon \frac{\partial \delta_\epsilon}{\partial \lambda} \end{aligned}$$

$$\begin{aligned} &+ O\left( \int_{\Omega} \theta_\epsilon^2 \delta_\epsilon^{p-1+\epsilon} + \int_{\Omega} \delta_\epsilon^{p-1+\epsilon} \theta_\epsilon \left| \lambda_\epsilon \frac{\partial \theta_\epsilon}{\partial \lambda} \right| + \int_{\Omega} \theta_\epsilon^{p+\epsilon} \delta_\epsilon + \right. \\ &\quad \left. \frac{\lambda_\epsilon^{(n-2)}}{(\lambda_\epsilon d_\epsilon)^n} \right) \end{aligned} \quad (19)$$

and we have to estimate each term of the right hand side of (18). Using the fact that  $\lambda_\epsilon \frac{\partial \delta_\epsilon}{\partial \lambda} = \frac{n-2}{2} \frac{(1-\lambda_\epsilon^2|x-a_\epsilon|^2)}{(1+\lambda_\epsilon^2|x-a_\epsilon|^2)} \delta_\epsilon$ ,

we derive that

$$\begin{aligned} & \int_B K \delta_\epsilon^{p+\epsilon} \lambda_\epsilon \frac{\partial \delta_\epsilon}{\partial \lambda} \\ &= \frac{n-2}{2} \lambda_\epsilon^{\frac{\epsilon(n-2)}{n}} K(a_\epsilon) c_0^{p+1+\epsilon} \int_{\mathbb{R}^n} \frac{1}{(1+|x|^2)^{n+\frac{\epsilon(n-2)}{n}}} \frac{1-|x|^2}{1+|x|^2} dx \\ &+ O\left(\frac{\lambda_\epsilon^{\frac{\epsilon(n-2)}{n}}}{(\lambda_\epsilon d_\epsilon)^n}\right) \\ &= \lambda_\epsilon^{\frac{\epsilon(n-2)}{n}} \left( c_2 K(a_\epsilon) \epsilon - \alpha_\epsilon \frac{c_3}{n^2} \frac{\Delta K(a_\epsilon)}{\lambda_\epsilon^2} + O\left(\epsilon^2 + \frac{1}{\lambda_\epsilon^3}\right) \right) + \\ & \quad O\left(\frac{\lambda_\epsilon^{\frac{\epsilon(n-2)}{n}}}{(\lambda_\epsilon d_\epsilon)^n}\right) \end{aligned} \tag{20}$$

For the other terms in (19), we write

$$\begin{aligned} & \int_B K \delta_\epsilon^{p+\epsilon} \lambda_\epsilon \frac{\partial \theta_\epsilon}{\partial \lambda} = K(a_\epsilon) \lambda_\epsilon \frac{\partial \theta_\epsilon}{\partial \lambda}(a_\epsilon) \int_B \delta_\epsilon^{p+\epsilon} \\ & \quad + O\left(\int_B \delta_\epsilon^{p+\epsilon} \frac{|x-a_\epsilon|^2}{\lambda_\epsilon^{(n-2)/2} d_\epsilon^n}\right) \\ &= \frac{n-2}{2} K(a_\epsilon) c_0^{p+1+\epsilon} \frac{H(a_\epsilon, a_\epsilon)}{\lambda_\epsilon^{(n-2)/2}} \int_B \left(\frac{\lambda_\epsilon}{1+\lambda_\epsilon^2|x-a_\epsilon|^2}\right)^{(p+\epsilon)(n-2)/2} \\ & \quad + O\left(\frac{\lambda_\epsilon^{\epsilon(n-2)/2} \log(\lambda_\epsilon d_\epsilon)}{(\lambda_\epsilon d_\epsilon)^n}\right) \\ &= \frac{n-2}{2} c_1 K(a_\epsilon) \frac{H(a_\epsilon, a_\epsilon)}{\lambda_\epsilon^{n-2}} \lambda_\epsilon^{\epsilon(n-2)/2} \\ & \quad + O\left(\frac{\lambda_\epsilon^{\epsilon(n-2)/2} \log(\lambda_\epsilon d_\epsilon)}{(\lambda_\epsilon d_\epsilon)^n}\right). \end{aligned} \tag{21}$$

and

$$\begin{aligned} & \int_B K \delta_\epsilon^{p-1+\epsilon} \theta_\epsilon \lambda_\epsilon \frac{\partial \delta_\epsilon}{\partial \lambda} \\ &= \theta_\epsilon(a_\epsilon) K(a_\epsilon) \int_B \delta_\epsilon^{p-1+\epsilon} \lambda_\epsilon \frac{\partial \delta_\epsilon}{\partial \lambda} \\ & \quad + O\left(\int_B \delta_\epsilon^{p+\epsilon} \frac{|x-a_\epsilon|^2}{\lambda_\epsilon^{(n-2)/2} d_\epsilon^n}\right) \end{aligned}$$

Using (5), we obtain

$$\begin{aligned} \int_B K \delta_\epsilon^{p-1+\epsilon} \theta_\epsilon \lambda_\epsilon \frac{\partial \delta_\epsilon}{\partial \lambda} &= \frac{n-2}{2} K(a_\epsilon) c_1 \frac{H(a_\epsilon, a_\epsilon)}{\lambda_\epsilon^{n-2}} \lambda_\epsilon^{\epsilon(n-2)/2} + \\ & \quad O\left(\frac{\lambda_\epsilon^{\epsilon(n-2)/2} \log(\lambda_\epsilon d_\epsilon)}{(\lambda_\epsilon d_\epsilon)^n}\right). \end{aligned} \tag{22}$$

(19), (20), (21) and additional integral estimates of the same type provide us with the expansion

$$\begin{aligned} & \int_\Omega K(P\delta_\epsilon)^{p+\epsilon} \lambda_\epsilon \frac{\partial P\delta_\epsilon}{\partial \lambda} \\ &= \frac{n-2}{2} \lambda_\epsilon^{\epsilon(n-2)/2} \left[ c_2 K(a_\epsilon) \epsilon \right. \\ & \quad \left. + 2c_1 K(a_\epsilon) \frac{H(a_\epsilon, a_\epsilon)}{\lambda_\epsilon^{n-2}} - c_3 \frac{\Delta K(a_\epsilon)}{\lambda_\epsilon^2} \right] \\ & \quad + O\left(c_3 + \frac{1}{\lambda_\epsilon^3} + \frac{\log(\lambda_\epsilon d_\epsilon)}{(\lambda_\epsilon d_\epsilon)^n}\right) \\ & \quad + \frac{1}{(\lambda_\epsilon d_\epsilon)^2} \quad (\text{if } n=3). \end{aligned} \tag{23}$$

We note that

$$\begin{aligned} & \int_\Omega K(P\delta_\epsilon)^{p-1+\epsilon} v_\epsilon \lambda_\epsilon \frac{\partial P\delta_\epsilon}{\partial \lambda} \\ &= \int_\Omega K(\delta_\epsilon^{p-1+\epsilon} \\ & \quad + O(\theta_\epsilon^{p-1+\epsilon} + \delta_\epsilon^{p-2+\epsilon} \theta_\epsilon)) v_\epsilon \lambda_\epsilon \frac{\partial P\delta_\epsilon}{\partial \lambda} \\ &= \int_\Omega K \delta_\epsilon^{p-1+\epsilon} v_\epsilon \lambda_\epsilon \frac{\partial \delta_\epsilon}{\partial \lambda} \\ & \quad - \int_\Omega K \delta_\epsilon^{p-1+\epsilon} v_\epsilon \lambda_\epsilon \frac{\partial \theta_\epsilon}{\partial \lambda} \\ & \quad - O\left(\int_\Omega \delta_\epsilon^{p-1+\epsilon} |v_\epsilon| \theta_\epsilon\right) \\ &= \int_\Omega K(\delta_\epsilon)^{p-1+\epsilon} v_\epsilon \lambda_\epsilon \frac{\partial \delta_\epsilon}{\partial \lambda} \\ & \quad - O\left(\frac{\lambda_\epsilon^{\epsilon(n-2)} |v_\epsilon|_{H_0^1}}{(\lambda_\epsilon d_\epsilon^2)^{(n-2)/2}} \left(\int_\Omega \delta_\epsilon^{\frac{8n}{n^2-4}}\right)^{(n+2)/2}\right) \\ & \quad + O\left(\lambda_\epsilon^{\epsilon(n-2)} |v_\epsilon|_{H_0^1} \left(\int_\Omega \delta_\epsilon^{2n/(n-2)}\right)^{(n+2)/(2n)}\right) \end{aligned}$$

Using (15) we find

$$\begin{aligned} & \int_\Omega K(P\delta_\epsilon)^{p-1+\epsilon} v_\epsilon \lambda_\epsilon \frac{\partial P\delta_\epsilon}{\partial \lambda} = \int_\Omega K \delta_\epsilon^{p-1+\epsilon} v_\epsilon \lambda_\epsilon \frac{\partial \delta_\epsilon}{\partial \lambda} + \\ & \quad O\left(\frac{\lambda_\epsilon^{\frac{\epsilon(n-2)}{2}} |v_\epsilon|_{H_0^1}}{(\lambda_\epsilon d_\epsilon)^{\frac{(n+2)}{2}}}\right) + O\left(\lambda_\epsilon^{\frac{\epsilon(n-2)}{2}} |v_\epsilon|_{H_0^1} \left[\frac{1}{(\lambda_\epsilon d_\epsilon)^{n-2}} (\text{if } n < \right. \right. \\ & \quad \left. \left. 6) + \frac{\log(\lambda_\epsilon d_\epsilon)}{(\lambda_\epsilon d_\epsilon)^4} (\text{if } n=6) + \frac{1}{(\lambda_\epsilon d_\epsilon)^{\frac{(n+2)}{2}}} (\text{if } n > 6)\right]\right) \end{aligned} \tag{24}$$

We also have, using Remark 4

$$\begin{aligned} & \int_{\Omega} K(\delta_{\epsilon})^{p-1+\epsilon} v_{\epsilon} \lambda_{\epsilon} \frac{\partial \delta_{\epsilon}}{\partial \lambda} \\ &= \lambda_{\epsilon}^{\frac{\epsilon(n-2)}{2}} c_0^{\epsilon} \int_{\Omega} K \delta_{\epsilon}^{p-1} v_{\epsilon} \lambda_{\epsilon} \frac{\partial \delta_{\epsilon}}{\partial \lambda} \\ &+ \int_{\Omega} K \left( \delta_{\epsilon}^{p-1+\epsilon} \right. \\ &\left. - c_0^{\epsilon} \lambda_{\epsilon}^{\frac{\epsilon(n-2)}{2}} \delta_{\epsilon}^{p-1} \right) v_{\epsilon} \frac{\partial \delta_{\epsilon}}{\partial \lambda} \\ &= O \left( \epsilon \int_{\Omega} K \log(1 \right. \\ &\left. + \lambda_{\epsilon}^2 |x - a_{\epsilon}|^2) \delta_{\epsilon}^p |v_{\epsilon}| \right) \end{aligned}$$

whence

$$\frac{\int_{\Omega} K(\delta_{\epsilon})^{p-1+\epsilon} v_{\epsilon} \lambda_{\epsilon} \frac{\partial \delta_{\epsilon}}{\partial \lambda}}{O(\epsilon)} = O \left( \epsilon |v_{\epsilon}|_{H_0^1} \right) = O(\epsilon). \tag{25}$$

Noticing that in addition  $\lambda_{\epsilon} \frac{\partial P \delta_{\epsilon}}{\partial \lambda} = O(\delta_{\epsilon})$  and

$$\int_{\Omega} \delta_{\epsilon}^{p-1+\epsilon} |v_{\epsilon}|^2 = O \left( \lambda_{\epsilon}^{\epsilon(n-2)/2} |v_{\epsilon}|_{H_0^1}^2 \right). \tag{26}$$

$$\int_{\delta < |v_{\epsilon}|} |v_{\epsilon}|^{p+\epsilon} \delta_{\epsilon} = O \left( \lambda_{\epsilon}^{\epsilon(n-2)/2} |v_{\epsilon}|_{H_0^1}^{p+1} \right). \tag{27}$$

(18), (23), (24),(25), (26), (27) and Lemmas 3 and 5 prove Proposition 6.

We are now able to prove the theorem.

**Proof of Theorem 1**

Arguing by contradiction, let us suppose that  $(P_{\epsilon})$  has a solution  $u_{\epsilon}$  as stated in the theorem. From Proposition 6, we have

$$\begin{aligned} \alpha_{\epsilon} c_1 \frac{H(a_{\epsilon}, a_{\epsilon})}{\lambda_{\epsilon}^{n-2}} - \alpha_{\epsilon} \frac{c_3}{n^2} \frac{\Delta K(a_{\epsilon})}{K(a_{\epsilon}) \lambda_{\epsilon}^2} + \alpha_{\epsilon} c_2 \epsilon &= o \left( \epsilon + \frac{1}{\lambda_{\epsilon}^2} + \right. \\ &\left. \begin{cases} \frac{1}{(\lambda_{\epsilon} d_{\epsilon})^{n-2}} & (\text{if } n \geq 4) \\ \frac{1}{\lambda_{\epsilon} d_{\epsilon}} & (\text{if } n = 4) \end{cases} \right) \end{aligned} \tag{28}$$

Notice that  $H(a_{\epsilon}, a_{\epsilon}) \sim d_{\epsilon}^{n-2}$  if  $d_{\epsilon} \rightarrow 0$  as  $\epsilon \rightarrow 0$  and  $H(a_{\epsilon}, a_{\epsilon}) \geq c > 0$  as  $\epsilon \rightarrow 0$  if  $d_{\epsilon} \not\rightarrow 0$  as  $\epsilon \rightarrow 0$ .

For  $n = 3$ , it follows from (28) that

$$c_1 \frac{H(a_{\epsilon}, a_{\epsilon})}{\lambda_{\epsilon}} + c_2 \epsilon = o \left( \epsilon + \frac{1}{\lambda_{\epsilon}} \right)$$

which is a contradiction.

For  $n = 4$  it follows from (28) that

$$c_1 \frac{H(a_{\epsilon}, a_{\epsilon})}{\lambda_{\epsilon}^2} - \frac{c_3}{16} \frac{\Delta K(a_{\epsilon})}{K(a_{\epsilon}) \lambda_{\epsilon}^2} + c_2 \epsilon = o \left( \epsilon + \frac{1}{\lambda_{\epsilon}^2} + \frac{1}{(\lambda_{\epsilon} d_{\epsilon})^2} \right)$$

which is a contradiction with assumption (ii) of the theorem.

For  $n \geq 5$ , it follows from (28) that

$$- \frac{c_3}{n^2} \frac{\Delta K(a_{\epsilon})}{K(a_{\epsilon}) \lambda_{\epsilon}^2} + c_2 \epsilon = o \left( \epsilon + \frac{1}{\lambda_{\epsilon}^2} \right)$$

also leads to a contradiction with assumption (iii).

**References**

- [1] A. Bahri, *Critical point at infinity in some variational problems*, Pitman Res. Notes Math. Ser. 182, Longman Sci. Tech. Harlow 1989.
- [2] A. Bahri and J. M. Coron, *On a nonlinear elliptic equation involving the Sobolev exponent : the effect of the topology of the domain*, Comm. Pure Appl. Math. 41 (1988), 253-294
- [3] A. Bahri, Y.Y. Li and O. Rey, *On a variational problem with lack of compactness: the topological effect of the critical points at infinity*, Calc. Var. and Part. Diff. Equ. 3 (1995), 67-94.
- [4] M. Ben Ayed, K. El Mehdi, M. Grossi and O. Rey, *A Nonexistence result of single peaked solutions to a supercritical nonlinear problem*, Comm. Contemporary Math., 2 (2003), 179-195.
- [5] M. Ben Ayed, K. Ould Bouh, *Nonexistence results of sign-changing solutions to a supercritical nonlinear problem*, Comm. Pure Applied Anal, 5 (2007), 1057-1075.
- [6] M. Del Pino, P. Felmer and M. Musso, *Two bubbles solutions in the supercritical Bahri-Coron's problem*, Calc. Var. Part. Diff. Equat., 16 (2003), 113-145.
- [7] Z. C. Han, *Asymptotic approach to singular solutions for nonlinear elliptic equations involving critical Sobolev exponent*, Ann. Inst. Henri Poincare (Analyse non-linear) 8(1991),159-174.
- [8] K. Ould Bouh, *Nonexistence result of sign-changing solutions for a supercritical problem of the scalar curvature type*, Advance in Nonlinear Studies (ANS), 12 (2012), 149-171.
- [9] O. Rey, *The role of Green's function in a nonlinear elliptic equation involving critical Sobolev exponent*, J. Funct. Anal. 89 (1990), 1-52.
- [10] O. Rey, *The topological impact of critical points at infinity in a variational problem with lack of compactness : the dimension 3*, Adv. Diff. Equ. 4 (1999), 581-616.