

# Some Separate Quasi-Asymptotics Properties of Multidimensional Distributions and Application

Nenad Stojanovic

Department of Mathematics, Faculty of Agriculture, University of Banja Luka, Banja Luka, Bosnia and Herzegovina

## Email address:

nenad.stojanovic@agro.unibl.org

## To cite this article:

Nenad Stojanovic. Some Separate Quasi-Asymptotics Properties of Multidimensional Distributions and Application. *Pure and Applied Mathematics Journal*. Vol. 9, No. 3, 2020, pp. 64-69. doi: 10.11648/j.pamj.20200903.13

Received: June 17, 2020; Accepted: July 3, 2020; Published: July 13, 2020

**Abstract:** Quasi-asymptotic behavior of functions as a method has its application in observing many physical phenomena which are expressed by differential equations. The aim of the asymptotic method is to allow one to present the solution of a problem depending on the large (or small) parameter. One application of asymptotic methods in describing physical phenomena is the quasi-asymptotic approximation. The aim of this paper is to look at the quasi-asymptotic properties of multidimensional distributions by extracted variable. Distribution  $T(x_0, x)$  from  $S'(\bar{R}_+^{1 \times R^n})$  has the property of the separability of variables, if it can be represented in form  $T(x_0, x) = \sum \varphi_i(x_0) \psi_i(x)$  where distributions,  $\varphi_i(x_0)$  from  $S'(\bar{R}_+^1)$  and  $\psi_i$  from  $S(R^n)$ ,  $x_0$  from  $\bar{R}_+^1$  and  $x$  is element  $R^n$  different values of do not depend on each other. Distribution  $T(x_0, x)$  the element  $S'(\bar{R}_+^{1 \times R^n})$  is homogeneous and of order  $\alpha$  at variable  $x_0$  is element  $\bar{R}_+^1$  and  $x = x_1, x_2, \dots, x_n$  from  $R^n$  if for  $k > 0$  it applies that  $T(kx_0, kx) = k^\alpha T(x_0, x)$ . The method of separating variables is one of the most widespread methods for solving linear differential equations in mathematical physics. In this paper, the results by V. S Vladimirov are used to present the proof of the basic theorems, regarding the quasi-asymptotic behavior of multidimensional distributions by a singular variable, with the application of quasi-asymptotics to the solution of differential equations.

**Keywords:** Distribution Spaces, Asymptotics, Separate Quasi-Asymptotics, Multidimensional Distributions

## 1. Introduction

We use  $S(\mathbb{R}_0^n)$  to mark the standard space of the Schwartz's rapidly decreasing functions, and  $S'(\mathbb{R}_0^n)$  to mark the corresponding space of the slowly increasing distributions [1, 7].

If  $f(t) \in S'$  and  $\varrho(k)$  is a positive and a continuous function for  $k > 0$ , distribution  $f(t)$  has quasi-asymptotics at infinity (at zero) with respect to positive function  $\varrho(k)$ , if the following is valid

$$\frac{1}{\varrho(k)} f(kt) \rightarrow g(t), \left( \frac{1}{\varrho(k)} f\left(\frac{t}{k}\right) \rightarrow g(t) \right) \quad (1)$$

$k \rightarrow \infty$  in  $S'(\mathbb{R}_0^n)$  with the distribution being  $g(t) S'(\mathbb{R}_0^n)$ , [1, 2-4, 7].

If  $g(t) = 0$  distribution  $f(t)$  has a trivial quasi-asymptotics at infinity (that is, at zero) with respect to positive function  $\varrho(k)$ . If (1) is true, function  $\varrho(k)$  occurs as an auto-modal function. If  $\varrho(k)$  is a positive and continuous function, and  $k \rightarrow \infty$ , then we say that  $\varrho(k)$  is an auto-modal

function, if for a real number  $\alpha > 0$  there exists

$$\lim_{k \rightarrow \infty} \frac{\varrho(ak)}{\varrho(k)} = a^\alpha \quad (2)$$

Where by it converges evenly along  $a$  on each compact semi-axes  $(0, \infty)$ . Distribution  $f(t) \in S'_+$  is asymptotically homogeneous with respect to function  $\varrho(k)$  of order  $\alpha$  if:

$$\frac{1}{\varrho(k)} f(kt) \rightarrow C \cdot f_{\alpha+1}(t) \text{ in } S'_+ \quad (3)$$

where the nucleus of fractional differentiation and integration  $f_\alpha(t) \in S'$  is defined by

$$f_\alpha(t) = \begin{cases} \frac{\theta(t) \cdot t^{\alpha-1}}{\Gamma(\alpha)}, & \text{if } \alpha > 0 \\ \frac{d^N}{dt^N} f_{\alpha+N}(t), & \text{if } \alpha \leq 0, \alpha + N > 0 \end{cases} \quad (4)$$

with  $\Gamma(\alpha)$  being the gamma function, and  $\theta(t)$  being the Heaviside function [1-3, 6, 7].

The fractional derivative, [3-5, 7], of order  $\alpha$  and

distribution of  $f(t) \in S'(\mathbb{R}_1)$  is defined by the formula

$$f^{(-\alpha)}(t) = f_\alpha(t) * f(t) \quad (5)$$

Distribution  $T(x_0, x) \in S'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$  has the property of the separability of variables, if it can be represented in form  $T(x_0, x) = \sum_i \varphi_i(x_0) \psi(x)$  where distributions  $\varphi(x_0) \in S'(\mathbb{R}_+^1)$  and  $\psi \in S(\mathbb{R}^n)$ ,  $x_0 \in \mathbb{R}_+^1$  and  $x \in \mathbb{R}^n$  for different values of  $i$  do not depend on each other, [15].

Distribution  $T(x_0, x) \in S'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$  is homogeneous and of order  $\alpha$  at variable  $x_0 \in \mathbb{R}_+^1$  and  $x = x_1, x_2, \dots, x_n \in \mathbb{R}^n$  if for  $k > 0$  it applies that  $T(kx_0, kx) = k^\alpha T(x_0, x)$ , [1, 3, 7, 8].

In other words, distribution  $T(x_0, x) \in S'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$  is homogeneous and of order  $\alpha$  at variable  $x_0$  and  $x$  if for each test function  $\phi(x_0, x) \in S(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$  the following is valid

$$\langle T(x_0, x), \phi\left(\frac{x_0}{k}, \frac{x}{k}\right) \rangle = k^{2\alpha+n+1} \langle T(x_0, x), \phi(x_0, x) \rangle,$$

$k > 0$ . Indeed,

$$\begin{aligned} \langle T(kx_0, kx), \phi(x_0, x) \rangle &= \\ &= \left\langle \begin{cases} \text{shift } kx_0 = x'_0 \Rightarrow x_0 = \frac{x'_0}{k}, \\ kx = x' \Rightarrow x = \frac{x'}{k} \end{cases} \right\rangle \\ &= \frac{1}{k^{n+1}} \langle T(x'_0, x'), \phi\left(\frac{x'_0}{k}, \frac{x'}{k}\right) \rangle = \left( \begin{matrix} x'_0 = x_0 \\ x' = x \end{matrix} \right) \\ &= \frac{1}{k^{n+1}} \langle T(x_0, x), \phi\left(\frac{x_0}{k}, \frac{x}{k}\right) \rangle \end{aligned} \quad (6)$$

For example, let it be that  $T(x_0, x) \in S'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$  in the form of  $T(x_0, x) = f(x_0) \times g(x)$  with distributions  $f(x_0) \in S'(\mathbb{R}_+^1)$ ,  $g(x) \in S'(\mathbb{R}_0^n)$  being homogenous and of order  $\alpha$ . Then, there is a number of equations that are valid:

$$\begin{aligned} \langle T(kx_0, kx), \phi(x_0, x) \rangle &= \langle f(kx_0) \times g(kx), \phi(x_0, x) \rangle \\ &= \langle f(kx_0), \langle g(kx), \phi(x_0, x) \rangle \rangle = \left( \begin{matrix} f(kx_0) = k^\alpha f(x_0) \\ g(kx) = k^\alpha g(x) \end{matrix} \right) \\ &= k^{2\alpha} \langle f(x_0), \langle g(x), \phi(x_0, x) \rangle \rangle \\ &= k^{2\alpha} \langle T(x_0, x), \phi(x_0, x) \rangle. \end{aligned} \quad (7)$$

From (6) and (7) we can see that the following equation is valid

$$\frac{1}{k^{n+1}} \langle T(x_0, x), \phi\left(\frac{x_0}{k}, \frac{x}{k}\right) \rangle = k^{2\alpha} \langle T(x_0, x), \phi(x_0, x) \rangle$$

and from here, there is

$$\langle T(x_0, x), \phi\left(\frac{x_0}{k}, \frac{x}{k}\right) \rangle = k^{2\alpha+n+1} \langle T(x_0, x), \phi(x_0, x) \rangle.$$

For example, distribution  $T(x_0, x) \in S'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$  in the form of  $T(x_0, x) = f(x_0) \times g(x)$  with  $f(x_0) \in S'(\mathbb{R}_+^1)$ ,  $g(x) \in S'(\mathbb{R}_0^n)$ , and  $f(x_0)$  being homogenous and of order  $\alpha$ , and test function  $\phi(x_0, x)$  in the fom of

$\phi(x_0, x) = \sum_i \varphi_i^1(x_0) \varphi_i^2(x)$  with  $\varphi_i^1(x_0) \in S(\mathbb{R}_+^1)$ ,  $\varphi_i^2(x) \in S(\mathbb{R}^n)$  and  $\varphi^1(x_0)$ , is a homogenous distribution of order  $\alpha$  and then, for  $(\forall i)$  the following is true

$$\begin{aligned} \langle T(kx_0, x), \phi(x_0, x) \rangle &= \langle f(kx_0) \times g(x), \phi(x_0, x) \rangle \\ &= \langle f(kx_0), \langle g(x), \phi(x_0, x) \rangle \rangle \\ &= \langle f(kx_0), \langle g(x), \sum_i \varphi_i^1(x_0) \varphi_i^2(x) \rangle \rangle \\ &= \sum_i \langle f(kx_0), \varphi_i^1(x_0) \rangle \langle g(x), \varphi_i^2(x) \rangle \\ &= k^\alpha \sum_i \langle f(kx_0), \varphi_i^1(x_0) \rangle \langle g(x), \varphi_i^2(x) \rangle \\ &= k^\alpha \langle f(x_0) g(x), \phi(x_0, x) \rangle = k^\alpha \langle T(x_0, x), \phi(x_0, x) \rangle \end{aligned}$$

since the set of functions  $\sum_i \varphi_i^1(x_0) \varphi_i^2(x)$  is dense in  $S(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$ . This is followed by the claim, because it is valid in a dense set, and with its continuity, it extends to entire set in  $S(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$ . Homogeneity by the second variable is similarly defined [1, 6].

The homogeneity of distribution  $T(x_0, x) \in S'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$  separable at variable  $x_0$  assuming that distribution  $\varphi_i^1(x_0) \in S(\mathbb{R}_+^1)$  is homogeneous and of order  $\alpha$  for each  $i$ , then, from these relations, it follows that

$$\begin{aligned} T(kx_0, x) &= \sum_i \varphi_i(kx_0) \psi(x) \\ &= k^\alpha \sum_i \varphi_i(x_0) \psi(x) = k^\alpha T(x_0, x). \end{aligned} \quad (8)$$

Homogeneity at variable  $x$  is similarly observed.

Let there be distribution  $T(x_0, x) \in S'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$ . Distribution  $T(x_0, x)$  with  $x_0 \in \mathbb{R}_+^1$  and  $x \in \mathbb{R}^n$  has quasi-asymptotic at infinity at variable  $x_0$  relative to auto-modal function  $\varrho$ , if there is distribution  $G(x_0, x) \neq 0$  such that

$$\lim_{k \rightarrow \infty} \frac{1}{\rho(k)} T(kx_0, x) = G(x_0, x) \text{ in } S'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n). \quad (9)$$

Quasi-asymptotics by the separated variable at zero is similarly defined, [1, 7].

Let us suppose a distribution  $T(x_0, x) \in S'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$ . Distribution  $T(x_0, x)$ ,  $x_0 \in \mathbb{R}_+^1$  and  $x \in \mathbb{R}^n$  has quasi-asymptotics at zero at variable  $x_0$  with respect to auto-modal function  $\varrho$ , if, and only if there is distribution  $G(x_0, x) \neq 0$  such that

$$\lim_{k \rightarrow \infty} \frac{1}{\rho(k)} T\left(\frac{x_0}{k}, x\right) = G\left(\frac{x_0}{k}, x\right) \neq 0 \quad (10)$$

in  $S'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$ .

For distributions from  $D'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$  (or  $S'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$ ) we define the fractional (rational) differentiation at variable  $x_0$  as a convolution  $f_\alpha(x_0)$  with  $f(x_0, x)$  at  $x_0$  by the following formula

$$f^{(\alpha)} = f_{-\alpha}(x_0) * f(x_0, x) \quad (11)$$

which belongs to  $D'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$  if  $f \in D'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$  that is,  $\mathcal{S}'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$  if  $f \in \mathcal{S}'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$ , (more in [3-6, 8, 12, 13]).

## 2. Some Quasi-Asymptotics Properties of Multidimensional Distributions

We provide proof of some of the basic theorems that apply to multidimensional distributions, and their formulaic presentation can be seen in [1].

**Theorem 1.** *If distribution  $T(x_0, x) \in \mathcal{S}'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$  is asymptotically homogeneous with respect to positive function  $\rho(k)$  at variable  $x_0$  or if the following is true*

$$\lim_{k \rightarrow \infty} \frac{1}{\rho(k)} T(kx_0, x) = G(x_0, x) \text{ in } \mathcal{S}'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n) \quad (12)$$

then  $\rho(k)$  is an auto-modal function.

**Proof:** Let (12) be true and let  $\phi(x_0, x) \in \mathcal{S}(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$  test function such that  $\langle G(x_0, x), \phi(x_0, x) \rangle \neq 0$ .

Then let the test function be of the following form  $\phi(x_0, x) = \sum_i \varphi_i^1(x_0) \varphi_i^2(x)$ , so that  $\varphi_i^1(x_0) \in \mathcal{S}(\overline{\mathbb{R}_+^1})$ ,  $\varphi_i^2(x) \in \mathcal{S}(\mathbb{R}^n) \forall i$ , are continuous functions with the following feature:

$$\text{supp } \varphi_i^1 \subset \overline{\mathbb{R}_+^1}, \text{supp } \varphi_i^2 \subset \mathbb{R}_+^n,$$

$$\text{supp } \phi = \text{supp } \varphi_i^1 \times \text{supp } \varphi_i^2 \subset (\overline{\mathbb{R}_+^1} \times \mathbb{R}^n), (\forall i),$$

$$K \subset \overline{\mathbb{R}_+^1} \text{ compact set.}$$

For  $\phi(x_0, x)$  and  $a \in K$  it applies that

$$\frac{1}{a} \phi\left(\frac{x_0}{a}, x\right) = \frac{1}{a} \sum_i \varphi_i^1\left(\frac{x_0}{a}\right) \varphi_i^2(x).$$

Now, the following is valid for distribution  $T(x_0, x) \in \mathcal{S}'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$  and test function

$$\phi(x_0, x) \in \mathcal{S}(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n);$$

$$\langle T(kx_0, x), \frac{1}{a} \phi\left(\frac{x_0}{a}, x\right) \rangle \xrightarrow{k \rightarrow \infty, a \in K} \langle G(x_0, x), \frac{1}{a} \phi\left(\frac{x_0}{a}, x\right) \rangle$$

For  $a \in K$ , and using  $(x_0 = ax'_0)$  the following is valid

$$\begin{aligned} & \frac{\rho(ak)}{\rho(k)} \langle \frac{T(kx_0, x)}{\rho(ak)}, \frac{1}{a} \phi\left(\frac{x_0}{a}, x\right) \rangle \\ &= \frac{\rho(ak)}{\rho(k)} \langle \frac{T(akx'_0, x)}{\rho(ak)}, \phi(x'_0, x) \rangle \\ &= \frac{\rho(ak)}{\rho(k)} \langle \frac{T(akx_0, x)}{\rho(ak)}, \phi(x_0, x) \rangle \\ & \xrightarrow{k \rightarrow \infty, a \in K} \langle G(x_0, x), \frac{1}{a} \phi\left(\frac{x_0}{a}, x\right) \rangle \end{aligned} \quad (13)$$

Further, if we replace  $k$  with  $ak, a \in K$ , the following is valid

$$\frac{1}{\rho(ak)} \langle T(akx_0, x), \phi(x_0, x) \rangle \xrightarrow{k \rightarrow \infty} \langle G(x_0, x), \phi(x_0, x) \rangle. \quad (14)$$

Using relations (12) and (13), we get the following relation

$$\frac{\rho(ak)}{\rho(k)} \xrightarrow{k \rightarrow \infty} \frac{\langle G(x_0, x), \frac{1}{a} \phi\left(\frac{x_0}{a}, x\right) \rangle}{\langle G(x_0, x), \phi(x_0, x) \rangle}. \quad (15)$$

From here, by inserting  $(ax'_0 = x_0)$  we get the following

$$\frac{\rho(ak)}{\rho(k)} \xrightarrow{k \rightarrow \infty} \frac{\langle G(ax'_0, x), \phi(x'_0, x) \rangle}{\langle G(x_0, x), \phi(x_0, x) \rangle}.$$

From here, we get the required relation

$$\frac{\rho(ak)}{\rho(k)} \xrightarrow{k \rightarrow \infty} \frac{\langle G(ax_0, x), \phi(x_0, x) \rangle}{\langle G(x_0, x), \phi(x_0, x) \rangle} = C(a).$$

From the existence of  $\lim_{k \rightarrow \infty} \frac{\rho(ak)}{\rho(k)} = C(a)$  following  $C(a) = a^\alpha$  and  $\rho(a) = a^\alpha L(a)$ , and Karamata L function [16], it follows that function  $\rho(k)$  is an auto-modal function, even in the case of multi-variable distributions.

**Theorem 2.** *Let distribution  $T(x_0, x) \in \mathcal{S}'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$  be asymptotically homogeneous with respect to positive function  $\rho(k)$  at variable  $x_0$ . In this case, if the order of auto-modal function  $\rho(k)$  is equal to  $\alpha$ , then distribution  $G(x_0, x)$  in the following equation*

$$\lim_{k \rightarrow \infty} \frac{1}{\rho(k)} T(kx_0, x) = G(x_0, x) \text{ is equal to}$$

$$G(x_0, x) = C f_\alpha(x_0) \times g(x), \text{ with } C \text{ being the constant.}$$

**Proof:** It has already been shown in the case of distributions of one variable [1],[7], that distribution  $G(x) \in \mathcal{S}_+'$  has the form of  $G(x) = C f_\alpha(x)$  with  $C$  being the constant, and  $f_\alpha(x)$  being the nucleus of fractional differentiation.

For  $G(x_0, x) \in \mathcal{S}'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$  let us suppose that

$f(x_0) \in \mathcal{S}'(\overline{\mathbb{R}_+^1})$ , and  $g(x) = \mathcal{S}'(\mathbb{R}^n)$ , and that distribution  $f(x_0)$  is homogeneous and of order  $\alpha$ , and  $G(x_0, x) = f(x_0) \times g(x)$ . Since for the function in the form of  $\varphi(x_0)\psi(x) \in \mathcal{S}'(\overline{\mathbb{R}_+^1} \times \mathbb{R}^n)$  the following applies

$$\lim_{k \rightarrow \infty} \langle \frac{1}{\rho(k)} T(x_0, x), \varphi(x_0) \rangle, \psi(x) \rangle$$

$$= \langle C g(x) f_{\alpha+1}(x_0), \psi(x) \rangle = C f_{\alpha+1}(x_0) \langle g(x), \psi(x) \rangle$$

so distribution  $g(x) = \mathcal{S}'(\mathbb{R}^n)$ ,  $G(x_0, x)$  is in the form of  $G(x_0, x) = C f_{\alpha+1}(x_0) \times g(x)$ .

**Theorem 3.** *If distribution  $T(x_0, x)$  is separated at variable forms  $x_0$ , then it has the following form:*

$T(x_0, x) = T_1 g_1(x) + T_2(x_0) g_2(x_0)$  and distribution  $T(x_0, x)$  has the quasi-asymptotics of order  $\alpha$  in relation to

function  $k^\alpha \rho(k)$  at a variable  $x_0$ , if  $T_1$  and  $T_2$  have the same quasi-asymptotics in relation to function  $\rho(k)$ . The reverse of the theorem is not valid.

**Proof.** Let us show that distribution  $T(x_0, x)$  has quasi-asymptotics of order  $\alpha$  with respect to  $\rho(k)$  if  $T_1$  and  $T_2$  have the same quasi-asymptotics. Let the test function  $\phi(x_0, x)$  be in the form of  $\phi(x_0, x) = \sum_i \varphi_i(x_0) \psi_i(x)$ . By the definition of quasi-asymptotics, the following applies:

$$\begin{aligned}
\frac{1}{\rho(k)} \langle T(kx_0, x), \phi(x_0, x) \rangle &= \left\langle \frac{T(kx_0, x)}{\rho(k)}, \sum_i \varphi_i(x_0) \psi_i(x) \right\rangle \\
&= \left\langle \frac{T_1(kx_0)g_1(x) + T_2(kx_0)g_2(x)}{k^\alpha \rho(k)}, \sum_i \varphi_i(x_0) \psi_i(x) \right\rangle \\
&= \left\langle \frac{T_1(kx_0)g_1(x)}{\rho(k)}, \sum_i \varphi_i(x_0) \psi_i(x) \right\rangle + \left\langle \frac{T_2(kx_0)g_2(x)}{\rho(k)}, \sum_i \varphi_i(x_0) \psi_i(x) \right\rangle.
\end{aligned}$$

Since  $\rho(k) = k^\alpha L(k)$  and  $T_1(kx_0) = k^\alpha T_1(x_0)$  and  $T_2(kx_0) = k^\alpha T_2(x_0)$  therefore

$$\begin{aligned}
&= \left\langle \frac{T_1(kx_0)g_1(x)}{k^\alpha L(k)}, \sum_i \varphi_i(x_0) \psi_i(x) \right\rangle + \left\langle \frac{T_2(kx_0)g_2(x)}{k^\alpha L(k)}, \sum_i \varphi_i(x_0) \psi_i(x) \right\rangle \\
&= \frac{1}{k^\alpha L(k)} \langle T_1(kx_0), \langle g_1(x), \sum_i \varphi_i(x_0) \psi_i(x) \rangle \rangle + \frac{1}{k^\alpha L(k)} \langle T_2(kx_0), \langle g_2(x), \sum_i \varphi_i(x_0) \psi_i(x) \rangle \rangle \\
&= \frac{1}{k^\alpha L(k)} \sum_i \langle T_1(kx_0), \varphi_i(x_0) \rangle \langle g_1(x), \psi_i(x) \rangle + \frac{1}{k^\alpha L(k)} \sum_i \langle T_2(kx_0), \varphi_i(x_0) \rangle \langle g_2(x), \psi_i(x) \rangle \\
&= \frac{k^\alpha}{k^\alpha L(k)} \sum_i \langle T_1(kx_0), \varphi_i(x_0) \rangle \langle g_1(x), \psi_i(x) \rangle + \frac{k^\alpha}{k^\alpha L(k)} \sum_i \langle T_2(kx_0), \varphi_i(x_0) \rangle \langle g_2(x), \psi_i(x) \rangle \\
&= \frac{1}{L(k)} \sum_i \langle T_1(kx_0), \varphi_i(x_0) \rangle \langle g_1(x), \psi_i(x) \rangle + \frac{1}{L(k)} \sum_i \langle T_2(kx_0), \varphi_i(x_0) \rangle \langle g_2(x), \psi_i(x) \rangle \\
&= \frac{1}{L(k)} \langle T_1(x_0)g_1(x), \sum_i \varphi_i(x_0) \psi_i(x) \rangle + \frac{1}{L(k)} \langle T_2(x_0)g_2(x), \sum_i \varphi_i(x_0) \psi_i(x) \rangle \\
&= \frac{1}{L(k)} \langle T_1(x_0)g_1(x), \phi(x_0, x) \rangle + \frac{1}{L(k)} \langle T_2(x_0)g_2(x), \phi(x_0, x) \rangle \\
&= \frac{1}{L(k)} \langle T_1(x_0)g_1(x) + T_2(x_0)g_2(x), \phi(x_0, x) \rangle.
\end{aligned}$$

This shows that distribution  $T(x_0, x) = T_1 g_1(x) + T_2(x_0)g_2(x)$  has quasi-asymptotics of order  $\alpha$  with respect to function  $k^\alpha \rho(k)$  at variable  $x_0$  if distributions  $T_1$  and  $T_2$  have the same quasi-asymptotics.

The reverse of the theorem is not valid. To show this, it is

enough to show that, for example, the following is not valid for distribution  $T_1(x_0) = x_0^{\alpha+1} + x_0^\alpha$  and  $T_2(x_0) = -x_0^{\alpha+1} + x_0^\alpha$  with respect to function  $k^\alpha \rho(k)$ . Indeed

$$\begin{aligned}
&\frac{1}{\rho(k)} \langle T(kx_0, x), \phi(x_0, x) \rangle \\
&= \left\langle \frac{T(kx_0, x)}{\rho(k)}, \sum_i \varphi_i(x_0) \psi_i(x) \right\rangle = \left\langle \frac{k^\alpha x_0^\alpha [kx_0(g_1(x) - g_2(x)) + g_1(x) + g_2(x)]}{\rho(k)}, \sum_i \varphi_i(x_0) \psi_i(x) \right\rangle \\
&= \left\langle \frac{k^\alpha x_0^\alpha [kx_0(g_1(x) - g_2(x)) + g_1(x) + g_2(x)]}{k^\alpha L(k)}, \sum_i \varphi_i(x_0) \psi_i(x) \right\rangle \\
&= \left\langle \frac{x_0^\alpha [kx_0(g_1(x) - g_2(x)) + g_1(x) + g_2(x)]}{L(k)}, \sum_i \varphi_i(x_0) \psi_i(x) \right\rangle \\
&= \left\langle \frac{x_0^\alpha [(kx_0 + 1)g_1(x) + (kx_0 - 1)g_2(x)]}{L(k)}, \sum_i \varphi_i(x_0) \psi_i(x) \right\rangle.
\end{aligned}$$

From here, it can be seen that  $T(x_0, x)$  has no quasi-asymptotics when  $k \rightarrow \infty$ .

Theorem 4. In order for distribution  $T(x_0, x) \in \mathcal{S}'(\overline{\mathbb{R}}_+^1 \times$

$\mathbb{R}^n)$  to be asymptotically homogeneous at infinity, with respect to auto-modal function  $\rho(k)$  at variable  $x_0$ , it is necessary, and it is also sufficient, that for each  $\beta \in \mathbb{R}$  its

fractional derivative  $T^{(-\beta)}(x_0, x)$  is asymptotically homogeneous with respect to  $k^\beta \rho(k)$ .

Proof: We define fractional differentiation in  $\mathcal{S}'(\overline{\mathbb{R}}_+^1 \times \mathbb{R}^n)$  with distribution  $T(x_0, x)$  at  $x_0$  as convolution of distribution  $f_\beta(x_0) \in \mathcal{S}'(\overline{\mathbb{R}}_+^1)$  and distribution  $T(x_0, x) \in \mathcal{S}'(\overline{\mathbb{R}}_+^1 \times \mathbb{R}^n)$  i.e.  $T^{(-\beta)}(x_0, x) = T(x_0, x) * f_\beta(x_0)$ . Using the property of distribution  $f_\beta(x_0)$  to be homogeneous and of order  $\beta - 1$ , that is, using the validity of the following  $f_\beta(kx_0) = k^{\beta-1} f_\beta(x_0)$ , we get the following:

$\lim_{k \rightarrow \infty} \frac{1}{k^\beta \rho(k)} \langle T^{(-\beta)}(kx_0, x), \phi(x_0, x) \rangle$ , from here, if we put that  $kx_0 = x'_0$ , we get

$$\begin{aligned} &= \lim_{k \rightarrow \infty} \frac{1}{k^{\beta+1} \rho(k)} \langle T^{(-\beta)}(x'_0, x), \phi\left(\frac{x'_0}{k}, x\right) \rangle \\ &= \lim_{k \rightarrow \infty} \frac{1}{k^{\beta+1} \rho(k)} \langle T^{(-\beta)}(x_0, x), \phi\left(\frac{x_0}{k}, x\right) \rangle. \end{aligned}$$

By using the definition of convolution

$$\begin{aligned} T(x_0, x) * f_\beta(x_0) &= \frac{1}{\Gamma(\beta)} \Theta(x_0) x_0^{\beta-1} * T(x_0, x) \\ &= \frac{1}{\Gamma(\beta)} \int_0^\infty (x, x_0)^{\beta-1} T(t, x) dt = T^{(-\beta)}(x_0, x). \end{aligned}$$

we can see that the last equation is precisely the  $\beta$  primitive integral for  $T(x_0, x)$ . Based on this, we have that  $T(x_0, x) \in \mathcal{S}'(\overline{\mathbb{R}}_+^1 \times \mathbb{R}^n, f_\beta(x_0) \in \mathcal{S}'(\overline{\mathbb{R}}_+^1)$ ,

$$\begin{aligned} &\langle T(x_0, x) * f_\beta(x_0), \phi(x_0, x) \rangle \\ &= \lim_{k \rightarrow \infty} \langle T(x_0, x) * f_\beta(\tau), \eta_k(x_0, \tau) \phi(x_0 + \tau, x) \rangle, \end{aligned}$$

with  $\{\eta_k\}$  being unit sequence. If there is a limes on the right-hand side for each series  $\{\eta_k, k \rightarrow \infty\}$  then the function from  $S(\mathbb{R}^2)$  which converges to number one in  $\mathbb{R}^2$  and this limit does not depend on the choice of series  $\{\eta_k, k \rightarrow \infty\}$  then we have that  $T(x_0, x) * f_\beta(x_0) \in \mathcal{S}'(\mathbb{R}^{n+1})$ . Based on this, the last equation transforms into

$$\begin{aligned} &\lim_{k \rightarrow \infty} \frac{1}{k^{\beta+1} \rho(k)} \langle T(x_0, x) * f_\beta(x_0), \phi\left(\frac{x_0}{k}, x\right) \rangle \\ &= \lim_{k \rightarrow \infty} \frac{1}{k^{\beta+1} \rho(k)} \langle T(x_0, x) \times f_\beta(\tau), \eta_k(x_0; \tau) \phi\left(\frac{x_0 + \tau}{k}, x\right) \rangle. \end{aligned}$$

Now, if we put that

$$\left( \frac{x_0 + \tau}{k} = \frac{\tau = k\tau'}{k} = \frac{x_0 + k\tau}{k} = \frac{x_0}{k} + \tau' \right)$$

the last equation transforms into the following form:

$$\begin{aligned} &\lim_{k \rightarrow \infty} \frac{1}{k^\beta \rho(k)} \langle T(x_0, x), \langle f_\beta(k\tau'), \phi\left(\frac{x_0}{k} + \tau', x\right) \rangle \rangle \\ &\lim_{k \rightarrow \infty} \frac{1}{k^\beta \rho(k)} \langle T(x_0, x), \langle f_\beta(k\tau), \phi\left(\frac{x_0}{k} + \tau, x\right) \rangle \rangle \end{aligned}$$

(since  $f_\beta(k\tau) = k^{\beta-1} f_\beta(\tau)$ )

$$\lim_{k \rightarrow \infty} \frac{k^{\beta-1}}{k^\beta \rho(k)} \langle T(x_0, x), \langle f_\beta(\tau), \phi\left(\frac{x_0}{k} + \tau, x\right) \rangle \rangle.$$

From the last equation, using the shift ( $x_0 = kx'_0$ ) we get the following

$$\begin{aligned} &\lim_{k \rightarrow \infty} \frac{1}{\rho(k)} \langle T(kx'_0, x), \langle f_\beta(\tau), \phi(x'_0 + \tau, x) \rangle \rangle \\ &\lim_{k \rightarrow \infty} \frac{1}{\rho(k)} \langle T(kx_0, x), \langle f_\beta(\tau), \phi(x_0 + \tau, x) \rangle \rangle \end{aligned}$$

$\lim_{k \rightarrow \infty} \frac{1}{\rho(k)} \langle T(kx_0, x), \psi(x_0, x) \rangle$ , where function

$\phi(x_0, x) = \psi(x_0, x) = \langle f_\beta(\tau), \phi(x_0 + \tau, x) \rangle$  creates the auto-morphism of space  $S(\overline{\mathbb{R}}_+^1 \times \mathbb{R}^n) \rightarrow S(\overline{\mathbb{R}}_+^1 \times \mathbb{R}^n)$ .

Theorem 5. Let it be that  $m \in \mathbb{N}_0$  and that  $T(x_0, x) \in \mathcal{S}'(\overline{\mathbb{R}}_+^1 \times \mathbb{R}^n)$  has quasi-asymptotics  $g(x_0, x)$  at variable  $x_0$  with respect to auto-modal function  $\rho(k), k \rightarrow \infty$  and let it be that  $x_0^m \in \mathcal{M}_{(\cdot)}$ , with  $\mathcal{M}_{(\cdot)}$  being the space of the multiplier of distributions, then distribution  $x_0^m \cdot T(x_0, x)$  also has quasi-asymptotics  $G(x_0, x) = x_0^m \cdot g(x_0, x)$  at  $x_0$  with respect to auto-modal function  $k^m \rho(k)$ .

Proof. There is

$$\begin{aligned} &\lim_{k \rightarrow \infty} \left\langle \frac{(kx_0)^m \cdot T(kx_0, x)}{k^m \cdot \rho(k)}, \phi(x_0, x) \right\rangle = \\ &= \lim_{k \rightarrow \infty} \left\langle \frac{k^m \cdot x_0^m \cdot T(kx_0, x)}{k^{|m|} \cdot \rho(k)}, \phi(x_0, x) \right\rangle \\ &= \lim_{k \rightarrow \infty} \left\langle \frac{T(kx_0, x)}{\rho(k)}, x_0^m \phi(x_0, x) \right\rangle \\ &= \lim_{k \rightarrow \infty} \left\langle \frac{T(kx_0, x)}{\rho(k)}, x_0^m \phi(x_0, x) \right\rangle = \langle g(x_0, x), x_0^m \phi(x_0, x) \rangle \\ &= \langle x_0^m g(x_0, x), \phi(x_0, x) \rangle = \langle G(x_0, x), \phi(x_0, x) \rangle. \end{aligned}$$

From here we find that  $G(x_0, x) = x_0^m \cdot g(x_0, x)$ .

### 3. Example of the Use of Quasi-Asymptotics to the Solutions of Differential Equations

Let  $L$  be a differential operator with constant coefficients  $a_\beta(x) = a_\beta$  and let  $f \in \mathcal{D}'$ , be such a distribution that convolution  $\mathcal{E} * f$  exists in  $\mathcal{D}'$  where  $\mathcal{E} \in \mathcal{D}'$  is the fundamental solution of equation  $L(D)\mathcal{E} = \delta(x)$ , [3, 6, 9, 11].

Then the solution  $u = \mathcal{E} * f$  of differential equation  $L(D)u = f(x), f \in \mathcal{D}'$  has quasi-asymptotics of order  $\alpha$  with respect to  $\rho(k) = k^\alpha L(k)$  (with  $L(k)$  being the Karamata slow-varying function), if distribution  $f \in \mathcal{D}'$  has such quasi-asymptotics,  $\mathcal{D}'$ -distribution space.

Proof: Let  $f$  have the quasi-asymptotics with respect to  $\rho(k) = k^\alpha L(k)$ . Then the following is valid

$$\frac{1}{\rho(k)} \langle f(kx), \phi(x) \rangle = \frac{1}{k\rho(k)} \langle f(x), \phi\left(\frac{x}{k}\right) \rangle$$

$$\begin{aligned}
 &= \frac{1}{k\rho(k)} \langle \delta(x) * f(x), \varphi\left(\frac{x}{k}\right) \rangle \\
 &= \frac{1}{k\rho(k)} \langle L(D)\mathcal{E} * f(x), \varphi\left(\frac{x}{k}\right) \rangle \\
 &= \frac{1}{k\rho(k)} \langle \left( \sum_{|\alpha|=0}^m a_\alpha D^\alpha \mathcal{E}(x) \right) * f(x), \varphi\left(\frac{x}{k}\right) \rangle \\
 &= \frac{1}{k\rho(k)} \langle \sum_{|\alpha|=0}^m a_\alpha D^\alpha (\mathcal{E} * f)(x), \varphi\left(\frac{x}{k}\right) \rangle \\
 &= \frac{1}{k\rho(k)} \langle L(D)(\mathcal{E} * f)(x), \varphi\left(\frac{x}{k}\right) \rangle \\
 &= \frac{1}{k\rho(k)} \langle L(D)u(x), \varphi\left(\frac{x}{k}\right) \rangle \\
 &= \frac{1}{k\rho(k)} \langle \sum_{|\alpha|=0}^m a_\alpha D^\alpha u(x), \varphi\left(\frac{x}{k}\right) \rangle \\
 &= \frac{1}{k\rho(k)} \sum_{|\alpha|=0}^m \langle D^\alpha u(x), a_\alpha \varphi\left(\frac{x}{k}\right) \rangle \\
 &= \frac{1}{k\rho(k)} \sum_{|\alpha|=0}^m (-1)^{|\alpha|} \langle u(x), D^\alpha \left( a_\alpha \varphi\left(\frac{x}{k}\right) \right) \rangle \\
 &= \frac{1}{k\rho(k)} \langle u(x), L^*(D) \varphi\left(\frac{x}{k}\right) \rangle \\
 &= \frac{1}{k\rho(k)} \langle u(kx), L(-D) \varphi(x) \rangle = \langle \frac{u(kx)}{\rho(k)}, L(-D) \varphi(x) \rangle.
 \end{aligned}$$

Therefore, we have the following:

$\frac{1}{\rho(k)} \langle f(kx), \varphi(x) \rangle = \langle \frac{u(kx)}{\rho(k)}, L(-D) \varphi(x) \rangle$ , and, as per assumption,  $f$  has the quasi-asymptotics, thus, distribution  $u$  has one also.

## 4. Conclusion

Most of the theorems proved in this paper on quasi-asymptotics of distributions at a separable variable have their analog in the case of one-dimensional distributions. In [1], Vladimirov showed a theorem that does not have a one-dimensional analog, the consequence of which is very important, and on the basis of which the application of separated quasi-asymptotics in to the solutions of differential equations.

## References

- [1] V. S. Vladimirov, N. Drozinov, B. I. Zavjalov, *Mnogomernie Tauberovi Teoremi dlja Oboboschenie funkci, Nauka, Moskva, 1986.*
- [2] N. Drozinov, B. I. Zavjalov, *Vvedenie v teoriju oboboschenih funkci, Matematicheskij institut im V. A. Steklov, RAN (MIAN), Moskva. 2006.*
- [3] S. Pilipović, B. Stanković, Vindas, J., *Asymptotic behavior of generalized functions, Series on Analysis, Applications and Computation, 5, World Sci-entific Publishing Co. Pte. Ltd., Hackensack, NJ, 2012.*
- [4] S. Pilipović, J. Toft, *Pseudo-Differential Operators and Generalized Functions, Birkhäuser; 2015.*
- [5] V. S. Vladimirov, *Oboboschenie funkci u matematice skoj fiziki, Nauka, 1979.*
- [6] N. Y. Drozhzhinov, B. I. Zav'yalov, *Asymptotically homogeneous generalized functions and boundary properties of functions holomorphic in tubular cones. Izv. Math., 70, (2006), 1117–1164.*
- [7] S. Pilipović, B. Stanković, *Prostori distribucija, SANU, Novi Sad, 2000.*
- [8] S. Pilipović, *On the Quasiasymptotic of Schwartz distributions. Math. Nachr. 141 (1988), 19-25.*
- [9] J. Vindas, S. Pilipović, *Structural theorems for quasiasymptotics of distributions at the origin, Math. Nachr. 282 (2.11), (2009), 1584–1599.*
- [10] J. Vindas, *The structure of quasiasymptotics of Schwartz distributions, Banach Center Publ. 88 (2010), 297-31.*
- [11] S. Pilipović, *Quasiasymptotic and the translation asymptotic behavior of distributions, Acta Math. Hungarica, 55 (3-4) (1990), 239-243.*
- [12] V. S. Vladimirov, *Uravnjenja matematicheskoy fiziki, Nauka, Moskva, 1981.*
- [13] Teofanov, N., *Convergence of multiresolution expansion in the Schwartz class, Math. Balcanica, 20, (2006), 101-111.*
- [14] S. Pilipović, B. Stanković, *Asymptotic Behavior or Generalized Function, Novi Sad, 2008.*
- [15] N. Stojanović, *Separirana kvaziasimptotika više-dimenzionih distribucija, PMF, Novi Sad, 2009-magistarska teza.*
- [16] M. Tomić, 'Jovan Karamata 1902-1967', *Bulletin T. CXXII de l'Acad'emie Serbe des Sciences et des Arts, No 26, 2001.*